



Effect of Salicylic Acid on Antioxidant Enzymes and Biochemical Contents of Date Palm Plantlets (*Phoenix dactylifera* L.) under Salt Stress Conditions

Naji Salim Jasim and Muntaha Abd-A. Ati

Date Palm Research Center-Basrah University- Basrah, Iraq
E-mail: ahmidnaji916@gmail.com

Abstract: The study was conducted to observe the effect of Salicylic acid (SA) on physiological and biochemical changes in date palm plantlets. Plantlets were irrigated water with different concentrations of salt stress (10, 15 and 20 ds m⁻¹), while the control treatment was irrigated by tap water only. The salt concentration significantly decreased chloroplast pigments, total chlorophyll and carotenoid, total soluble carbohydrates compared with increased levels of proline contents and antioxidant activities (catalase and peroxidase). The treatments of date palm plantlets with salicylic acid (1.5 mM) under salt conditions led to increase of leaf chemical composition and antioxidant enzymes activity, catalase and peroxidase compared with treatments under salt stress only and control.

Keywords: Date palm plantlets, Salicylic acid, Salinity, Antioxidant enzymes, Chlorophyll, Carotenoid

Date palm (*Phoenix dactylifera* L.) is one of the Arecaceae family of economic importance typically grown in arid and semi-arid parts of the globe. Various environmental stresses effect the development and productivity (Shao et al 2008). One of the most influencing factors is salt stress which affects the plant growth and production of secondary metabolites (Nikolova and Ivancheva 2005). One-third of the areas of land in the world's cultivated are seriously affected by high salt stress (Kaya et al 2002). The salts stress of 8-10 ds m⁻¹ decreased chlorophyll contents total soluble, carbohydrates and increased proline and accumulation of Na and Cl in *Cassia absus* L. (Hussain et al 2009). Salts increased protein content and antioxidant enzyme activity CAT and POD on *Momrdica charantia* (Agrawal and Shahee 2007) and on sunflower (Ebrahimian and Bybordi 2012).

Salicylic acid (SA) is an endogenous regulators of growth with a phenolic nature, it contributes in regulating the physiological processes of the plants and has a role as natural inductor of thermogenesis (Horva'th et al 2007). One of important strategies to alleviated plants against different stress is use of plant growth regulators (PGR) which regulating many of physio-biochemical processes like protein synthesis, antioxidant enzymes, and photosynthesis, which are involved in the stress tolerance mechanism directly or indirectly (Akram and Ashraf 2011). Under salt stress 8-16 ds m⁻¹ plant height, leaves numbers, fresh and dry weights of leaves decreased, while treatments of SA 30 mg l and GA3 on Date palm *Phoenix dactylifera* L. increased these estimations and increased proline, superoxide dismutase (POD), peroxidase (POX) and catalase (CAT) (Darwesh and Mohamed 2009, Darwesh 2014). Many trials have shown

that the use of salicylic acid in drought trees enhances the activity of antioxidant enzymes such as POD, SOD and CAT (Hayat et al 2008).

The current study was undertaken to find out whether the application of salicylic acid (SA) on date palm plantlet can alleviate the harmful effects of salinity on chemical contents and antioxidant enzymes CAT and POX.

MATERIAL AND METHODS

This study was conducted in greenhouse and laboratories of Date Palm Research Center, the University of Basra during 2017 to 2018. Three levels of salinity stress 10, 15 and 20 ds m⁻¹ alone and application of while salicylic acid at 1.5 mM were evaluated along with control (0 ds m⁻¹ salt).

Plantlets of the date palm were ten months old. The different versions mentioned above were applied to the plants for 21 days depending on the soil water content of the study soil. Each treatment consists of three replicates (2plantlets pot⁻¹). The following physiological characteristics measurements were recorded after the completion of the experiment.

Total chlorophyll and total carotenoid: The total chlorophyll was estimated by method given by Aron (1949) from fresh leaves were determined by U.V. spectrophotometer (CECL 2021, ENGLAND) at wave max 663nm, The total Carotenoid contents from fresh leaves were determined by U.V. spectrophotometer (CECL 2021, ENGLAND) at wave max 470nm (Lichtenthaler and Wellburn 1983).

Proline content: Proline was estimated by Bates et al (1973) method the standard curve (mg g⁻¹ DW).

Determination of soluble carbohydrate content: Fales (1951) and Schlegel (1956) method was used for estimation of soluble carbohydrate content in aqueous solution

Measurement of catalase (CAT) activity: The assessment of the activity of the catalase (EC: 1.11.1.6) was according to the method given by Chance and Maehly (1955).

Peroxidase (POX) activity: MacAdam et al (1992) procedure was used to assess Peroxidase (EC: 1.11.1.7) activity.

RESULTS AND DISCUSSIONS

Photosynthetic pigments: Results of photosynthetic pigments (Fig. 1 and 2) indicated significant reduction in total chlorophyll and carotenoid contents under salt stress at 15 and 20 ds m⁻¹ which were 9.24 and 7.38 mg 100 g⁻¹ FW for chlorophyll respectively and 0.029 and 0.0215 mg 100g⁻¹ FW for carotenoid respectively. This reduction of pigments contents could be related to the toxic effects of salinity on the pigments biosynthesis elevating their degradation and/or causing damage to the chloroplast thylakoid. Darwesh et al (2006) observed that salinity from 10000-14000 ppm decreased chlorophyll contents in leaves on date palm. Furthermore, Abd.El-Samed et al (2011) found that chlorophyll contents decreased under 30-90 mM NaCl on *Zea mays* and broad bean *Vicia faba*. This inhibition of photosynthesis pigments was recovered by the addition of 1.5mM salicylic acid. Exogenous application of SA at 1.5 mM combination with salt concentration 10, 15 and 20 ds m⁻¹ caused an increased in total chlorophyll and carotenoid which were 11.721, 13.238 and 9.762 mg 100 g⁻¹ FW for chlorophyll and 0.037, 0.039 and 0.033 mg 100 g⁻¹ FW, respectively. Al-Mayahi (2015) observed that the chlorophyll content reduction in date palm leaves under salt stress. A decline in the level of photosynthesis pigments were due to the formation of protolytic enzymes such as chlorophyllase which is responsible of destruction of chlorophyll under high stress condition. Schutz and Fangmier (2001) observed that the reduction of chlorophyll owing to stress is due to the increase in reactive oxygen species (ROS) production in the cell. SA application causes ROS scavenging and that can enhance chlorophyll biosynthesis and prevent degradation of chlorophyll and enhanced content in date palm plantlets.

Proline content: Both salt stress and salicylic acid affected significantly proline content. In addition, the interaction effect of both salt and salicylic acid on proline content of plantlets was significant. The salt increased greatly proline content in treated plants. Proline content of the plantlets under salt stresses concentrations 10, 15 and 20 ds.m⁻¹ were 18.41, 29.24 and 27.49 µg g⁻¹ respectively in comparison with control, which was 17.67 µg g⁻¹ FW (Fig. 3). The application

of SA in combination with salt concentrations increased proline content which were 37.49, 40.66 and 34.24 µg g⁻¹. Dash and Panda (2001) reported that when NaCl concentration and duration of stress increased proline content in *Phaseolus mungo*. Other study found that NaCl at 50-150 mM increased proline contents in *Salventia natans* L. (Jampeetong and Brix 2009). The application of SA caused enhancement in the synthesis of proline which might be the adaptive mechanism of date palm plantlets under salt stress condition. These results are in agreement with finding of previous studies (El-Beltagi et al 2017, Agamy et al 2013). The increase in proline content in plants allowed water stressed plants to maintain low water potential, the accumulation of proline involved in osmo-regulation seemed to enable additional water to be absorbed from the environment (Gebaly et al 2013).

Soluble carbohydrate contents: The applying 1.5mM SA on the control plants elevated the content of leaf soluble carbohydrate which was 23.99 mg g⁻¹ DW compared to control plant (without SA) which was 15.62 mg g⁻¹ DW. With salt stress, the soluble carbohydrate content of plantlet leaves was substantially high with increased salt concentrations (Fig. 4). Exposure plantlets were irrigated with growth regulator (SA) at 1.5 mM in combination with salt stress significant more in leaves content of total carbohydrates in plants which were 26.368, 23.226 and 22.307 mg g⁻¹ DW for 10, 15 and 20 ds m⁻¹ respectively compared with treated plants at the same concentrations which were 22.153, 22.920 and 19.088 mg g⁻¹ respectively. Salt stresses, however, resulted in a significant increase in the content of the soluble carbohydrate of date palm leaves. With increasing salt concentrations the rate of rise in soluble carbohydrate content has been reduced, demonstrating a function for soluble carbohydrate in osmotic adjustment. Pe'rez-Lo'pez et al (2010) indicated that osmotic adjustment could involve the accumulation of soluble carbohydrates in crops under stress circumstances. The accumulation of soluble carbohydrates was commonly recorded as a reaction to salinity or drought (Cheeseman 1988). The carbohydrates constitute one of the dry matter's primary organic constituents were influenced by salt stress. The complete carbohydrate content was decreased by growing salinity concentrations. Reducing the content of carbohydrates may be due to reduced photosynthesis and increased photorespiration under salt stress (Namich et al 2007). The application of SA to date palm plantlets under salt conditions increased the contents of total soluble carbohydrates as compared with untreated stressed plants. Photosynthetic pigments were increased in response to salicylic acid treatments thus enhance the polysaccharides biosynthesis

and total carbohydrates significantly which are utilized in the sunflower plants growth (Dawood et al 2012). The application of SA might activate the metabolic consumption of soluble carbohydrates to form new cell constituents as a mechanism to alleviated the growth of date palm plantlets. The high concentration of carbohydrates with its role in reducing water content helps prevent oxidative damages and the

preservation of protein structure during water shortage. Carbohydrates also play a molecular role in sugar-responsible genes, which give different physiological responses such as defensive responses and cellular expansion (Simaei et al 2011). Moreover, accumulation of carbohydrate play a key role in alleviating the salinity stress, either via osmotic adjustment, as Ackerson (1985).




Fig. 1. SA effects on the total chlorophyll content (mg 100 g⁻¹) under salt condition




Fig. 4. SA effect on the carbohydrate content (mg g⁻¹) under salt condition




Fig. 2. SA effects on the Carotenoid content (mg 100g⁻¹) under salt condition




Fig. 5. SA effect on the catalase (unit g⁻¹) activity under salt condition




Fig. 3. SA effect on the proline content (µg g⁻¹) under salt condition




Fig. 6. SAeffect on the peroxidase (unit g⁻¹) activity under salt condition

Catalase (CAT) and Peroxidase (POD): The CAT and POD activity increased during salt stress and this increase was positively related to 10 and 15 ds.m⁻¹ salt concentrations. Additionally, the results showed that high levels of antioxidant enzymes CAT and POD activity occurred in plants at 10 and 15 ds m⁻¹ salt concentrations with SA combination at 1.5mM. Control crops revealed the smallest concentrations of CAT and POD operations. The enzymes of antioxidants include catalase (CAT), superoxide dismutase (SOD), peroxidase (POX) (Zhu 2001, Ashraf 2002). The study of Mittler (2002), showed that salinity induces and increases the activity of these antioxidant enzymes. CAT involving degradation of hydrogen peroxide and prevention of oxidative damage. The resistance to environmental stress may therefore depend at least partially on the production by enhancing the antioxidant defense system (Azevedo et al 2006). CAT and POD markedly increased under salt stress 40-200 mM (Abdulwahid 2012 on date palm). Increasing CAT and POD activities with SA application at 1.5mM was the highest under salt pressure, exogenous application of SA can control the activity of intracellular antioxidant enzymes such as POX, SOD and boost plant sensitivity to environmental stress (Sakhabutdinova et al 2004). Study of Saruhan et al (2012) discovered that the application of SA increased the activity of antioxidant enzymes such as CAT, POD and APX in plants.

CONCLUSIONS

Salt stress was significantly reduced chemical compositions chloroplast pigments (total chlorophyll and carotenoid), total soluble carbohydrate, antioxidant activity (CAT and POD) and increased proline content. The adverse effect of salinity can be alleviated by applying SA, thereby raising all above physiological characteristics.

REFERENCES

- Abd El-Samad HM, Shaddad MAK and Barakat N 2011. Improvement of plants salt tolerance by exogenous application of amino acids. *Journal of Medicinal Plants Research* **5**(24): 5692-5699.
- Abdulwahid AH 2012. Investigation of the effect of salt stress on the antioxidant enzyme activities on leaves of Date Palm (*Phoenix dactylifera* L.) seedling. *Advances in Agriculture and Botany-International Journal of Bioflux Society* **4**(3): 94-102.
- Ackerson RC 1985. Osmoregulation of cotton in response to water stress III. Effects of phosphorus fertility. *Journal of Plant Physiology* **77**: 309-312.
- Agamy RA, Hafez EE and Taha TH 2013. Acquired resistant motivated by salicylic acid applications on salt stressed tomato (*Lycopersicon esculentum* Mill.). *American-Eurasian Journal of Agricultural and Environmental Sciences* **13**(1): 50-57.
- Agarwal S and Shaheen R 2007. Stimulation of antioxidant system and lipid peroxidation by abiotic stresses in leaves of *Momordica charantia*. *Brazilian Journal of Plant Physiology* **19**(2): 149-161.
- Akram NA and Ashraf M 2011. Pattern of accumulation of inorganic elements in sunflower (*Helianthus annuus* L.) plants subjected to salt stress and exogenous application of 5- amino lenuinic acid, *Pakistan Journal of Botany* **43**(1): 521-530.
- Al-Mayahi AM 2015. The compound effects of NaCl and gibberellic acid on the growth and development of in vitro date palm (*Phoenix dactylifera* L.) plantlets cv. Showathy. *Journal of King Saud, University of Agriculture Science* **26**(2): 1-33.
- Arnon DI 1949. Copper enzymes in isolated chloroplast. Polyphenyl oxidase in *Beta vulgaris*. *Journal Plant Physiology* **24**: 1-5.
- Ashraf M, Karim F and Rasul E 2002. Interactive effects of gibberellic acid (GA3) and salt stress on growth, ion accumulation and photosynthetic capacity of two spring wheat (*Triticum aestivum* L.) cultivars differing in salt tolerance. *Plant Growth Regulation* **36**: 49-59.
- Azevedo N, Prisco AD, Eneas-Filho JT, De Abreu CEB and Gomes-Filho E 2006. Effect of salt stress on antioxidative enzymes and lipid peroxidation in leaves and roots of salt- tolerant and salt-sensitive maize genotypes. *Environmental and Experiment Botany* **56**: 87-94.
- Bates LS, Waldern RP and Teare ID 1973. Rapid determination of free proline for water stress studies. *Plant Soil* **39**: 205-207.
- Chance B and Maehly AC 1955. Assay of Catalases and Peroxidases, In: *Methods in Enzymology*, (Eds.): Colowick SP and Kaplan NO. Academic Press. 380 P.
- Cheeseman JM 1988. Mechanisms of salinity tolerance in plants. *Plant Physiology* **117**: 547-550.
- Darwesh RSS and Mohamed HF 2009. The adverse effect of GA3 (gibberellic acid) on salinity of *Phoenix dactylifera* L. plantlets in vitro rooting stage. 4 *Conference on Recent Technologies in Agriculture*, pp: 663-669.
- Darwesh RSS, El-Tantawy AA, Arafa AMS and El-Banna AE 2006. *Effect of salts stress on growth and development in vitro culture, acclimatization Stage on Phoenix dactylifera L. and Arecastrum romanzoffianum Becc. seedlings in greenhouse*. Ph.D. Thesis, Faculty of Agriculture, Cairo University, Egypt, pp: 55-65.
- Darwesh RSS 2014. Exogenous supply of salicylic acid and IAA on morphology and biochemical characteristics of date plantlets exposed to salt stress. *Middle East Journal of Agriculture Research* **3**(3): 549-559.
- Dash M and Panda SK 2001. Salt stress induced changes in growth and enzyme activities in germinating *Phaseolus mungo* seeds. *Biologia Plantarum* **44**: 587-589.
- Dawood MG, Sadak MS and Hozayen M 2012. Physiological role of salicylic acid in improving performance, yield and some biochemical aspects of sunflower plant grown under newly reclaimed sandy soil. *Australian Journal of Basic and Applied Sciences* **6**(4): 82-89.
- Ebrahimian E and Bybordi A 2012. Effect of salinity, salicylic acid, silicium and ascorbic acid on lipid peroxidation, antioxidant enzyme activity and fatty acid content of sunflower. *African Journal of Agriculture Research* **7**(25): 3685-3694.
- El-Beltagi HS, Ahmed SH, Namich AAM and Abdel-Sattar RR 2017. Effect of salicylic acid and potassium citrate on cotton plant under salt stress. *Fresenius Environmental Bulletin* **26**(1): 1091-1100.
- Fales FW 1951. The assimilation and degradation of carbohydrates by yeast cells. *Journal of Biological and Chemical* **193**: 113-124.
- Gebaly SG, Ahmed FMM and Namich AAM 2013. Effect of spraying some organic, amino acids and potassium citrate on alleviation of drought stress in cotton plant. *Journal of plant production, Mansoura University* **4**(9): 1369-1381.
- Hayat S, Hasan SA, Fariduddin Q and Ahmad A 2008. Growth of tomato (*Lycopersicon esculentum*) in response to salicylic acid under water stress. *Journal of Plant Interaction* **3**: 297-304.
- Horva'th E, Szalai G and Janda T 2007. Induction of abiotic stress tolerance by salicylic acid signaling. *Journal of Plant Growth Regulation* **26**: 290-300.

- Hussain K, Majeed M and Nisar FF 2009. Growth and ionic adjustments of chaksu (*Cassia absus* L.) under NaCl stress. *American-Eurasian Journal of Agriculture and Environmental Sciences* **6**(5): 557-560.
- Jampeetong A and Brix H 2009. Effects of NaCl salinity on growth, morphology, photosynthesis and proline accumulation of *Salvinia natans*. *Aquatic Botany* **91**(3): 181-186.
- Kaya C, Kirnak H, Higgs D and Saltati K 2002. Supplementary calcium enhances plant growth and fruit yield in strawberry cultivars grown at high (NaCl) salinity. *Science of Horticultural* **26**: 807-820.
- Lichtenthaler HK and Wellburn WR 1983. Determination of total carotenoids and chlorophyll a and b of leaf extracts in different solvents. *Biochemical Society Transactions* **11**: 591-592.
- MacAdam JW, Nelson CJ and Sharp RE 1992. Peroxidase activity in the leaf elongation zone of tall fescue. *Plant Physiology* **99**: 872-878.
- Mittler R 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sciences* **7**: 405-410.
- Namich AAM, Kassem MM and Gebaly SG 2007. Effect of irrigation with saline water on some cotton cultivars. *Journal of Agriculture Sciences, Mansoura University* **32**(7): 5117-5136.
- Nikolova MT and Ivancheva SV 2005. Quantitative flavonoid variations of *Artemisia vulgaris* L. and *Veronica chamaedrys* L. In relation to altitude and polluted environment. *Acta Biologica Szegediensis* **49**:29-32.
- Pe´rez-Lo´pez U, Robredo A, Lacuesta M, Mu˜oz-Rueda A and Mena-Petite A 2010. Atmospheric CO₂ concentration influences the contributions of osmolytes accumulation and cell wall elasticity to salt tolerance in barley cultivars. *Journal of Plant Physiology* **167**: 15-22.
- Sakhabutdinova AR, Fatkhudinova DR and Shakirova FM 2004. Effect of salicylic acid on the activity of antioxidant enzymes in wheat under conditions of salination. *Applied Biochemical and Microbiology* **40**: 501-505.
- Saruhan N, Saglam A and Kadioglu A 2012. Salicylic acid pretreatment induces drought tolerance and delays leaf rolling by inducing antioxidant systems in maize genotypes. *Acta Physiologia Plant* **34**: 97-106.
- Schlegel HG 1956. Die verwertung organischer sauren durch Chlorella in lichen. *Planta* **47**: 510-514.
- Schutz M and Fangmeir E 2001. Growth and yield response of spring wheat (*Triticum aestivum* L. cv. Minaret) to elevated CO₂ and water limitation. *Environmental Pollution* **11**:187-194.
- Shao HB, Chu LY and Jaleel CA 2008. Water-deficit stress induced anatomical changes in higher plants. *CR Biologies* **331**(3): 215-225.
- Simaei M, Khavari-Nejad RA, Saadatmand S, Bernard F and Fahimi F 2011. Interactive effects of Salicylic acid and nitric oxide on soybean plants under NaCl salinity. *Russian Journal of Plant Physiology* **58**(5): 783-390.