

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/8132005>

Thiosugar Nucleosides. Synthesis and Biological Activity of 1,3,4-Thiadiazole, Thiazoline and Thiourea Derivatives of 5-Thio-D-Glucose.

Article in *Nucleosides Nucleotides & Nucleic Acids* · February 2004

DOI: 10.1081/NCN-200034040 · Source: PubMed

CITATIONS

16

READS

156

3 authors:



Najim Al-Masoudi

University of Basrah & University of Konstanz (Germany)

270 PUBLICATIONS 2,378 CITATIONS

[SEE PROFILE](#)



Yaseen A. Al-Soud

Al al-Bayt University

123 PUBLICATIONS 1,653 CITATIONS

[SEE PROFILE](#)



Wasfi A. Al-Masoudi

University of Basrah

55 PUBLICATIONS 101 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



5-Nitroimidazole: Synthetic Approaches and Pharmacological Importance [View project](#)



Let's Save the World and Humanity [View project](#)

Thiosugar Nucleosides. Synthesis and Biological Activity of 1,3,4-Thiadiazole, Thiazoline and Thiourea Derivatives of 5-Thio-D-Glucose

Najim A. Al-Masoudi,^{1,*} Yaseen A. Al-Soud,² and Wasfi A. Al-Masoudi³

¹Department of Chemistry, University of Konstanz, Konstanz, Germany

²Department of Chemistry, College of Science,
University of Al al-Bayt, Al-Mafraq, Jordan

³Department of Chemistry, College of Arts and Science,
Almergab University, Tarhona, Libya

ABSTRACT

New acylated 5-thio- β -D-glucopyranosylimino-disubstituted 1,3,4-thiadiazols **8**, and **11** were prepared, via spontaneous rearrangements, by cycloaddition of the glycosyl isothiocyanate **2** with the reactive intermediates 1-aza-2-azoniaallene hexachloroantimonates **4** and **6**, respectively. Reaction of **2** with aminoacetone or chloroethylamine afforded the acylated 5-thio- β -D-glucopyranosyl-4-imidazoline-2-thione nucleoside **16** and glucopyranosylamino-2-thiazoline derivative **18**, respectively. Deblocking of **8**, **11**, **17** and **19** furnished the free nucleoside analogues **9**, **12**, **18** and **20**, respectively. Analogously, treatment of **2** with chloroethylamine in the 1:2 ratio afforded the thioureyldisaccharide **21**. No in vitro antiviral activity against HIV-1, HIV-2, human cytomegalovirus (HCMV), has been found for the new synthesized compounds.

Key Words: Antitumor activity; Cumulenes; Glycosylimino-1,3,4-thiadiazoles; Glycosyl isothiocyanates; Glycosylthiourea.

*Correspondence: Najim A. Al-Masoudi, EuroMed, P.O. Box 10 05 52, Konstanz D-78405, Germany; Fax: +49-(0)7531-34435; E-mail: Najim.Al-Masoudi@gmx.de.

INTRODUCTION

Because isothiocyanates^[1] and glycosylisothiocyanate^[2–4] are versatile synthetic synthons in organic chemistry, their stereoselective syntheses of different heterocyclic derivatives, via nucleophilic and cycloadditions, such as *N*-nucleosides,^[1,5–7] glycosylamino-heterocycles, glycosyl-thioureas,^[8–11] glycosyl-guanidines and glycosyl derivatives of β -cyclodextrines^[12–17] with potential pharmacological properties,^[18] are of continuing interest. On the other hand, 5-thio-D-glucose is an important synthon in carbohydrate chemistry since exhibited interesting biological activities, such as inhibition of D-glucose transport across membranes (inhibition of spermatogenesis) and of carbohydrate-metabolising enzymes.^[19–21]


Recently, we reported syntheses of 2,5- and 2,3-dihydro-(glycosylimino)-1,3,4-thiadiazole and 3'-(disubstituted 1,3,4-thiadiazolimino)-2',3'-dideoxythymidines formed by reaction of a glycosyl and 3'-thymidine isothiocyanate, respectively, with certain cumulene salts.^[15,6] We now describe the results of a more investigation of reactive cumulenes, aminoacetone or chloroethylamine with acylated 5-thio-D-glucose isothiocyanate.

RESULTS AND DISCUSSION

In our recent work, some reactive cumulenes have been used in the synthesis of new types of glycosyl 1,2,4-triazole *C*-nucleosides,^[6,22–24] and glycosylimino-1,3,4-thiadiazole nucleosides^[15,6] from cycloaddition with the glycosyl nitrile and isothiocyanate, respectively. The starting material 2,3,4,6-tetra-*O*-acetyl-5-thio- β -D-glucopyranosyl isothiocyanate (**2**) was prepared, in 39% yield, by treatment of the corresponding bromo derivative **1**^[25] with trimethylsilyl isothiocyanate in the presence of SnCl₄ as a Lewis acid. The heteroallene **4** carrying a *tert*-butyl group were used to synthesize electrically neutral 2-substituted 1,2,4-triazoles.^[26,27] Compound **4**, which formed at low temperature from **3**^[6,24] and SbCl₅, reacted with isothiocyanate **2** via C = S cycloaddition to give the nitrilium ion **7a** as an intermediate. The latter is cyclized to the thiadiazolium salt **7b** giving, after the loss of the *tert*-butyl group (as isobutene), the inseparable iminium salt **7c**. The *tert*-butyl group was probably lost during or after the rearrangement. In situ deprotection of **7c** with saturated aqueous NaHCO₃ afforded thiadiazole **8** (67%). Deblocking of **8** with NaOMe in MeOH afforded after purification the free glycosylimino derivative **9** (82%). The *concerted* cycloaddition to isothiocyanate could have been occurred across the C = S bond of **2** to give **8** [*cf.* structure **14**, δ_C : C = N (152.0); N = C–S (149.7)]^[28,29] and not the C = N bond, which furnishes an isomeric 4,5-dihydro-1*H*-1,3,4-triazole-5-thione (*cf.* structure **13**). The ¹³C chemical shift of **8** spectral data indicated the **2** reacts as *S*-nucleophile and not as *N*-nucleophile, from the ¹³C NMR spectrum data in comparison to those of **14**. Furthermore, L'abbé et al. reported values in the range of δ_C 151–164 ppm for C-2 and of δ_C 149–155 for C-5 for several thiadiazoles of type **14**,^[30] which are in agreement with data shown by compound **8** (δ_C : 158.0 for C-5 and 147.2 for C-1).

Analogously, the α,α -dichloroazo compound **5**^[31] was obtained by chlorination of the bis-hydrazone, and the 1-aza-2-azaniaallene salt **6**^[32–34] by treatment of **5** at –60°C with SbCl₅. Addition of **2** to the reactive intermediate **6**, by employing the methods of Jochims et al.,^[16] lead to a color change of the orange suspension

of **6** between -60°C and $+23^{\circ}\text{C}$ indicating a cycloaddition reaction. The resulting 1,3,4-thiadiazolium hexachloroantimonate **10a** lost its CMe_2Cl group and furnished the protonated salt **10b**. Deprotection afforded 1,3,4-thiadiazole **11** (41%) (Scheme 1). Deblocking of **11** with NaOMe in MeOH afforded **12** in 83% yield. Again, the concerted cycloaddition of **2** with the reactive cumulene **6** would occurred via $\text{C}=\text{S}$ double bond, as shown from the ^{13}C NMR spectrum [δ_{C} : C-5 (106.9); C-2 (176.8)] [cf. $\text{C}_5\text{-Me}_2$ (106.9); $\text{C}=\text{N}$ (174.3)] and not across the $\text{C}=\text{N}$ double bond to give an



Scheme 1. Reagent and conditions: (i) SbCl_5 , CH_2Cl_2 , -60°C to 23°C , 7 h, CH_2Cl_2 , 23°C ; (ii) aq. NaHCO_3 , 2 h, 23°C , (iii) NaOMe, MeOH, 16 h, 23°C .

isomeric mixture of 4,5-dihydro-1*H*-1,3,4-triazole-5-thione (*cf.* structure of **15**], while for several triazoles **16** C = S shifts close to δ_C 188 and shifts of the saturated ring carbon (C-5) of δ_C 109–110 were found.^[28] Accordingly compound **11** is a thia-diazole. The large coupling constants ($J_{1',2'}$ and $J_{4',5'} \sim 9$ –10 Hz) is indicating of the β -configuration, and the 4C_1 conformation of the sugar moiety.


Next, our attempt to synthesize the *N*-nucleoside 5-methyl-1-(2,3,4,6-tetra-*O*-acetyl-5-thio- β -D-glucopyranosyl)-4-imidazoline-2-thione (**17**), in 51% yield, by cycloaddition reaction of the corresponding glycosyl isothiocyanate **2** with aminoacetone hydrochloride in acetone at 23°C. Deblocking of **17** afforded the free nucleoside **18** (86%). The structures of **17** and **18** were identified by their 1H NMR, ^{13}C NMR, and mass spectrum. The 1H NMR spectrum is characterized by the presence of H-1' and H-4' as doublet and triplet at δ_H 5.80 ($J_{1',2'} = 9.3$ Hz), and δ_H 5.40 ($J_{4',5'} = 10.0$ Hz), indicating for the β -configuration and 4C_1 conformation of the sugar moiety. The rotating nuclear Overhauser effect (ROE)^[35–37] between H-1' and H-5' is an additional proof for β -configuration. The two singlets at δ_H 2.36 and δ_H 10.69 were attributed to the methyl group at C-5 and NH group of the imidazoline ring. The ribosylation of **17** occurred at the *N*-site of the imidazoline ring, and this was proved by the selected homo- and heteronuclear NMR study. Thus, gradient selected HMBC^[38] spectrum allowed via $^2J_{C,H}$ and $^3J_{C,H}$ couplings the assignment of all the carbon atoms. C = S and C-5 groups at δ_C 163.0 and δ_C 34.5, respectively, were identified from their $^2J_{C,H}$ correlations to H-1' at δ_H 5.80. C-4 at δ_C 127.0 was identified from its $^3J_{C,H}$ correlation with H-1' as well. Furthermore, these data are in agreement with those of the related nucleosides.^[7]

Sainsbury^[39] had reported that reaction of β -haloalkylamines with alkyl(aryl)-isothiocyanate furnished 2-alkyl(alkyl)aminothiazolines. Therefore, reaction of **2** with an excess of chloroethylamine for 12 h at 23°C, followed by treatment with aqueous solution of $NaHCO_3$ gave after chromatographic purification an a foam, tentatively identified as 2-(2,3,4,6-tetra-*O*-acetyl-5-thio- β -D-glucopyranosyl)amino-2-thiazoline (**19**) (82%). Deblocking of **19** with NaOMe in MeOH afforded the free aminoglycoside (**20**) (82%). The structures of **19** and **20** were assigned from their 1H NMR, which showed similar pattern, and mass spectra. H-1' and H-4' of **19** were appeared as doublet of doublets at δ_H 4.97 ($J_{1',2'} = 8.8$ Hz, $J_{NH,H-1'} = 5.5$ Hz) and δ_H 5.30 ($J_{4',5'} = 9.2$ Hz), respectively, giving an proof for the β -configuration and 4C_1 -conformation of the sugar moiety. The doublet at δ_H 8.20 were attributed to C_1 -NH, while the triplets at δ_H 3.60 and 3.20 ($J = 7.2$ Hz) were assigned to CH_2 -5 and CH_2 -4, respectively. These data are in agreement with those of the related oxygen analogue prepared previously.^[7]

Under the above condition, by using chloroethylamine and **2** in the 1:2 ratio the thioureylenedisaccharide **21** was obtained (71%). The structure of **21** was based on analytical UV and 1H NMR data. The shift at lower wave number of NH ($\Delta\nu = 252$ cm^{-1}) and C = N ($\Delta\nu = 20$ –40 cm^{-1}), with respect to other non-bonded thiazolines, 2-amino-2-thiazolines and 2-amino-2-oxazolines,^[40] indicates a strong intramolecular hydrogen bond is present in compound **21**. Furthermore, these data is coincident with those reported for *N*-(2,3,4,6-tetra-*O*-acetyl-5-thio- β -D-glucopyranosyl)-*N'*-(2-thiazolin-2-yl)thiourea^[41] (Scheme 2).

Biological Activity

Compounds **9**, **12**, **20** and **21** were evaluated for their anti-HIV activity in vitro using the IIBB strain for HIV-1 and the ROD strain for HIV-2, and monitored by the



Scheme 2. Reagents and conditions: (i) $\text{NH}_2\text{CH}_2\text{COMe.HCl}/\text{NaHCO}_3$, Me_2CO , 30 min, 23°C ; (ii) $\text{ClC}_2\text{H}_4\text{NH}_2.\text{HCl}/\text{H}_2\text{O}/\text{ether}$, M NaOH , 15 h, 23°C .

inhibition of the virus-induced cytopathic effect in human MT-4 lymphocyte cells at non-toxic concentrations, as well as for the human anti-cytomegalovirus (HCMV, causing pneumonia and CNS diseases) in human embryonic lung (HEL) cells with AD-169 and Davis strain and were found to be inactive in comparison to the anti-HIV activity of Delviridine.^[42] In conclusion, the 4-thio-glycofuranosyl moiety carrying the similar heterocycles might optimize the antiviral activity and then could balance both the hydrophobic and hydrophilic requirements of our target molecules.

Experimental

General Procedure

See Refs. [6,22–24].

2,3,4,6-Tetra-*O*-acetyl-5-thio- β -D-glucopyranosyl isothiocyanate (2). A solution of **1**^[25] (4.27 g, 10 mmol), trimethylsilyl isothiocyanate (1.31 g, 10 mmol) and distilled SnCl_4 (2.61 g, 10 mmol) in CH_2Cl_2 (60 mL) was boiled under reflux for 13 h. After addition of further trimethylsilyl isothiocyanate (0.66 g, 5.0 mmol) and SnCl_4 (2.61 g, 10 mmol), the mixture was boiled for another 12 h. After cooling, the mixture was neutralized by shaking with water (50 mL) and excess of NaHCO_3 , followed by filtration and the organic layer was separated. The aqueous layer was extracted with CH_2Cl_2 (2×30 mL) and the combined organic extracts were dried (Na_2SO_4), treated with charcoal, filtered (*celite*) and evaporated to dryness to give an anomeric mixture of the glycosyl isothiocyanate **2** as a dark orange syrup. The crude product was poured onto a column of silica gel (40 g) using toluene-ethyl acetate (3:1) as eluent. Eluted first was a syrup, as an unidentified product, while further elution yielded a syrup which crystallized at 5°C . Recrystallization twice from EtOH yielded the pure β -isomer (1.58 g, 39%); m.p. $96\text{--}99^\circ\text{C}$; $[\alpha]_{\text{D}} + 6^\circ$ (c 1.2, CHCl_3); ν_{max} (CCl_4) 2019 cm^{-1} (NCS); δ_{H} (CDCl_3): 5.18 (dd, 1H, $J_{4,5} = 10.5$ Hz, H-4); 5.10 (t, 1H, H-2); 4.98 (dd, 1H, $J_{3,4} = 9.5$ Hz, H-3); 4.42 (d, 1H, $J_{1,2} = 10.6$ Hz, H-1); 4.22 (dd, 1H, $J_{6,6'} = 12.0$ Hz,

H-6'); 4.07 (dd, 1H, Hz, H-6); 3.25 (ddd, 1H, $J_{5,6} = 5.5$ Hz, $J_{5,6'} = 3.4$ Hz, H-5); 2.09, 2.00, 1.96, 1.87 (4xs, 12H, 4xOAc). Anal Calc. for $C_{15}H_{19}N_2S_2O_8$ (405.4): C, 44.44; H, 4.72; N., 3.45. Found: C, 44.18; H, 4.61; N., 3.19. MS: m/z (FAB) 406 (M + H)⁺.

General procedure for the preparation of the acylated 5-thio- β -D-glucopyranosylimino-disubstituted 1,3,4-thiadiazols 8, and 11. A solution of $SbCl_5$ (0.90 g, 3.0 mmol) was added dropwise to a stirred, cooled ($-60^\circ C$) solution of **2** (0.41 g, 1.0 mmol) and **3** or **5** (3.0 mmol) in CH_2Cl_2 (15 mL). After stirring the mixture at $-60^\circ C$ for 1 h, then at $0^\circ C$ for 1 h, and finally at $23^\circ C$ for 10 min, the solvent was evaporated to dryness. The brown residue was dissolved in MeCN (15 mL). After cooling the mixture to $0^\circ C$, an aqueous solution of $NaHCO_3$ [2.52 g (30 mmol) in H_2O (10 mL)] was added slowly. The mixture was stirred at $23^\circ C$ for 30 min and filtered. The organic solvent was evaporated and the aqueous layer was extracted with CH_2Cl_2 (3×30 mL). The combined organic extracts were washed with water (3×10 mL), dried (Na_2SO_4), filtered, and evaporated to dryness. The amorphous residue was purified on a column of SiO_2 (20 g) using first CH_2Cl_2 and then CH_2Cl_2 -MeOH (99:1) as eluents.

2,3-Dihydro-5-methyl-2-(2,3,4,6-tetra-*O*-acetyl-5-thio- β -D-glucopyranosylimino)-3-(2,4,6-trichlorophenyl)-1,3,4-thiadiazole (8). From **3** (0.98 g). Yield: 0.43 g, (67%). δ_H ($CDCl_3$): 7.41 (dd, $J = 2.2$ Hz, ArH); 5.43 (t, 1H, $J_{4',5'} = 9.0$ Hz, H-4'); 5.25 (t, 1H, $J_{2',3'} = 9.0$ Hz, H-2'); 5.17 (t, 1H, $J_{3',4'} = 9.0$ Hz, H-3'); 4.40 (d, 1H, $J_{1',2'} = 8.9$ Hz, H-1'); 4.26 (dd, 1H, $J_{6',6''} = 11.5$ Hz, H-6''); 4.10 (dd, 1H, $J_{5',6'} = 4.9$ Hz, H-6'); 3.47 (ddd, 1H, $J_{5',6''} = 3.5$ Hz, H-5'); 2.42 (s, 3H, C_5 -Me); 2.03, 1.99, 1.96, 1.88 (4xs, 12H, 4xOAc). δ_C ($CDCl_3$): 170.4, 170.1, 169.1, 168.9 (C = O); 158.0 (C_5 = N); 147.2 (C_1 = N); 136.4, 136.0, 135.7, 133.0, 128.8, 128.5 (Ar-C); 74.2 (C-2'); 73.6 (C-3'); 72.0 (C-4'); 64.5 [$^1J(^{13}C$ - 1H) = 152.0 Hz, (C-1')]; 61.4 (C-6'); 42.2 (C-5'); 20.6–20.3 (4xCOMe). Anal Calc. for $C_{23}H_{24}Cl_3N_3S_2O_8$ (640.9): C, 43.10; H, 3.77; N., 6.56. Found: C, 42.78; H, 3.62; N., 6.32. MS: m/z (FAB) 663/665 (M + Na)⁺.

2,3-Dihydro-5-methyl-2-(5-thio- β -D-glucopyranosylimino)-3-(2,4,6-trichlorophenyl)-1,3,4-thiadiazole (9). A solution of **8** (0.30 g, 0.47 mmol) in 0.3 M NaOMe solution (10 mL) was stirred at $23^\circ C$ for 9 h. The solution was neutralized with AcOH to pH 5 and then evaporated to dryness. The residue was partitioned between water (10 mL) and ether (10 mL) and the aqueous layer was evaporated to dryness and the residue was co-evaporated with EtOH (4×10 mL). The residue was purified by column chromatography on (5 g) using, in gradient, MeOH (0–10%) and CH_2Cl_2 to give **9** (0.18 g, 82%), m.p. 149 – $152^\circ C$. δ_H ($DMSO-d_6/D_2O$): 7.84 (dd, $J = 2.2$ Hz, ArH); 3.97 (d, 1H, $J_{1',2'} = 9.6$ Hz, H-1'); 3.89 (dd, 1H, $J_{6',6''} = 12.0$ Hz, H-6''); 3.80 (dd, 1H, $J_{5',6'} = 3.5$ Hz, H-6'); 3.54 (dd, 1H, $J_{4',5'} = 10.4$ Hz, H-4'); 3.49 (dd, 1H, $J_{2',3'} = 9.2$ Hz, H-2'); 3.27 (dd, 1H, $J_{3',4'} = 9.2$ Hz, H-3'); 3.01 (ddd, 1H, $J_{5',6''} = 6.0$ Hz, H-5'); 2.40 (s, 3H, C_5 -Me). Anal Calc. for $C_{15}H_{16}Cl_3N_3S_2O_4$ (472.8): C, 38.11; H, 3.41; N., 8.89. Found: C, 37.76; H, 3.32; N., 8.. MS: m/z (FAB) 663/665 (M + Na)⁺.

2,5-Dihydro-5,5-dimethyl-2-(2,3,4,6-tetra-*O*-acetyl-5-thio- β -D-glucopyranosylimino)-1,3,4-thiadiazole (11). From **5** (0.55 g). Yield: 0.58 g, (41%). δ_H ($CDCl_3$): 5.56 (t, 1H, $J_{4',5'} = 9.5$ Hz, H-4'); 5.47 (t, 1H, $J_{2',3'} = 9.5$ Hz, H-2'); 5.33 (t, 1H,

$J_{3',4'} = 9.5$ Hz, H-3'); 4.81 (d, 1H, $J_{1',2'} = 9.2$ Hz, H-1'); 4.27 (dd, 1H, $J_{6',6''} = 12.0$ Hz, H-6''); 4.16 (dd, 1H, $J_{5',6'} = 5.0$ Hz, H-6'); 3.56 (ddd, 1H, $J_{5',6''} = 3.5$ Hz, H-5'); 2.08, 2.50 [2xs, 6H, C₅-(Me₂)]; 2.00, 1.97, 1.92, 1.84 (4xs, 12H, 4xOAc). δ_C (CDCl₃): 176.8 (C-2); 170.3, 170.1, 169.0, 168.7 (C = O); 106.9 (C-5); 74.4 (C-2'); 73.9 (C-3'); 72.2 (C-4'); 64.8 [$^1J(^{13}\text{C}-^1\text{H}) = 153.0$ Hz, (C-1')]; 61.6 (C-6'); 42.6 (C-5'); 27.8, 27.1 [C₅-(Me₂)]; 20.5–20.3 (4xCOMe). Anal Calc. for C₁₈H₂₅N₃S₂O₈ (475.5): C, 45.46; H, 5.03; N, 8.84. Found: C, 45.49; H, 4.94; N, 8.61. MS: m/z (FAB) 476 (M + H)⁺.

2,5-Dihydro-5,5-dimethyl-2-(5-thio- β -D-glucopyranosylimino)-1,3,4-thiadiazole (11). From tetracatate **11** (0.40 g, 0.84 mmol) as described for analogue **9**. The oily product was purified on a SiO₂ column, using in gradient MeOH (0–10%) and CH₂Cl₂ to give **15** (0.21 g, 83%), as a foam. δ_H (DMSO-*d*₆/D₂O): 4.01 (d, 1H, $J_{1',2'} = 9.5$ Hz, H-1'); 3.85 (dd, 1H, $J_{6',6''} = 12.0$ Hz, H-6''); 3.82 (dd, 1H, $J_{5',6'} = 3.7$ Hz, H-6'); 3.60 (dd, 1H, $J_{4',5'} = 10.0$ Hz, H-4'); 3.54 (dd, 1H, $J_{2',3'} = 9.0$ Hz, H-2'); 3.29 (dd, 1H, $J_{3',4'} = 9.2$ Hz, H-3'); 3.11 (ddd, 1H, $J_{5',6''} = 5.5$ Hz, H-5'); 2.11, 2.52 [2xs, 6H, C₅-(Me₂)]. Anal Calc. for C₁₀H₁₇N₃S₂O₄ (307.4): C, 39.07; H, 5.57; N, 13.67. Found: C, 38.72; H, 5.48; N, 13.47. MS: m/z (FAB) 330 (M + Na)⁺.

5-Methyl-1-(2,3,4,6-tetra-*O*-acetyl-5-thio- β -D-glucopyranosyl)-4-imidazoline-2-thione (17). A solution of aminoacetone hydrochloride (0.30 g, 2.13 mmol) in water (15 mL) was neutralized with NaHCO₃ (0.17 g, 2.13 mmol) and added to a solution of **2** (0.82 g, 2.0 mmol) in acetone (30 mL) under nitrogen. The reaction mixture was stirred at 23°C for 30 min, the evaporated to dryness. The residue was partitioned between CH₂Cl₂ (3 \times 30 mL) and water (30 mL), and the combined organic extracts were dried (Na₂SO₄), filtered and evaporated to dryness. The crude foam was purified on a SiO₂ column (15 g) using, in gradient, MeOH (0–0.1%) and CH₂Cl₂ to give **17** (0.47 g, 51%) as a foam. δ_H (CDCl₃): 10.69 (s, 1H, NH); 6.13 (s, 1H, H-4); 5.80 (d, 1H, $J_{1',2'} = 9.3$ Hz, H-1'); 5.73 (t, 1H, $J_{2',3'} = 9.5$ Hz, H-2'); 5.40 (t, 1H, $J_{4',5'} = 10.0$ Hz, H-4'); 5.23 (t, 1H, $J_{3',4'} = 9.2$ Hz, H-3'); 4.26 (dd, 1H, $J_{6',6''} = 12.0$ Hz, H-6''); 4.07 (dd, 1H, $J_{5',6'} = 5.0$ Hz, H-6'); 3.46 (ddd, 1H, $J_{5',6''} = 3.7$ Hz, H-5'); 2.36 (s, 3H, C₅-Me); 2.01, 1.98, 1.94, 1.86 (4xs, 12H, 4xOAc). δ_C (CDCl₃): 170.4, 169.6, 169.3, 169.2 (C = O); 163.0 (C = S); 127.0 (C-4); 112.2 (C-5); 74.4 (C-2'); 73.6 (C-3'); 72.3 (C-4'); 63.0 (C-1'); 61.8 (C-6'); 42.2 (C-5'); 34.5 (C₅-Me); 20.6, 20.5, 20.4, 20.4 (4xCOMe). Anal Calc. for C₁₈H₂₄N₂S₂O₈ (460.5): C, 46.95; H, 5.25; N, 6.08. Found: C, 46.76; H, 5.16; N, 5.86. MS: m/z (FAB) 461 (M + H)⁺.

5-Methyl-1-(5-Thio- β -D-glucopyranosyl)-4-imidazoline-2-thione (18). From **17** (0.20 g, 0.43 mmol) as described for analogue **9**. Yield: 0.05 g, (86%), m.p. 146–150°C (from EtOH). δ_H (DMSO-*d*₆/D₂O): 6.17 (s, 1H, H-4); 5.31 (d, 1H, $J_{1',2'} = 9.0$ Hz, H-1'); 3.98 (t, 1H, $J_{2',3'} = 9.0$ Hz, H-2'); 3.88 (dd, 1H, $J_{5',6'} = 10.0$ Hz, H-4'); 3.52 (dd, 1H, $J_{3',4'} = 9.2$ Hz, H-3'); 3.23 (dd, 1H, $J_{6',6''} = 12.0$ Hz, H-6''); 3.14 (dd, 1H, $J_{5',6'} = 5.5$ Hz, H-6'); 3.09 (ddd, 1H, $J_{5',6''} = 3.5$ Hz, H-5'); 2.38 (s, 3H, C₅-Me). Anal Calc. for C₁₀H₁₆N₂S₂O₄ (292.4): C, 41.08; H, 5.52; N, 9.58. Found: C, 40.81; H, 5.40; N, 9.29. MS: m/z (FAB) 315 (M + Na)⁺.

2-(2,3,4,6-Tetra-*O*-acetyl-5-thio- β -D-glucopyranosyl)amino-2-thiazoline (19). To a solution of chloroethylamine hydrochloride (0.20 g, 1.72 mmol) in water (3 mL),

ether (4 mL) and M NaOH (4 mL) were added. The organic layer was decanted and the aqueous solution was extracted with ether (4 × 4 mL). The combined extracts were dried (MgSO₄) and then **2** (1.86 g, 4.58 mmol) was added. After stirring for 12 h at 23°C, the solvent was evaporated to dryness to give **19** as a crude hydrochloride. The residue was stirred for 2 h at 23°C in CH₂Cl₂ (50 mL) and aqueous solution of NaHCO₃ [3.78 g, 45.0 mmol, in water (50 mL)]. The organic layer was separated and the aqueous layer was extracted with CH₂Cl₂ (3 × 50 mL). The combined organic extracts were dried (Na₂SO₄), filtered and evaporated to dryness. The residue was poured onto a SiO₂ column (25 g) using, in gradient, MeOH (0–0.1%) in CH₂Cl₂ to give **19** as a foam (1.68 g, 82%). δ_{H} (CDCl₃): 8.20 (d, 1H, $J_{\text{NH,H-1'}}$ = 5.5 Hz, NH); 5.64 (t, 1H, $J_{2',3'}$ = 9.2 Hz, H-2'); 5.37 (t, 1H, $J_{3',4'}$ = 9.2 Hz, H-3'); 5.30 (t, 1H, $J_{4',5'}$ = 9.2 Hz, H-4'); 4.97 (d, 1H, $J_{1',2'}$ = 8.8 Hz, H-1'); 4.28 (dd, 1H, $J_{5',6'}$ = 4.7 Hz, H-6'); 4.16 (dd, 1H, $J_{6',6''}$ = 12.0 Hz, H-6''); 3.56 (ddd, 1H, $J_{5',6''}$ = 3.5 Hz, H-5'); 3.60 (t, 2H, J = 7.2 Hz, CH₂-5); 3.20 (t, 2H, J = 7.2 Hz, CH₂-4); 2.02, 1.99, 1.96, 1.88 (4xs, 12H, 4xOAc). Anal. Calc. for C₁₇H₂₄N₂S₂O₄ (448.5): C, 45.52; H, 5.39; N., 6.25. Found: C, 45.21; H, 5.30; N., 6.01. MS: m/z (FAB) 449 (M + H)⁺.

2-(5-Thio-β-D-glucopyranosyl)amino-2-thiazoline (20). From tetracetate **20** (0.35 g, 0.78 mmol) as described for analogue **11**. Yield: 0.18 g, 82%), m.p. 151–154°C. δ_{H} (DMSO-*d*₆/D₂O): 4.30 (d, 1H, $J_{1',2'}$ = 10.0 Hz, H-1'); 3.76 (dd, 1H, $J_{5',6'}$ = 6.0 Hz, H-6'); 3.70 (t, 1H, $J_{2',3'}$ = 9.0 Hz, H-2'); 3.59 (t, 2H, J = 7.1 Hz, CH₂-5); 3.55 (dd, 1H, $J_{6',6''}$ = 12.0 Hz, H-6''); 3.25 (pt, 1H, $J_{4',5'}$ = 10.0 Hz, H-4'); 3.19 (t, 2H, J = 7.1 Hz, CH₂-4); 3.07 (pt, 1H, $J_{3',4'}$ = 8.5 Hz, H-3'); 2.78 (m, 1H, $J_{5',6'}$ = 3.5 Hz, H-5'). Anal. Calc. for C₉H₁₆N₂S₂O₄ (280.4): C, 38.56; H, 5.75; N., 9.99. Found: C, 38.24; H, 5.66; N., 9.70. MS: m/z (FAB) 281 (M + H)⁺.

***N,N'*-Bis(2,3,4,6-tetra-*O*-acetyl-5-thio-β-D-glucopyranosyl)-*N*-(2-thiazolin-2-yl)thiourea (21).** To a solution of 2-chloroethylamine hydrochloride (0.06 g, 0.52 mmol) in pyridine (2 mL) was added **2** (0.42 g, 1.04 mmol). The mixture was kept at 23°C for 2 h, and then poured into ice–water. The solid product was poured onto column of silica gel (25 g), using, in gradient, ether (0–50%) and hexane to afford a white foam (0.63 g, 71%), tentatively identified as **21**. δ_{H} (CDCl₃): 13.0 (d, 1H, $J_{\text{NH,H-1'}}$ = 7.8 Hz, N'H); [5.68–5.40 (m, 2H, H-2', H-3')]_{A,B}; [5.33–5.29 (m, 1H, H-4')]_{A,B}; [5.00 (dd, 1H, $J_{1',2'}$ = 10.5 Hz, H-1')]_A; [4.95 (m, 2H, H-5')]_{A,B}; 4.62 (dt, 1H, = NCHH'); 4.48 (dt, 1H, J = 11.0 Hz, = NCHH'); [4.27 (m, 2H, H-6'')]_{A,B}; 4.21 (m, 1H, H-1')_B; [4.15–4.01 (m, 2H, H-6')]_{A,B}; 3.60 (t, 2H, J = 7.2 Hz, CH₂-5); 3.55 [(m, 2H, H-5')]_{A,B}; 3.01 (t, 2H, J = 7.0 Hz, S-CH₂); [2.04, 2.02, 2.00, 1.99, 1.96, 1.92, 1.90, 1.87, 1.85 (9xs, 24H, 8xOAc)]_{A,B}. Anal. Calc. for C₃₂H₄₃N₃S₄O₁₆ (854.0): C, 45.01; H, 5.08; N., 4.92. Found: C, 44.72; H, 4.92; N., 4.56. MS: m/z (FAB) 855 (M + H)⁺.

ACKNOWLEDGMENTS

We thank Professor E. De Clercq of Rega Institute for Medical Research, Catholic University of Leuven, Belgium for anti-viral screening.

REFERENCES

1. Mukerjee, A.K.; Ashare, R. Isothiocyanates in the chemistry of heterocycles. *Chem. Rev.* **1991**, *91*, 1–24.
2. Witczak, Z.J. Monosaccharides isothiocyanates and thiocyanates: synthesis, chemistry, and preparative applications. *Adv. Carbohydr. Chem. Biochem.* **1986**, *44*, 91–145.
3. Fuentes, J.; Pradera, A.M.; Robina, I. Preparations of 2-deoxyglycopyranosyl isothiocyanates. *Tetrahedron* **1991**, *47*, 5797–5810, and references therein.
4. Fuentes Mota, J.; Garcia Fernández, J.M.; Ortiz Mellet, C.; Pradera Adrián, M.A.; Cuevas Lorite, T. Synthesis of *N*-heterarylthiourea derivatives of carbohydrates. *J. Carbohydr. Chem.* **1990**, *9*, 837–851 and references therein.
5. Al-Soud, Y.A.; Al-Masoudi, N.A. Synthesis of 3'-1,2,4-triazolo- and 3'-1,3,4-thiadiazoliminothymidines. *Heteroat. Chem.* **2003**, *14*, 298–303.
6. Al-Masoudi, N.A.; Hassan, N.A.; Al-Soud, Y.A.; Schmidt, P.; Gaafer, A.E.-D.M.; Weng, M.; Marino, S.; Schoch, A.; Amer, A.; Jochims, J.C. Syntheses of *C*- and *N*-nucleosides from 1-aza-2-azoniaallene and 1,3-diaza-2-azoniaallene. *J. Chem. Soc., Perkin Trans. 1*, **1998**, 947–953.
7. Fuentes, J.; Moreda, W.; Ortiz, C.; Robina, I.; Ewlsh, C. Partailly protected D-glucopyranosyl isothiocyanates. Synthesis and transformations into thiourea and heterocyclic derivatives. *Tetrahedron* **1992**, *48*, 6413–6424.
8. Caballero, R.B.; Mota, J.F. A new method for the preparation of acylated glycosylamines and their transformations into glycosyl isothiocyanates and *NN'*-diglycosylthioureas. *Carbohydr. Res.* **1986**, *154*, 280–288.
9. Hanessian, S.; Pernet, A.G. Synthesis of naturally occurring *C*-nucleosides, their analoges, and functionalized *C*-glycosyl pPrecursors. *Adv. Carbohydr. Chem. Biochem.* **1976**, *33*, 111–188.
10. López, Ó.; Maya, I.; Ulgar, V.; Robina, I.; Fuentes, J.; Fernández-Bolaños, J.G. Expeditious synthesis of cyclic isourea derivatives of β -D-glucopyranosylamine. *Tetrahedron Lett.* **2002**, *43*, 4313–4316.
11. Maya, I.; López, Ó.; Fernández-Bolaños, J.G.; Robina, I.; Fuentes, J.A. A practical one-pot synthesis of *O*-unprotected glycosyl thioureas. *Tetrahedron Lett.* **2001**, *42*, 4313–4316.
12. Webb, T.R.; Mitsuya, H.; Broder, S.J. 1-(2,3-Anhydro- β -D-lyxofuranosyl)cytosine derivatives as potential inhibitors of the human immunodeficiency virus. *J. Med. Chem.* **1988**, *31*, 1475–1479, and references therein.
13. Reitz, A.B.; Tuman, R.W.; Marchione, C.S.; Jordan, A.D.; Bowden, C.R.; Maryanoff, B.E. Carbohydrate biguanides as potential hypoglycemic agents. *J. Med. Chem.* **1989**, *32*, 2110–2116.
14. Parrot-Lopez, H.; Galons, H.; Coleman, A.W.; Mahuteau, J.; Miocque, M. Victorised transport of drugs: synthesis of a new glycosyl derivatives of β -cyclodextrin. *Tetrahedron Lett.* **1992**, *33*, 209–212.
15. Suhadolink, R.J. *Nucleosides as Biological Probes*; Wiley Interscience: New York, 1979.
16. Garcia Fernández, J.M.; Ortiz, Mellet, C. Chemistry and developments of *N*-thiocarbonyl carbohydrate derivatives sugars isothiocyanates, thioamides,

- thioureas, thiocarbamates and their conjugates. *Adv. Carbohydr. Chem. Biochem.* **1999**, *55*, 35–135.
17. Gasch, C.; Pradera, M.A.; Salameh, B.A.B.; Molina, J.L.; Fuentes, J. Isothiocyanato derivatives of sugars in the stereoselective synthesis of spironucleosides and spiro-C-glycosides. *Tetrahedron: Asymmetry* **2001**, *12*, 1267–1277.
 18. Fernández-Bolaños, J.G.; Zafra, E.; López, Ó.; Robina, I.; Fuentes, J. Stereoselective synthesis of imidazolidine, and imidazole C- and N-pseudonucleosides. *Tetrahedron: Asymmetry* **1999**, *10*, 1267–3023 and references therein.
 19. *The Carbohydrates; Chemistry and Biochemistry, Thiosugars and Derivatives*, 2nd Ed.; Horton, D., Wander, J.D., Pigman, W., Eds.; Academic Press: New York, 1980; Vol. IB, 799–842.
 20. *The Merck Index*, 12th Ed.; Budavari, S., O'Neil, M.J., Smith, A., Heckelman, P.E., Kinneary, J.F., Eds.; Merck and Co.: Whitehouse Station, NJ, 1996; Nr. 9470.
 21. Fernández-Bolaños, J.G.; Al-Masoudi, N.A.; Maya, I. Sugars having sulfur in the ring. *Adv. Carbohydr. Chem. Biochem.* **2001**, *57*, 21–98, and references therein.
 22. Al-Soud, Y.A.; Al-Masoudi, W.A.; El-Halawa, R.A.; Al-Masoudi, N.A. Synthesis and antiviral activity of some 1,2,4-triazole C-nucleosides from 1-(chloroalkyl)-1-aza-2-azoniaallene salts. *Nucleosides Nucleotides* **1999**, *18*, 1985–1994.
 23. Al-Masoudi, N.A.; Al-Soud, Y.A.; Geyer, A. Synthesis and spectroscopic analysis of some acyclic C-nucleosides and the homo-C-analogues from 1-(chloroalkyl)-1-aza-2-azoniaallene salts. *Tetrahedron* **1998**, *55*, 751–758.
 24. Al-Masoudi, N.A.; Al-Soud, Y.A.; Lagoja, I.M. Synthesis and reactions of 1,5- and 1,3-dialkyl derivatives of (D-manno-pentitol-1-yl)-1H-1,2,4-triazole nucleosides derived from 1-(chloroalkyl)-1-aza-2-azoniaallene salts. *Carbohydr. Res.* **1999**, *318*, 67–74.
 25. Strumpel, M.K.; Buschmann, J.; Szilagyi, L.; Györgydeak, Z. Synthesis and structural studies of anomeic 2,3,4,6-tetra-O-acetyl-5-thio-D-glucopyranosyl azide. *Carbohydr. Res.* **1999**, *318*, 91–97, references therein.
 26. Wang, Q.; Jochims, J.C.; St. Köhlbrandt, L.; Dahlenburg, M.; Al-Talib, A.; Hamed, A.E. 1,2,4-Triazolium salts from the reaction of 1-aza-2-azoniaallene salts with nitriles. *Synthesis* **1992**, 710–718.
 27. Wang, Q.; Amer, A.; Mohr, S.; Ertel, E.; Jochims, J.C. [3 + 2]-Cycloadditions of 1-aza-2-azoniaallene cations to multiple bonds. *Tetrahedron* **1993**, *49*, 9973–9986, and references therein.
 28. El-Gazzar, A.-R.B.A.; Scholten, K.; Guo, Y.; Weißenbach, K.; Hitzler, M.G.; Roth, G.; Fischer, H.; Jochims, J.C. Cycloaddition of 1-aza-2-azoniaallene cations to isothiocyanates. *J. Chem. Soc., Perkin Trans. 1* **1999**, 1999–2010.
 29. L'abbé, G.; Verhelst, G.; Huybrechts, L.; Toppet, S. Tosylation of 2-(mono-substituted)amino-1,3,4-thiadiazoles. *J. Heterocycl. Chem.* **1977**, *14*, 515–516.
 30. L'abbé, G.; Verhelst, G.; Toppet, S. Trapping of thiaziridinimines with imines and nitriles. *J. Org. Chem.* **1976**, *41*, 3403–3406.
 31. Al-Soud, Y.A.; Wirschum, W.; Hassan, N.A.; Maier, G.-M.; Jochims, J.C. Reaction of 1-(chloroalkyl)-1-aza-2-azoniaallene salts with alkenes: preparation of cyclic azo, (azoalkyl)azonium, and formazanum compounds. *Synthesis* **1998**, 721–728.
 32. Moon, M.W. The chlorination of aldehyde and ketone phenylhydrazones. *J. Org. Chem.* **1972**, *37*, 383–385.

33. Moon, M.W. The chlorination of alkyl glycoxyate phenylhydrazones. *J. Org. Chem.* **1972**, *37*, 386–390.
34. Moon, M.W. The synthesis and properties of phosgene phenylhydrazones. *J. Org. Chem.* **1972**, *37*, 2005–2009.
35. Bothner, A.A.; Stephensen, R.L.; Lee, J.; Warren, C.D.; Jeanloz, R.W. Structure determination of the tetracsaccharide: transient nuclear overhauser effects in the rotating frame. *J. Am. Chem. Soc.* **1984**, *106*, 811–813.
36. Bax, A.; Davies, D.G. Practical aspects of two-dimensional transverse NOE spectroscopy. *J. Magn. Reson.* **1985**, *63*, 207–213.
37. Geisinger, C.; Ernst, R.R. Frequency offset effects and their elimination in NMR rotating-frame cross relaxation spectroscopy. *J. Magn. Reson.* **1987**, *75*, 261–271.
38. Summers, M.F.; Marzili, L.G.; Bax, A. Complete ^1H and ^{13}C assignments of coenzyme B₁₂ through the use of new two-dimensional NMR experiments. *J. Am. Chem. Soc.* **1986**, *108*, 4285–4294.
39. Sainsbury, M. *Rodd's Chemistry of Carbon Compounds, Thiazoles*; Ansell, M.F., Ed.; Elsevier: Amsterdam, 1986; 429–431. IV^c.
40. Katritzky, A.R.; Ambler, A.P. *Physical Constants for Organic Compounds*; C.R.C. Press: Cleveland, 1972; B-930.
41. Avalos, M.; Babiano, R.; Cintas, P.; Fuentes, J.; Jimenez, J.L.; Palacios, J.C. Reaction of glycosylisothiocyanates with 2-chloroethylamine. Synthesis and structure of *N*-nucleoside analogues. *Nucleosides Nucleotides* **1990**, *9*, 137–149.
42. Romero, D.L.; Olmsted, R.A.; Poel, T.J.; Morge, R.A.; Biles, C.; Keiser, B.J.; Kopta, L.A.; Friis, J.M.; Hosley, J.D.; Stefanski, K.J.; Wishka, D.G.; Evans, D.B.; Morris, J.; Stehle, R.G.; Sharma, S.K.; Yagi, Y.; Voorman, R.L.; Adams, W.J.; Tarpley, W.G. Targeting delaviridine/ateviridine resistance HIV-1, 2: identification of (alkylamino)piperidine-containing bis(heteroaryl)piperazines as broad spectrum reverse transcriptase inhibitors. *J. Med. Chem.* **1996**, *39*, 3769.

Received January 26, 2004

Accepted July 2, 2004

Copyright of Nucleosides, Nucleotides & Nucleic Acids is the property of Marcel Dekker Inc. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.