

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/312590146>

Analysis of Water Quality Using Physico-Chemical Parameters in the Shatt AL-Arab Estuary, Iraq

Article in *Open Journal of Marine Science* · September 2015

CITATIONS

2

READS

141

6 authors, including:



[Hamid T. Al-Saad](#)

University of Basrah

255 PUBLICATIONS 1,000 CITATIONS

[SEE PROFILE](#)



[Abdulzahra Alhello](#)

University of Basrah

42 PUBLICATIONS 101 CITATIONS

[SEE PROFILE](#)



[Mohsin Alhilo](#)

Rutgers, The State University of New Jersey

9 PUBLICATIONS 60 CITATIONS

[SEE PROFILE](#)



[Wesal Fakhri Hassan](#)

University of Basrah

42 PUBLICATIONS 121 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Pollution indexes in Shatt Al-Arab River [View project](#)



Polycyclic Aromatic Hydrocarbons in Some Fishes from the Iraqi Marine Waters [View project](#)

OPEN  ACCESS

ISSN1927-6648

International Journal of

Marine Science



Published by BioPublisher
<http://biopublisher.ca>

2015
Volume5

Publisher

Sophia Publishing Group

Edited by

Editorial Team of International Journal of Marine Science

Email: edit@ijms.biopublisher.ca

Website: <http://ijms.biopublisher.ca>

Address:

11388 Stevenston Hwy,

PO Box 96016,

Richmond, V7A 5J5, British Columbia

Canada

International Journal of Marine Science (ISSN 1927-6648) is an open access, peer reviewed journal published online by BioPublisher.

The journal publishes all the latest and outstanding research articles, letters and reviews in all areas of marine science, the range of topics containing the advancement of scientific and engineering knowledge regarding the sea; from chemical and physical to biological oceanography, from estuaries and coastal waters to the open ocean; as well as including fisheries, socio-economic science, co-management, ecosystems and other topical advisory subjects.



BioPublisher, operated by Sophia Publishing Group (SPG), is an international Open Access publishing platform that publishes scientific journals in the field of life science. Sophia Publishing Group (SPG), founded in British Columbia of Canada, is a multilingual publisher.

Open Access

All the articles published in International Journal of Marine Science are Open Access, and are distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



BioPublisher uses CrossCheck service to identify academic plagiarism through the world's leading plagiarism prevention tool, iParadigms, and to protect the original authors' copyrights.

Latest Content

[Ethological and phenetic characterization of sea turtle's fishing stock in Jardines del Rey \(Cuba\)](#)

I. Lee-González, A. Ruiz-Urquiola, E. Pérez-Bermúdez, B.L.J. Petric

[Study of impact of climatic variability on the sea surface temperature and Chlorophyll-a concentration using Statistical analysis on Satellite derived data for The arabian sea](#)

Naeem Syed, Syed Ahmed

[First report on mass reproductive swarming of a polychaete worm, *Dendronereis aestuarina* \(Annelida, Nereididae\) Southern 1921, from a freshwater environment in the south west coast of India](#)

P.R. Jayachandran, M.P. Prabhakaran, C.V. Asha, Akhilesh Vijay, S. Bijoy Nandan

[Community Structure of Birds at Islands in Karon River in Ahvaz City in Spring and Summer 2014](#)

Behrouz Behrouzi-Rad, Pyvand Maktabi, Alireza Jafarnejad

[Phenotypic Plasticity and Genetic Variation of Two Wild Populations of Green tiger Shrimp \(*Penaeus semisulcatus*-De Haan, 1844\)](#)

D.H.N. Munasinghe, J.D.M. Senevirathna

[Fishers' attitude towards performance of Beach Management Units \(BMUs\) in regulating fishery and reducing poverty: case study of two BMUs Lake Victoria, Tanzania](#)

Luomba Onyango

[Economic performance efficiency of gillnet based on mesh size and net color](#)

Fatemeh Radfar, Houshang Ansari, Mohammad Gerami, Mehdi Dastbaz

[Characteristics of sub-bottom profile acquired in Shatt Al-Arab River, Basrah-Iraq](#)

H.A. Alaamer

[Cold Lugol's solution as an alternative of formaldehyde-base preservative in preserving stomach content of shellfish](#)

Tan Soon, Julian Ransangan

[Marine Fishes Captured in within and outside potential Fishing zones off Ratnagiri coast, Maharashtra, India](#)

Tingote R. S., Pawar M.B., Mane U.H.

[Nutritional value of important commercial fish from Iraqi waters](#)

A.A. Hantoush, Q.H. Al-Hamadany, A.S. Al-Hassoon, H.J. Al-Ibadi

[Changes in the Population of Wintering Waterbirds in Gomishan Wetland at Caspian Sea Coast, Iran](#)

Behrouz Behrouzi-Rad, Ramazanali Ghaeimi

[Colonization of formulated substrates of different composition by benthic macroinvertebrate community in an estuarine ecosystem: a case study of response of benthic invertebrates to substrate alteration](#)

Rolande. Uwadiae, Deborah Felix

[Effect on white gut and white feces disease in semi intensive *Litopenaeus vannamei* shrimp culture system in south Indian state of Tamilnadu](#)

Durai V, B. Gunalan, P. Johnson, M. L. Maheswaran, M. Pravinkumar

[Occurrence of *Lepocreadioides orientalis* Park, 1939 and *Lepocreadioides* sp. \(Trematoda: Lepocreadiidae\) from *Psettodes erumei* and *Sillago sihama* from the Iraqi Marine water](#)

Majid Bannai, Essa T. Muhammad

[Seasonal Variation of Environmental Factors Influencing Benthic Macrofaunal Diversity in Parangipettai and Cuddalore Coast \(Southeast coast of India\)](#)

Sundaravarman. K, A. Saravanakumar, M. Pravinkumar, K. Kathiresan

[Analysis on Global Eel Aquaculture Conditions](#)

Sun Zhaoqun, Wan Rong, Zhu Yugui, Talib Muhammad, Moshin Muhammad

[Contribution of Fish Production and Trade to the Economy of Pakistan](#)

Muhammad Mohsin, Mu Yongtong, Khadim Hussain, Aamir Mahmood, Sun Zhaoqun, Kiran Nazir, Wang Wei

[Sexual differentiation and variations sexual characteristics *Rapana venosa* \(Valenciennes, 1846\)](#)

Igor P. Bondarev

[Socio-economics status and adaptations of purse seine fishermen in Bali coastal village, Indonesia](#)

Achmad Zamroni

[Potential risk of some heavy metals in *Pampus chinensis* \(Euphrasen\) Chinese silver pomfret Stromateidae collected from Karachi Fish Harbour, Pakistan](#)

Quratulan Ahmed, Levent Bat

[Distribution and concentration several types of heavy metal correlated with diversity and abundance of microalgae at Tallo Estuary, Makassar, South Sulawesi, Indonesia](#)

Nita Rukminasari

[Environmental factors structuring the fish assemblage distribution and production potential in Vembanad estuarine system, India](#)

C. V. Asha, R. I. Cleetus, P. S. Suson, S. Nandan

[Diversity and seasonal changes of zooplankton communities in the Shatt Al-Arab River, Basrah, Iraq, with a special reference to Cladocera](#)

M.F. Abbas, S.D. Salman, S.H. Al-Mayahy

[Determination of heavy metals in low value food fish from commercial landing centers \(India\) by Inductively Coupled Plasma Optical Emission Spectrometer](#)

M. Pravinkumar, A. R. Logesh, C. Vishwanathan, G. Ponnusamy, V. Elumalai, S.M. Raffi, K. Kathiresan

[Spatio-temporal variations of macrobenthic annelid community of the Karnafuli River Estuary, Chittagong, Bangladesh](#)

Md. Habibur Rahman Molla, Md. Al-Imran, Md. Aktaruzzaman, Shamindranath Mandol, Mohammad Saydul Islam Sarkar, M. Shafiqul Islam

[The Effect of Biostimulation and Biostimulation-Bioaugmentation on Biodegradation of Oil-Pollution on Sandy Beaches Using Mesocosms](#)

Yeti Darmayati, Harpasis S. Sanusi, Tri Prartono, Dwi Andreas Santosa, Ruyitno Nuchsin

[Sustainable management of Fisheries in Tamil Nadu](#)

J.S. Amarnath, A. Mouna

[Adsorption of copper\(II\) and lead\(II\) ions from aqueous solutions by porcellanite](#)

A.Y. Hmood, T.E. Jassim

[Length-weight relationship of coral reef associated fishes of Cuddalore, southeast coast of India](#)

N. Jayaprabha, S. Purusothaman, M. Srinivasan

[Reconstructing the past fluctuations of Urmia Lake](#)

Homayoun Khoshnavan, Anahita Jabbari

[Note on the Genus *Marettia* gray, 1855 and Description of *Marettia planulata* \(Lamarck, 1816\) \(Echinoidea, Spatangoida\), off Thoothukudi Coast of Gulf of Mannar, India \(08° 35' 22.5" N 78° 27' 40.9 E\) – \(310 M\)](#)

T. Vaitheeswaran, T. Rajasekaran, S. Balasubramani

[Checklist of intertidal benthic macrofauna of a brackish water coastal lagoon on east coast of India: The Chilika lake](#)

Debasish Mahapatro, R.C. Panigrahy, S. Panda, R.K. Mishra

[Survival of Tropical Benthic Amphipod *Grandidierella bonnieroides* Stephensen 1948 on Different Sediment Particle Size: Implications for Ecotoxicological Testing](#)

Dwi Hindarti, Zainal Arifin, Tri Prartono, Ety Riani, Harpasis S. Sanusi

[Coastal sedimentary morphology of Urmia Lake](#)

Homayoun Khoshnavan, Anahita Jabbari

[Recruitment Pattern, Virtual Population Analysis \(Vpa\) and Exploitation Status of *Lethrinus lentjan* \(Lacepede, 1802\) Exploited in Thoothukudi Coast, Tamil Nadu, India](#)

Vasantharajan M., Jawahar P., Venkatasamy M.

[Comparative Study on Macrobenthic Community Structure with Special Reference to Oligochaetes During Drought and Flooded Phases in a Tropical Kole Wetland, India](#)

Vineetha S., Bijoy Nandan S., Rakhi Gopalan K.P.

[Diet and trophic status of fish landed by tropical artisanal bait fishermen, Mida Creek Kenya](#)

Kihia C.M., Hendrick Y., Muthumbi A., Okondo J., Nthiga A., Njuguna V.M.

[Temporal Variation of Population Structure of the Invasive *Isognomon bicolor* \(Mollusca, Bivalvia\), *Brachidontes solisianus* \(Mollusca, Bivalvia\) and *Ulva* spp. \(Chlorophyta, Ulvales\) Biomass, Pernambuco – Brazil](#)

Guimaraens M.A., Dias V.

[A New Record of Sea Snail *Cochlespira Travancorica Travancorica* \(E. A. Smith, 1896\) \(Family: Cochlespiridae\) \(Mollusca: Gastropoda: Turridae\) off Thoothukudi Coast of Gulf of Mannar, Southeast Coast of India \(08° 53.6'N 78° 16'E and 08° 53.8'N 78° 32'E\) \(310 M\)](#)

Vaitheeswaran T., Jayakumar N., Venkataramani V.K.

[Seasonal Variation in Occurrence of Heavy Metals in *Perna viridis* from Manora Channel of Karachi, Arabian Sea](#)

Qari Rashida., Ajiboye Olufemi., Manzoor Rana., Afridi Abdul Rahim.

[Integrating Knowledge to Assess Coastal Vulnerability to Sea-Level Rise In El- Mex Bay, Egypt: The Application of the DIVA Tool](#)

El-Raey M., Nasr S., Hendy Dina M.

[Interrelationship Between Planktonic Diatoms and Selected Governing Physicochemical Parameters of the Hooghly Estuary, Bay of Bengal](#)

De T. K., Mukherjee A., Das S., Chakraborty S., De M., Maiti T.K.

[Habitat Inventory Parameters and Drainage Analysis of River Siang in Arunachal Pradesh](#)

Das B.K., Kar D.

[Analysis of Water Quality Using Physico-Chemical Parameters in the Shatt AL-Arab Estuary, Iraq](#)

AL-Saad H. T., Alhello A. A., AL-Kazaeh D. K., Al-Hello M. A., Hassan W. F., Mahdi S.

Analysis of Water Quality Using Physico-Chemical Parameters in the Shatt AL-Arab Estuary, Iraq

AL-Saad H. T.[✉], Alhello A. A., AL-Kazaeh D. K., Al-Hello M. A., Hassan W. F., Mahdi S.

Dept. Of Marine Environmental Chemistry, Marine Science Center-University of Basra, Iraq

✉ Corresponding author email: htalsaad@yahoo.com

International Journal of Marine Science, 2015, Vol.5, No.49 doi: [10.5376/ijms.2015.05.0049](https://doi.org/10.5376/ijms.2015.05.0049)

Received: 01 Jul., 2015

Accepted: 01 Aug., 2015

Published: 30 Aug., 2015

Copyright © 2015 AL-Saad et al., This is an open access article published under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Preferred citation for this article: AL-Saad H. T., Alhello A. A., AL-Kazaeh D. K., Al-Hello M. A., Hassan W. F and Mahdi S., 2015, Analysis of Water Quality Using Physico-Chemical Parameters in the Shatt AL-Arab Estuary, Iraq, International Journal of Marine Science, 5(49): 1-9

Abstract Six sampling stations, Al-Qurna, paper factory, Al- Sindebad, AL-Ashar, Al-Sebah and Al-Fao , representing different sectors of Shatt AL-Arab estuary were selected to the distribution of the Physical and chemical parameters. Water sample analyzed from August 2008 to July 2009, Dissolved Oxygen (DO), Turbidity, Total Suspended Solids (TSS), Total Dissolved Solids (TDS), Electrical Conductivity (EC) , temperature, salinity and pH as well as nutrients were also analyzed . The mean range of the following parameters were recorded: pH (7.56-7.84), EC (1.29-3.22 mmohs/cm²), Ca (87.18-130.26 mg/l), Mg (60.35-111.17 mg/l), Cl (304.7-753.31 mg/l), TSS (11-38.58 mg/l), TDS (891-2040.42 mg/l), DO (5.16-10.05 mg/l), turbidity (4.57-39.03 FTU), water temperature (21.09-22.47 °C), air temperature (23.54-35.26 °C), SO²⁻ (285.73-663.89 mg/l), HCO₃ (204.39-255.22 mg/l), and Nutrient(NO₂, NO₃, PO₄). Also the seasonal variation of all the parameters were monitored during this study and the result showed some fluctuation in some of them during different seasons at different locations of the estuary The results indicate that the water is no-polluted and can be used for domestic, irrigation and fisheries.

Keywords *Shatt AL-Arab estuary; Physico-Chemical Parameters; Monthly variation*

Introduction

As water is one of the most important compounds of the ecosystem, but due to increased human population, industrialization, use of fertilizers in the agriculture and man-made activity, all these factors effect on water quality. The natural aquatic resources are causing heavy and varied pollution in aquatic environment leading to pollute water quality and depletion of aquatic biota. It is therefore necessary that the quality of drinking water should be checked at regular interval, It is difficult to understand the biological phenomena fully because the chemistry of water reveals much about the metabolism of the ecosystem and explain the general hydro - biological relationship (Basavaraja et al., 2011). The Shatt al Arab River is formed after the confluence of the Euphrates and the Tigris Rivers near the city of Qurna in southern Iraq. The southern part of the river constitutes the border between Iran and Iraq until it discharges into the Arabian Gulf, with a total length of 192 km, the Shatt al Arab widens over its course, expanding from a width of 250-300 m near the Euphrates-Tigris confluence to almost 700 m near the city of Basrah and more than 800 m as it approaches the river mouth.

An area of 145,190 km² drains directly to the Shatt al Arab region downstream of the Euphrates-Tigris confluence (excluding the Euphrates and Tigris Basin areas). Several tributaries join the Shatt al Arab during its course, most importantly the Karkheh and the Karun Rivers. The Shatt al Arab Delta area is classified as estuarine-deltaic because the river's sediment seeps into a shallow, narrow part of the Arabian Gulf. The Shatt al Arab Delta is 140 km wide and splits into more than 10 branches. The landscape is characterized by green marshy areas, lakes, lagoons and estuaries, bordered by irrigated lands and date palm plantations and surrounded by desert (UN-ESCWA and BGR2013).

Shatt AL-Arab depends on some water sources tributaries for feeding. These tributaries are Al-Hammar, Al-Huyaza and Al-Qurna marshes and Al-Karoon River which flows in its left side. Shatt AL-Arab also affected by the climate tide comes from Arabian Gulf which reduces discharge of Shatt AL-Arab water to the sea. Shatt AL-Arab's two banks are characterized to be low in its flow area reaching Basrah , its full with water channels which are used for irrigation agriculture land (Hussain et al., 1991) studies on water quality and

different land use to prevent the salinization in this soil which irrigated with Shatt Al-Arab water.

Water quality characteristics of an aquatic environment are of great significance for the proper understanding of distribution, growth and physiological function of the biotic community inhabiting the area. Understanding of water quality is also a very important factor in the semi-enclosed systems where nutrients and pollutants may be concentrated and where the growth and proliferation of plankton is largely dependent on the environmental and physico-chemicals variables which can either support or limit their production capacities. Although scientific measurements are used to define water quality, it is not simple to say that water is good or bad. So the determination is typically made, relative to the purpose of the water. Whether it is for drinking or to wash a car or for some other purpose, poor water quality can pose a health risk for people and ecosystem.

As described by (Tabari et al., 2011) that one major goal of surface water quality data collection may be the estimation of magnitude of changes in the concentration of various constituents. Such estimates may be to determine water quality impacts over time due to human activities, changes brought about by regulations, improvements to waste water treatment facilities, or land use changes. Also, the goal of water quality assessment includes: accurately assessing pollution level identifying the key pollutants' effect on the quality of water and future development trend (Al-Hejuje 2015).

This field study was initiated to assess the extent and magnitude of water quality threats of the Shatt AL-Arab-estuary. Six sampling locations were selected to cover the distribution of the pollutants in this estuary. The physical and chemical aspects of the water were assessed which will provide valuable indications of the overall health of the ecosystem of the Shatt AL-Arab estuary.

Materials and Methods

Water sampler were used to collected water samples monthly, from August 2008 to July 2009. from 6 fixed stations as shown in Figure. (1).The stations on different sectors of Shatt Al-Arab, are Al-Qurna, paper factory, Al- Sindebad, Al-Ashar, Al-Sebah and Al-Fao (indicate the number - station 1-6) .The air and water

temperatures were measured by simple thermometers with a range of 10-100°C graduate at 0.2°C.; pH was measured using pH meters model HANNA HI-9821, and Electrical Conductivity (EC) was measured using portable digital conductivity meters (WTW 3301).The TDS, TSS, HCO₃ and SO₄²⁻ and DO, Mg, Ca and Cl were analyzed according to APHA (2005) standard method. Nutrient (NO₃, NO₂, PO₄,) were measured by using Parson et al. (1984) methods.

Result and Discussion

Water is a natural resource with limited and uneven distribution in time and space. All forms of life and all human activities are dependent on water. Water resources in Shatt al-Arab estuary are of great importance to human life and economy and are the main source of meeting the demand for drinking water, for irrigation of lands and industries. Lack of water is considered as a limiting factor of socio-economic development of a country. The rates of biological and chemical processes depend on temperature. Aquatic organisms from microbes to fish are dependent on certain temperature ranges for their optimal health. Optimal temperatures for fish depend on the species: some survive best in colder water, whereas others prefer warmer water. Benthic macro invertebrates are also sensitive to temperature and will move in the stream to find their optimal temperature. If temperatures are outside this optimal range for a prolonged period of time, organisms are stressed and can die.




Figure 1 .Map of the Shatt Arab Estuary showing the sampling locations

Results of the present study showed that the air temperature in all stations were high during summer and low during winter. Temperature at Al-Qurna ranged from 12.0 to 38.8°C. paper factory ranged from 14.8-36.4°C, Al- Sindebad ranged from 14.0-33.0°C, Al-Ashar ranged from 12.0-34.0°C, in Al-Sebah ranged from 16.5-40.3°C and in Al-Fao ranged from 17.4 -39.0°C .

Water temperature varies in accordance with ambient air temperature. The spatial variations in water temperature among stations may be due to the variation in the sampling time at each station, or the depth of water at each station .The minimum (12°C) and maximum (45°C) water temperature were recorded during winter and summer months respectively, with a mean water temperature of (27.83°C) (Tables 2). The seasonal mean values of water temperature during winter and summer were (13.6°C) and (44°C) respectively. The highest variations in water temperature between winter and summer was depending mainly on the variations in the air temperature , which was the highest in summer because of the long day period , and the lowest in winter because of the short day period. A significant correlation ($r = 0.926$, $p < 0.01$) was found between water and air temperatures. These results agree with 13-33.4°C obtained by Al-Hejuje 2015 .

The value of pH in natural water is affected by geological shed water and balance of CO₂, H₂CO₃ and CO₃. It ranged from 4.5-8.0 (Gems 1997), effected by processes which can be changed the concentration of dissolved CO₂. pH value effected on transpiration process between nutrient type and mineral it affection pollution toxic , which containing basic and acidic compound because of the ionization process (Gems, 1977). PH affects many chemical and biological processes in the water. For example, different organisms flourish within different ranges of pH. The largest variety of aquatic animals prefers a range of 6.5-8.0. pH outside this range reduces the diversity in the stream because it stresses the physiological systems of most organisms and can reduce reproduction. Low pH can also allow toxic elements and compounds to become mobile and "available" for uptake by aquatic plants and animals. This can produce conditions that are toxic to aquatic life, particularly to sensitive species of fish. Changes in acidity can be caused by atmospheric deposition (acid rain), surrounding rock,

and certain wastewater discharges Variations in pH were very large during the study period .The minimum (7. 0) and maximum (8.91) pH was recorded during autumn and winter months respectively, with a mean pH of 7.92 . The mean values of pH during winter and summer were 7.92 and 8.21, respectively This result agrees with Al- Kazaeh (2014)(7.4-7.9) and Al-Hejuje (2015) (7.4-8.52) .

DO levels fluctuate seasonally and over a 24-hour period. They vary with water temperature and altitude. Cold water holds more oxygen than warm water and water holds less oxygen at higher altitudes. Thermal discharges, such as water used to cool machinery in a manufacturing plant or a power plant, raise the temperature of water and lower its oxygen content. Aquatic animals are most vulnerable to low DO levels in the early morning on hot summer days when stream flows are low, water temperatures are high, and aquatic plants have not been producing oxygen since sunset. Fluctuations in DO were wide along Shatt Al-Arab during winter and summer. The minimum (5.3 mg/L) and maximum (9.3 mg/L). DO were recorded during summer and winter months, respectively, with a mean DO of 7.63 mg/L . The mean values of DO during summer and winter were 6.37 and 8.57 mg/L, respectively. The dissolved oxygen content of water is influenced by the sources, raw water temperature and chemical or biological processes taking place in the aquatic system. The reduction of dissolved oxygen at some stations (especially at Al-Ashar) during Summer and Autumn seasons can be due to the increasing in the untreated domestic sewage which polluted the water with reducing substances (such as nitrite) or due to organic pollutants which fasten the consumption of dissolved oxygen in water during warm months Low levels of dissolved oxygen in some stations are a sign of pollution and then , it is an essential parameter to be taken into consideration when the objective is to assess the water quality (ALHejuje ,2015).

Total solids suspended levels varied over wide range of 2.50-332.00 mg/L (Table 1). The minimum (4 mg/L) and maximum (187 mg/L) levels were recorded during summer months, respectively, with a mean value of 59.26 mg/L. The seasonal mean values of total solid suspended during winter and summer were 33 and 73.83 mg/L, respectively (Table.2). Fluctuation

Table 1 Regional distributions of physical and chemical parameters in six stations of Shatt AL-Arab Estuary during Aug.2008-Jul.2009 (All parameters determine in mg/L, while Turbidity in FTU, and EC in dS/m)

Season	Station	Month	PH	EC	TDS	TSS	Turb.	Ca	Mg	Cl	HCO ₄	NO ₂	NO ₃	PO ₄	SO ₄
Summer	1	Aug.08	7.60	2.15	1930	60	63	80	59	355	58	1.03	10.32	2.74	340
Autumn	1	Sep.08	7.10	2.3	2510	96	40	128	44	2057	183	0.58	8.85	1.58	467
Autumn	1	Oct.08	7.20	2.38	1568	48	17	80	413	390	268	0.12	17.74	2.49	369
Autumn	1	Nov.08	7.70	2.12	2588	62	26	248	194	660	366	0.98	3.2	2.40	459
Winter	1	Des.08	8.18	2.78	1334	46	18	160	112	638	305	5.39	36.81	2.84	591
Winter	1	Jan.09	8.28	2.7	1596	20	2	360	753	851	976	2.57	37.75	1.82	947
Winter	1	Feb.09	8.18	2.75	1992	48	22.6	160	49	532	305	1.37	18.57	2.21	4227
Spring	1	Mar.09	8.58	2.84	1948	42	12.7	176	78	496	183	2.55	41.78	2.38	378
Spring	1	Apr.09	8.34	1.56	1432	72	1.36	160	24	241	183	0.47	78.6	3.88	137
Spring	1	May-09	8.32	1.3	490	52	31.9	48	49	269	183	0.97	40.34	0.12	240
Summer	1	Jun.09	7.87	1.47	1282	187	52	200	24	461	98	1.86	24.52	2.68	80
Summer	1	Jul.09	8.05	1.14	1084	62	67	32	68	291	305	0.9	38.17	3.65	206
Summer	2	Aug.08	7.30	4.07	2700	52	42	120	48	638	83	1.24	12.38	13.63	505
Autumn	2	Sep.08	7.00	2.46	1726	80	50	120	58	1702	244	1.18	8.32	3.93	404
Autumn	2	Oct.08	7.10	2.47	1594	52	48	80	97	888	220	0.07	14.7	3.22	561
Autumn	2	Nov.08	7.50	2.78	2852	62	17	176	126	589	286	1.68	3.66	9.75	492
Winter	2	Des.08	8.03	2.79	966	34	17	104	268	674	268	6.6	31.69	4.00	653
Winter	2	Jan.09	7.89	2.88	1826	18	3	240	899	496	915	1.93	23.38	6.37	534
Winter	2	Feb.09	8.02	2.88	2056	50	21.2	120	170	355	610	1.35	19.26	2.86	4462
Spring	2	Mar.09	8.38	2.95	2142	48	9.29	184	92	1028	183	3.52	43.78	7.85	314
Spring	2	Apr.09	8.12	1.68	2748	152	0.8	80	49	262	183	0.47	78.54	7.53	186
Spring	2	May-09	8.05	1.39	1044	50	27.1	64	24	284	220	1.31	39.71	0.18	193
Summer	2	Jun.09	7.97	1.63	1150	52	32.8	200	49	425	85	1.57	24.98	7.81	80
Summer	2	Jul.09	7.19	2.48	1370	75	50	64	40	815	390	0.48	22.83	18.92	245
Summer	3	Aug.08	7.60	2.24	1557	4	37	96	64	383	87	9.11	81.61	6.46	602
Autumn	3	Sep.08	7.30	2.37	1564	68	36	120	34	603	317	1.16	7.41	2.61	435
Autumn	3	Oct.08	7.10	2.41	1536	104	163	160	316	568	195	0.49	21.31	2.65	480
Autumn	3	Nov.08	7.60	2.49	1200	48	18	144	180	547	244	1.83	6.78	2.60	509
Winter	3	Des.08	7.98	2.83	1826	22	11	152	268	674	268	7.26	42.34	5.33	620
Winter	3	Jan.09	8.33	2.75	1748	18	2	160	899	461	732	1.77	22.72	2.60	377
Winter	3	Feb.09	8.09	2.58	1984	68	16	160	146	355	610	1.32	10.33	1.43	523
Spring	3	Mar.09	8.39	2.88	2200	68	27.2	184	83	567	146	2.25	39.68	3.97	383
Spring	3	Apr.09	8.14	1.66	3092	36	0.84	80	49	262	183	0.54	78.58	7.13	178
Spring	3	May-09	7.96	1.31	636	34	20.8	56	39	284	195	1.33	48.24	0.61	253
Summer	3	Jun.09	8.12	1.5	910	136	24.7	120	73	390	61	0.81	43.57	4.88	110
Summer	3	Jul.09	8.27	1.44	780	45	56	48	54	241	268	1.22	19.51	3.98	174
Summer	4	Aug.08	7.60	2.31	1510	48	29	80	42	425	58	11.02	110.12	5.78	1256
Autumn	4	Sep.08	7.30	2.34	766	62	46	104	53	815	183	1.19	8.58	3.10	435
Autumn	4	Oct.08	7.10	2.42	1598	148	97	40	170	568	195	0.06	17.62	2.65	772
Autumn	4	Nov.08	7.70	2.54	2668	68	24	160	272	490	244	0.88	3.54	2.85	409
Winter	4	Des.08	8.05	2.85	1406	32	15	144	117	709	256	7.26	37.98	4.18	724
Winter	4	Jan.09	8.65	2.75	1786	20	2	240	680	603	915	1.93	22.74	2.40	449
Winter	4	Feb.09	8.27	2.81	1900	40	11.6	160	73	355	488	2.1	10.97	0.52	464

Continuing table 1

Season	Station	Month	PH	EC	TDS	TSS	Turb.	Ca	Mg	Cl	HCO ₄	NO ₂	NO ₃	PO ₄	SO ₄
Spring	4	Mar.09	8.59	2.83	2102	52	4.76	168	117	496	146	3.27	40.4	1.76	405
Spring	4	Apr.09	8.33	1.79	3248	36	2.20	80	49	284	159	0.75	73.88	4.70	228
Spring	4	May-09	7.97	1.33	926	30	11.2	48	44	248	195	2.00	48.28	0.43	284
Summer	4	Jun.09	8.26	1.5	898	156	26.1	200	24	319	61	1.23	38.44	3.84	95
Summer	4	Jul.09	8.09	1.46	732	92	27.37	56	44	241	281	2.46	23.73	3.74	178
Summer	5	Aug.08	7.60	2.23	1594	78	35.0	96	67	425	53	7.89	78.86	2.98	1674
Autumn	5	Sep.08	7.30	2.33	1426	128	47.0	96	53	638	244	1.18	7.270	3.10	413
Autumn	5	Oct.08	7.10	2.45	1644	48	35.0	120	73	497	195	0.11	17.89	3.02	470
Autumn	5	Nov.08	7.70	2.43	2706	62	28.0	160	253	518	244	0.81	4.120	2.20	407
Winter	5	Des.08	7.93	2.87	1916	22	6.00	152	117	744	244	10.34	35.47	5.16	691
Winter	5	Jan.09	8.46	2.8	1794	30	4.00	320	437	638	732	1.95	27.24	3.12	642
Winter	5	Feb.09	8.26	2.97	1910	40	13.4	160	146	390	488	1.45	17.34	0.52	523
Spring	5	Mar.09	8.59	2.85	5566	52	5.84	200	97	532	134	3.45	36.86	2.55	326
Spring	5	Apr.09	8.18	1.74	3048	84	0.99	80	73	269	146	0.75	76.96	7.05	205
Spring	5	May-09	7.97	1.3	142	34	16.5	56	15	262	183	1.62	41.41	0.43	230
Summer	5	Jun.09	8.19	1.51	890	10	31.9	160	73	389	85	1.13	42.40	4.21	100
Summer	5	Jul.09	8.07	1.48	704	36	59.0	56	44	262	317	2.02	28.29	3.82	200
Summer	6	Aug.08	7.60	2.25	1540	26	52.0	96	75	425	58	6.99	67.40	2.27	2106
Autumn	6	Sep.08	7.40	2.33	1388	54	41.0	112	49	532	244	1.09	8.400	3.23	301
Autumn	6	Oct.08	7.20	2.44	1668	56	48.0	120	170	604	195	0.37	18.96	4.00	590
Autumn	6	Nov.08	7.70	2.51	2600	62	26.0	200	97	483	232	0.80	3.700	4.15	477
Winter	6	Des.08	7.91	2.86	1796	16	3.00	160	117	738	281	6.93	39.72	5.34	766
Winter	6	Jan.09	8.91	2.81	1820	8	3.00	200	535	603	976	2.20	26.65	3.12	345
Winter	6	Feb.09	8.27	2.86	1604	62	21.7	160	146	355	610	2.10	18.08	0.26	669
Spring	6	Mar.09	8.54	2.82	2048	76	6.62	176	97	674	146	3.55	40.42	2.38	669
Spring	6	Apr.09	8.36	1.78	3168	64	5.06	80	194	268	146	0.66	73.78	6.48	163
Spring	6	May-09	8.01	1.35	2400	54	15.8	40	39	248	183	2.66	47.35	0.49	239
Summer	6	Jun.09	8.16	1.51	890	142	23.42	200	122	425	85	1.69	39.15	3.78	150
Summer	6	Jul.09	8.14	1.46	672	68	66.0	48	49	241	305	4.05	31.83	0.14	167

in NO₃-N varied over a wide range of 3.2-110.12 mg/L (Table 1). The minimum (3.2 mg/L) and maximum (110.12 mg/L) NO₃-N were recorded during autumn months with a mean NO₃-N of 9.93 mg/L (Table 2). The seasonal mean values of NO₃-N during winter and summer were 26.61 and 41.01 mg/L, respectively (Table 2).

Fluctuations in PO₄-P (mg/L) values were similar to NO₃-N. The minimum (0.12 mg/L) and maximum (18.92 mg/L) PO₄-P levels were observed during summer months with a mean value of 5.30 mg/L (Table 2). The seasonal mean values of PO₄-P(3.73 mg/L) during winter and summer were 3.00 and 5.3 mg/L, respectively as shown in Table. 2.

The E.C values in this study range from 1.14-4.07 dS/m.s while the EC values for natural water are between 0.5 – 1.5 dS/m (Gems 1997) , that is why Shatt AL-Arab is considered as a saline water in comparison with other rivers of the world. Previous studies show that EC. range from 1.469-2.364 dS/m during 1980 (Polservice 1980) .while it range from 2.282-2.510 dS/m during 2004 (Al-Mayahi 2005). 1.53-5.3 dS/m (Al-Kazaeh 2014) ,1.57-10.85 dS/m Al-Hejuje (2015).

In stream water, dissolved solids consist of calcium, chlorides, nitrate, phosphorus, iron, sulfur, and other ions particles that will pass through a filter with pores of around 2 microns (0.002 cm) in size. Suspended solids include silt and clay particles, plankton, algae,

Table 2 Seasonal variations of physical and chemical parameters in Shatt AL-Arab Estuary

	Station	Summer	Autumn	Winter	Spring		Station	Summer	Autumn	Winter	Spring
Water temp-rature (°C)	1	30.00	19.00	9.00	22.00	Ca	1	104.00	152.00	226.60	128.00
	2	31.00	18.00	10.00	23.00		2	128.00	125.30	154.60	109.30
	3	32.00	20.00	10.00	23.00		3	88.00	141.30	157.30	106.60
	4	33.00	21.00	11.00	24.00		4	112.00	101.30	181.30	98.60
	5	33.00	22.00	12.00	24.00		5	104.00	125.30	210.60	112.00
	6	34.00	22.00	14.00	25.00		6	114.60	144.00	173.30	98.60
D.O mg/L	1	5.30	6.30	9.30	8.40	Mg	1	50.58	217.18	304.56	50.22
	2	5.90	7.80	8.50	8.30		2	45.31	93.96	445.86	55.08
	3	6.20	6.50	8.70	7.60		3	63.36	176.58	437.76	56.70
	4	6.80	6.80	8.10	8.50		4	36.77	165.20	289.98	69.66
	5	6.90	7.90	8.30	8.90		5	58.34	126.36	233.28	57.22
	6	7.10	8.10	8.50	8.40		6	81.62	105.26	265.68	110.16
PH	1	7.99	7.33	8.21	8.41	Cl	1	368.68	1035.51	673.55	335.71
	2	7.48	7.20	7.98	8.18		2	626.28	1059.46	508.11	524.66
	3	7.99	7.33	8.10	8.16		3	337.93	572.45	496.30	371.04
	4	7.98	7.36	8.32	8.29		4	268.50	624.41	555.38	333.68
	5	7.95	7.36	8.55	8.24		5	361.92	551.46	590.83	354.50
	6	7.96	7.43	8.36	8.30		6	363.95	539.35	562.08	369.70
E.C mS/cm	1	1.58	2.26	2.71	1.90	HCO3	1	153.60	272.46	528.66	183.00
	2	2.72	2.57	2.85	2.00		2	186.14	250.00	597.80	195.20
	3	1.72	2.42	2.72	1.95		3	138.96	252.13	536.80	174.80
	4	1.75	2.43	2.80	1.98		4	133.30	207.40	553.06	166.73
	5	1.74	2.40	2.88	1.96		5	152.02	227.70	488.00	154.53
	6	1.73	2.42	2.84	1.98		6	149.57	223.60	622.20	158.60
TUR FTU	1	60.60	27.60	14.20	15.32	SO4	1	208.63	431.55	1921.88	251.50
	2	41.60	38.30	13.70	12.39		2	276.77	485.49	1852.89	230.75
	3	32.50	72.30	9.06	16.28		3	295.40	474.63	506.46	271.49
	4	27.49	55.60	9.50	6.05		4	509.66	538.71	545.62	305.49
	5	41.90	36.60	7.80	7.77		5	658.19	430.01	618.49	253.40
	6	47.14	38.30	9.20	9.16		6	807.60	456.03	593.56	337.28
TSS mg/l	1	100.00	68.60	38.00	55.30	NO3	1	24.34	9.93	31.04	53.57
	2	59.60	64.60	34.00	83.30		2	20.07	8.89	24.78	54.01
	3	61.60	73.30	36.00	46.00		3	48.23	11.83	25.13	55.50
	4	98.60	92.60	30.60	39.30		4	57.43	9.91	23.90	54.19
	5	41.30	79.30	30.60	56.60		5	49.85	9.75	26.68	51.74
	6	78.60	57.30	28.60	64.60		6	46.13	10.35	28.15	53.85
TDS mg/l	1	1432.00	2222.00	1640.60	1290.00	PO4	1	3.03	2.16	2.29	2.12
	2	1740.00	2057.30	1616.00	1978.00		2	13.45	5.63	4.41	5.19
	3	1082.00	1433.30	1852.60	1976.00		3	5.11	2.62	3.12	3.90
	4	1046.00	1677.30	1697.30	2092.00		4	4.45	2.87	2.37	2.30
	5	1062.00	1925.30	1873.30	2918.60		5	3.67	1.77	2.93	3.34
	6	1034.00	1885.30	1740.00	2538.60		6	2.06	3.79	2.91	3.12

fine organic debris, and other particulate matter. These are particles that will not pass through a 2-micron filter. Higher concentrations of suspended solids can serve as carriers of toxics, which readily cling to suspended particles. A high concentration of total solids will make

drinking water unpalatable and might have an adverse effect on people who are not used to drinking such water. Levels of total solids that are too high or too low can also reduce the efficiency of wastewater treatment plants, as well as the operation of industrial processes

that use raw water. Total solids also affect water clarity. Higher solids decrease the passage of light through water, thereby slowing photosynthesis by aquatic plants. Water will heat up more rapidly and hold more heat; this, in turn, might adversely affect aquatic life that has adapted to a lower temperature regime. Sources of total solids include industrial discharges, sewage, fertilizers, road runoff, and soil erosion. TDS range from 900-5965 mg/L during (1975-2000) (Hasan and Mahmood 2003) for Shatt AL-Arab water and within the study period from (142 and 5566.00) in Al Fao and paper factory respectively. High concentrations of TDS in spring and autumn seasons as compared with that in the winter season may be due to the addition of solids from runoff water and sewage effluents which increase the TDS values in the river. This result was in agreement with Hassan et al. (2010) and Al-Hejuje (2015) who found that TDS values in the Euphrates and Shatt Al-Arab river respectively increased in winter months and decreased in summer months.

The total suspended solids (TSS) values range between (11-18 mg/L) during (1978) which considered as a low value that may be because of the sedimentation process in marshes. These result can be observed during the study period, the mean value of TSS (66.25, 60.42, 54.25, 65.33, 52, 57.33) mg/L for six station respectively. value in Tigris (turbid water) 64 mg/L and in Al-Qurna location is 13 mg/L rising in Sibah to 44 mg/L and 147 mg/L in Al-Fao this location effected with Karone sediment and sea tide.

Higher turbidity increases water temperatures because suspended particles absorb more heat. This, in turn, reduces the concentration of dissolved oxygen (DO) because warm water holds less DO than cold. Higher turbidity also reduces the amount of light penetrating the water, which reduces photosynthesis and the production of DO. Suspended materials can clog fish gills, reducing resistance to disease in fish, lowering growth rates, and affecting egg and larval development. As the particles settle, they can blanket the stream bottom, especially in slower waters, and smother fish eggs and benthic macro invertebrates. Sources of turbidity include: Soil erosion, Waste discharge Urban runoff, eroding estuary banks, Large numbers of bottom feeders (such as carp), which stir up bottom sediments, excessive algal growth. One of the important light parameter which is of well relation with

clearness of water that considered as an indication to suspended mater in water , it values range between (0.80-163.00) FTU for study area Shatt Al-Arab river has high turbid water which exceeded the permissible value (5 NTU) in aquatic ecosystems.

Calcium & Magnesium (Ca , Mg) elements exist widely in water. In natural water Ca concentration range between (10 – 100)mg/L causing water hardness. Mg concentration Range between (5-50)mg/L , and if exceed 125 mg/L causing diarrhea and also softening hardness. during (1968-2000) (Hasan and Mahmood , 2003). Ca concentration ranges from 136 mg/L to 160 mg/L Mg concentration Range from 96 mg/L to 223 mg/L While the range between (212.1-280) mg/L for Ca and (67.4-691.4) mg/L for mg during 2004 (Al- Mayah, 2004). In this study Ca concentration range from 32.00 to 360 mg/L while Mg concentration range from 14.58 to 899 mg/L .Ca and mg concentrations Considered as a high value but they are still within the standard of WHO (2011), in Shatt Al-Arab river.

Bicarbonate (HCO_3) Alkalinity of the most natural and treated water are Ca , mg bicarbonate when pH degree for this water does not exceed 8.3. during (1988-2000) HCO_3 conc. Ranges (183-244) mg/L and during 2004 it ranges (133.8-131.2)mg/L. in this study its conc. Range (53.46-976) mg/L. this rising in HCO_3 concentration Is accompanied by rising in pH and Ca, Mg conc. That refer to an expected in the chemical constitution of Shatt Al-Arab water which may be a result of water pollution.

High concentration of Cl is caused by the geological components which contain chloride. Except that high concentration of Cl is an indication for high pollution of sewage water , some industrial waste , because of tide see or any saline water which have a bad effect on modern pipes, agriculture and factories. during (1988 – 2001) Cl range between (137 – 1060) mg/L ,and during the study period range (241.06-2056.6) mg/L for Qurna and Al-Ashar and that refer to the increase of industrial waste in river water (Table 1)

Nitrates are a form of nitrogen, which is found in several different forms in terrestrial and aquatic ecosystems. These forms of nitrogen include ammonia (NH_3), nitrates (NO_3), and nitrites (NO_2). Nitrates are essential plant nutrients, but in excess amounts they can cause significant water quality problems. Together

with phosphorus, nitrates in excess amounts can accelerate eutrophication, causing dramatic increases in aquatic plant growth and changes in the types of plants and animals that live in the stream. This, in turn, affects dissolved oxygen, temperature, and other indicators. Excess nitrates can cause hypoxia (low levels of dissolved oxygen) and can become toxic to warm-blooded animals at higher concentrations (10 mg/L or higher) under certain conditions. It is the most oxidized form of nitrogen, it comes from chemical fertilizers, agriculture lands, industrial water and sewage water. Nitrate estimation helps in the observation of the form and oxidation degree in stream and ground water. In natural water (unpolluted) contains low nitrate, in case of increasing its concentration. Causes (methaemoglobinemia) for young children and also causes Eutrophication; which causes secondary pollution.

Nitrate concentration. Range (3.2-110.12) mg/L during study period which considered higher values compared to the study result during 2004 which ranges from (3 -10) mg/L in Shatt AL-Arab water. Results found that most of downstream stations of Shatt AL-Arab river showed higher nitrogen compound compared with the upstream stations because they are more impacted by pollutants from domestic sewage released from Basrah city center. Also, climatic changes as well as reduction in water inflow to the river have resulted in alteration in nitrogen compounds concentrations, this conclusion was in agreement with Hameed and Aljorany (2011) and Al-Hejuje (2015).

Phosphorus (PO_4) has a complicated story. Pure, "elemental" phosphorus (P) is rare. In nature, phosphorus usually exists as part of a phosphate molecule (PO_4). Phosphorus in aquatic systems occurs as organic phosphate and inorganic phosphate. Organic phosphate consists of a phosphate molecule associated with a carbon-based molecule, as in plant or animal tissue. Phosphate that is not associated with organic material is inorganic. Inorganic phosphorus is the form required by plants. Animals can use either organic or inorganic phosphate. Both organic and inorganic phosphorus can either be dissolved in the water or suspended (attached to particles in the water column). It comes from the weathering and process on pH degree. When pH range (6-8) the boundary forms 10% $H_2PO_4^{-1}$ and 90% HPO_4^{-2} this form is the main nutrient for organisms. Phosphate compounds more in the natural and sewage

water and cause a secondary pollution that makes basic nutrition for increasing organisms growth which depends on light with undesirable amounts some times. Phosphate concentration range (0.07 -3.67) mg/L during (2004) and in Al-Qurna reach (0.193) mg/L to (5.15) mg/L for Tigris before joining Euphrates. The concentrations were range from 0.12 mg/L at station 1 to 18.92 at station 2. The decrease of phosphorus concentrations, which is more distant from the discharge points of effluents, may be due to the self-purification capacity of the receiving stream. Nitrates and Phosphates appear with gradual increases especially during the months of April and May when fertilizers in lands around the river are used and accidentally spill into water.

Sulfate (SO_4) is formed from earth crust and organic compounds containing sulfate, which come from the industrial waste its concentration. Range from several grams to some thousand grams. WHO considered (400) mg/L as the highest limit for drinking water. In Shatt Al-Arab water sulfate concentration. Ranges from (80.15- 4462.23) mg/L being the lowest in Al-Sebah and the highest in paper factory. While it reached (674) mg/L for Shatt Al-Arab water during 2004 (Al-Mayahi, 2005). and High concentrations of sulfate recorded in Winter may be due to high petroleum compounds used as fuel and released sulfate compounds to the environments, or may be due to wet precipitation during Winter season. High concentration of sulfate may be due to electrical power plants (Al-Najebbia power plant and Turkish power plant) that released sulfate compound to the environment, also the agricultural activities at these areas have a role in the increasing sulfate compounds (Al-Hejuje, 2015).

As conclusion results showed some fluctuation in some of the data during different seasons at different locations of the estuary, also the results indicate that the water is Non-polluted and can be used for Domestic, Irrigation and Fisheries.

References

- Al-Hejuje M.M.,2015," Application of Water Quality and Pollution Indices to Evaluate the Water and Sediments Status in the Middle Part of Shatt Al-Arab River" . Ph.D. Thesis . University of Basrah, college of science, pp.214
- Al-Kazaeh D. K. K.,2014, Chemical and Physical properties of common water in area and evaluation degree for Irrigation in Basra , Journal of Basra Research (Scientific) N.40 ,Part 2,B.
- Al-Mayahi A .,2004, Salinity balance of tided flow Shatt Al-Arab water way. Basra city. Summary report.
- Al-Mayahi M.A.T.,2005, " Concentration of total hydrocarbons in some region of Shatt Al-Arab River and its effected on benthic algae ".M .Sc

- Thesis , Fisheries and Marine Resources Department , College of Agriculture - University of Basrah , 61 pp.
- APHA, American Public Health Association .,2005 , "Standard methods for the examination of water and wastewater" , 21st Edition .Washington ,DC. USA.
- Basavaraja Simpi S.M., Hiremat K.N., Murthy K.N., Chandras H., Anil N., Patel E., and Puttiah., 2011, Analysis of Water Quality Using Physico-Chemical Parameters Hosahalli Tank in Shimoga District, Karnataka, India Global Journal of Science Frontier Research, 11(3):15-25
- Gems, 1997 , Water operational guide. Third edition prepared under the joint sponsorship of UNEP. WHO; UNESCO.WMO
- Hameed A.H., and Aljorany Y.S.,2011, Investigation on nutrient behavior along Shatt Al-Arab river , Basrah , Iraq , J. Appl. Sci. Res.,7(8): 1340-1345
- Hasan K. H., and Mahmood A., 2003, The hydrochemical factors in Shatt Al-Arab River (Incidents of conference to water Transport Company). Baghdad 21-22.Oct
- Hassan F.M., Saleh M.M., and Salman J.M., 2010," A study of physicochemical parameters and nine heavy metals in the Euphrates River , Iraq ," E- Journal of Chemistry ,7(3):685-692
- <http://dx.doi.org/10.1155/2010/906837>
- Hussain N. A., Al-Saboonich A.,Al-Saad H.T., Karem H., and Hamid I., 1991, Shatt Al-Arab scientific study. Univ. of Basra. Marin science Center, Publication. No. (10)
- Parson T.S., Mita Y., and Lall G.M., 1984, Manual of chemical and biological methods for sea water analysis. Pergamon Press, Oxford.
- PolSERVICE, 1980, Shatt Al-Arab project. Feasibility report Draft. Vol.1 part A, republic of Iraq. Ministry of Irrigation. General Establishment for studies and design. 11 sp. 29 fig. 35 tab.
- Tabari H., Safar M., and Mohammad A., 2011, Long-term variations of water quality parameters in the Maroon River, Iran. Environment. Monitoring Assessment , 177:273-287
- <http://dx.doi.org/10.1007/s10661-010-1633-y>
- UN-ESCWA and BGR (United Nations Economic and Social Commission for Western Asia; Bundesanstalt für Geowissenschaften und Rohstoffe), 2013, Inventory of Shared Water Resources in Western Beirut. CHAPTER 5 - SHATT AL ARAB, KARKHEH AND KARUN RIVERS.
- WHO: World Health Organization, 2011, "Guidelines for Drinking-Water Quality". 4th edition . Geneva 27, Switzerland. (<http://www.who.int>)



Reasons to publish in BioPublisher *A BioScience Publishing Platform*

- ★ Peer review quickly and professionally
- ☆ Publish online immediately upon acceptance
- ★ Deposit permanently and track easily
- ☆ Access free and open around the world
- ★ Disseminate multilingual available

Submit your manuscript at: <http://biopublisher.ca>

