

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281370843>

Spatial Analysis on the Concentrations of Air Pollutants in Basra Province (Southern Iraq)

Article in *Open Journal of Air Pollution* · September 2015

DOI: 10.4236/ojap.2015.43013

CITATIONS

10

READS

353

7 authors, including:



Shukri I Al-Hassen
University of Basrah

49 PUBLICATIONS 40 CITATIONS

SEE PROFILE



Abdul Wahab
Superior University

45 PUBLICATIONS 293 CITATIONS

SEE PROFILE



Adnan Ateeq
Southern Technical University -Iraq

23 PUBLICATIONS 28 CITATIONS

SEE PROFILE



Hamid T. Al-Saad
University of Basrah

255 PUBLICATIONS 1,000 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



FLOW INJECTION ANALYSIS (FIA) IN IRAQ [View project](#)



Climate Change [View project](#)

Spatial Analysis on the Concentrations of Air Pollutants in Basra Province (Southern Iraq)

Shukri I. Al-Hassen¹, Abdul Wahab A. Sultan², Adnan A. Ateek², Hamid T. Al-Saad^{3*}, Salah Mahdi³, Abdulzahra A. Alhello³

¹Department of Geography, University of Basra, Basra, Iraq

²Technical College, Southern Technical University, Basra, Iraq

³Department of Environmental Chemistry, University of Basra, Basra, Iraq

Email: *htalsaad@yahoo.com

Received 15 July 2015; accepted 28 August 2015; published 31 August 2015

Copyright © 2015 by authors and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

This paper aims to analyze the geographic distribution of air pollutant concentrations in Basra Province, Southern Iraq, and to cartographically determine the spatial variation of air pollution levels as well as to recognize the hottest spots of air pollution within the study area, and conclude that the levels of air pollution in the study area are spatially varied, with an irregular spatial pattern and some hotspots.

Keywords

Air Pollution, Air Pollutants, Gaseous Emission, Spatial Analysis, Basra

1. Introduction

Air pollution may be defined as any atmospheric condition in which certain substances are present in such concentrations that they can produce harmful effects on man and his environment. An air pollutant, however, is any gas or substance (such as SO_x, NO_x, CO, and HCs) or particulate matter (such as smoke, dust, fumes, and aerosols) that leads to ambient air contamination. A pollutant may originate from natural or anthropogenic sources, or both. Pollutants occur throughout much of the troposphere; however, pollution close to the earth's surface within the boundary layer is of most concern because of the relatively high concentrations resulting from sources at the surface [1] [2].

Air pollutant concentrations depend mainly on the total mass of pollution emitted into the atmosphere, to-

*Corresponding author.

gether with the atmospheric conditions that affect its fate and transport. Obviously, air pollution has many and varied sources, including cars, smokestacks, and other industrial inputs into the atmosphere as well as wind erosion of soil. Large emissions from both anthropogenic and natural sources over long periods enhance concentrations, as do the chemical and physical properties of these pollutants. For example, when nitrogen oxides and hydrocarbons in car exhaust are emitted into warm, sunlit air, they readily form ozone molecules (O_3). Similarly, the solubility of a pollutant affects how efficiently it is removed by rainfall. In addition, atmospheric conditions have a major effect upon pollutants once these pollutants are emitted into (e.g., nitrogen oxides from car exhaust) or formed within (e.g., O_3) the atmosphere. Pollution dispersal is controlled by atmospheric motion, which is affected by wind, stability, and the vertical temperature variation within the boundary layer. Stability, in turn, influences both air turbulence and the depth at which mixing of polluted air takes place ([2]; see also [3]-[7]).

In the study area (Basra province), located in southern Iraq, air pollution is of major public concern, which is currently the object of extensive scientific research. Studies of Al-Asadi [8], Al-Mayahi [9], Al-Imarah *et al.* [10], Al-Saad *et al.* [11], Garabedian [12], Al-Hassen [13], Qassim [14], Douabul *et al.* [15], Sultan *et al.* [16], Karmalla *et al.* [17], Abdullah and Hussien [18], and Al-Hassen *et al.* [19], are an example of some local-scale research in this respect.

The study area is rich in petroleum and many existing industrial and human activities as the main sources of gaseous emissions around it (Figure 1). These are emission sources to contaminate the ambient air in Basra. According to above mentioned studies, Basra was recorded, in the last two decades, high quantities of gaseous emissions causing elevated levels of outdoor air pollution, and that concentration of some air pollutants was at risk to the public health. One of the most important driving forces is atmospheric conditions in this region.



Figure 1. Photography of some gaseous emission sources in the study area. (a) Petrochemical plant; (b) Najybia power station; (c) Emissions of petroleum exploitation near Burchesya; (d) Iranian Abadan refinery offshore seeba.

The previous studies, however, are focused on values of gaseous pollutant concentration and its implications on the human health in the study area; thus the objective of the present study is to spatially analyze the geographic distribution of air pollutant concentrations in Basra, and to cartographically determine the spatial variation of air pollution levels on the map of this region. The major aim is to recognize the hottest spot of air pollution within whole areas of Basra province.

Basra has a hot desert climate (Köppen climate classification *BWh*), like the rest of the surrounding region, though it receives slightly more precipitation than inland locations due to its location near the coast. During the summer months, from June to August, Basra is consistently one of the hottest cities on the planet, with temperatures regularly exceeding 40°C (104°F) and approaching 45°C (113°F) in July. In winter Basra experiences mild weather with average high temperatures around 20°C (68°F). On some winter nights, minimum temperatures are below 0°C (32°F). High humidity—sometimes exceeding 90%—is common due to the proximity to the marshy Persian Gulf [20].

2. Materials and Methods

As shown in **Figure 2**, seventeen sampling stations were chosen in the study area. The selected stations divided to cover the eastern and western region within the study area, and the geographic distribution of sampling stations was taken in consideration in the vicinity of human settlements and involved a variety of local environments (see **Table 1**).




Figure 2. Map of the study area (Basra Province, Southern Iraq), showing the selected sampling stations. Note: Numbers of sampling stations correspondences their names listed in **Table 1** and **Table 2**.

Table 1. Type of environment in the selected sampling stations, and the major sources of gaseous emissions in the study area.

No.	Sampling station	Type of environment	Major source of air pollution
1	Midaina	Agricultural/industrial/urban	Petroleum exploitation and urban contaminates
2	Qurna	Agricultural/industrial/urban	Petroleum exploitation and urban contaminates
3	Dayer	Agricultural/industrial/urban	Petroleum exploitation and urban contaminates
4	Garmatt Ali	Agricultural/urban	Power station
5	Ashar	Urban	Urban contaminates
6	Abo Khaseeb	Agricultural/urban	Urban contaminates
7	Seeba	Agricultural/industrial	Petroleum industry and exploitation
8	Faw	Agricultural/urban	Urban contaminates
9	Ras Absha	Marine	Natural contaminates
10	Shatt Al-Basra	Riverine	Natural and urban contaminates
11	Burchesya	Industrial	Petroleum industry and exploitation
12	Petrochemical plant area	Industrial	Petroleum industry
13	South gas plant area	Industrial	Petroleum industry
14	Safwan	Urban	Urban contaminates
15	Khor Al-Zubayer	Industrial/urban	Industrial and urban contaminates
16	Gas terminal	Industrial/marine	Petroleum industry and shipping
17	Umm Qasr	Urban/marine	Urban contaminates

The selected stations were in order to monitor the concentrations of gaseous pollutants released into ambient air of the study area. A variety of gaseous pollutants such as carbon monoxide (CO), carbon dioxide (CO₂), sulfate oxides (SO_x), nitrogen oxides (NO_x), ozone (O₃), petroleum hydrocarbons (HCs), methane (CH₄), hydrogen sulfide (H₂S), and formaldehyde (HCHO), measured in this work. Concentrations of HCs, NO_x, SO_x, HCHO, O₃, and CO₂ measured using the portable detection instrument of Drager Chip-Measurement System, Germany, whereas the portable instrument of RK1 Gas Monitoring Eagle II, USA, detected the pollutants of CH₄, H₂S, and CO.

Fieldwork carried out during a one daytime of spring 2015, and all measurements were done at the same time in purpose of obtaining on readings approaching to reality. Therefore, the working team divided into two groups, one of them work in the eastern region (Midaina, Qurna, Dayer, Garmatt Ali, Ashar, Abo Khaseeb, Seeba, Faw, and Ras Absha), while the other group work in the western region (Shatt Al-Basra, Burchesya, Petrochemical plant area, South Gas Plant area, South Gas Plant area, Safwan, Khor Al-Zubayer, Gas terminal, Umm Qasr). The procedure of measuring was as described by Douabul *et al.* [15].

3. Results and Discussion

Table 2 lists the obtained results for concentrations of some air pollutants in the given study area and period. These concentrations recorded from the direct readings displayed on the screen of both employed detectors. The values were adjusted in a statistic form to make a more geographic mode. Thus, the values have been graphically and cartographically represented in **Figure 3** and **Figure 4**, respectively. In this study, data analysis and explanation will conducted in the terms of each given element with the emphasis on spatial variation and geographic distribution of air pollution, as follows:

1) Carbon monoxide (CO) is a colorless, odorless, and tasteless gas that is slightly less dense than air. It is toxic to humans when encountered in concentrations above about 35 ppm. In the atmosphere, it is spatially variable and short lived, having a role in the formation of ground-level ozone [21].

Table 2. Concentrations of air pollutants measured in the study area based on the selected sampling stations, 2015.

No.	Sampling station	CO ppm	CO ₂ ppm	NO _x ppm	SO _x ppm	H ₂ S ppm	HCs ppm	CH ₄ ppm	HCHO ppm	O ₃ ppb	Index*
1	Midaina	4.16	215.12	0.72	0.54	1.53	1.62	8.52	0.42	0.02	25.85
2	Qurna	6.25	286.45	0.83	0.64	1.82	1.93	8.86	0.35	0.03	34.12
3	Dayer	8.92	260.21	0.86	0.63	1.92	3.25	10.21	0.46	0.02	31.83
4	Garmatt Ali	10.23	280.38	0.95	1.25	1.98	5.28	9.52	0.63	0.06	34.47
5	Ashar	12.32	225.32	0.65	0.92	1.21	12.21	13.28	0.92	0.12	29.66
6	Abo Khaseeb	20.63	250.12	0.83	1.28	1.26	24.28	14.25	1.23	0.14	34.89
7	Seeba	30.23	280.11	1.45	2.28	2.68	31.23	25.68	1.86	0.23	41.63
8	Faw	10.24	240.61	0.79	1.68	2.06	22.52	13.34	0.66	0.06	32.44
9	Ras Absha	2.52	180.32	0.42	0.43	1.25	1.74	5.54	0.24	0.01	21.38
10	Shatt Al-Basra	10.68	210.11	0.65	0.72	1.12	7.25	10.11	0.72	0.04	26.82
11	Burchesya	40.23	310.27	4.25	10.23	6.24	30.21	22.65	1.52	0.12	69.30
12	Petrochemical plant area	16.23	200.10	1.31	1.65	1.32	10.53	13.21	0.31	0.06	27.19
13	South gas plant area	20.53	220.31	1.45	3.21	3.1	18.23	10.25	0.62	0.16	30.87
14	Safwan	18.22	210.53	0.93	1.24	2.5	10.82	9.34	0.52	0.13	28.24
15	Khor Al-Zubayer	14.28	226.3	1.23	2.52	1.8	11.23	9.93	0.68	0.09	29.78
16	Gas terminal	18.34	228.23	1.86	4.38	2.1	22.31	16.83	0.84	0.11	32.77
17	Umm Qasr	16.12	180.98	0.98	3.49	3.5	24.63	12.46	0.75	0.07	26.99
	Mean	15.30	235.61	1.15	2.18	2.19	14.01	12.58	0.74	0.08	31.49

Data based on Fieldwork. *Index means a sum of values for the selected parameters divided by its number. It may be expressed, in the other meaning, the intensity of pollution.

As listed in **Table 2** and **Figure 3** & **Figure 4(a)**, the Burchesya sampling station records the highest value of CO is 40.23 ppm, while the lowest is 2.52 ppm in Ras Abasha station. The mean concentration of CO is 15.30 ppm within the all sampling stations.

2) Carbon dioxide (CO₂) is a colorless, odorless gas vital to life on Earth. Carbon dioxide exists in the Earth's atmosphere as a trace gas at a concentration of about 0.04 percent (400 ppm) by volume. It is present in deposits of petroleum oil and natural gas [22].

Table 2 and **Figure 3** & **Figure 4(b)** report that the maximum concentration of CO₂ is 310.27 ppm recorded in the Burchesya sampling station, while the minimum concentration is 180.32 ppm in the Ras Abasha station. In general, the sampling stations in the eastern region of the study area registers values higher than those that in the western stations. The mean concentration of CO₂ is 235.61 ppm.

3) Nitrogen oxide (NO_x) is a prominent air pollutant; it may refer to a binary compound of oxygen and nitrogen, or a mixture of such compounds. This reddish-brown toxic gas has a characteristic sharp, biting odor and is a prominent air pollutant [23].

Table 2 and **Figure 3** & **Figure 4(c)** indicate that the highest concentration of NO_x registered in the study area was 40.23 at the Burchesya sampling station, this may be a record value to compare with given in the previous studies yet. The lowest concentration was 0.42 in the Ras Abasha station. The mean concentration of NO_x is 1.15 ppm.

4) Sulfur oxide (SO_x) refers to many types of sulfur and oxygen containing compounds such as SO, SO₂, SO₃, S₇O₂, S₆O₂, S₂O₂, etc. SO_x is a toxic gas with a pungent, irritating, and rotten smell [24].

In this study, as shown in **Table 2** and **Figure 3** & **Figure 4(d)**, the maximum concentration of SO_x was 10.23 ppm in the Burchesya sampling station, while the minimum concentration is 0.42 ppm in the Ras Abasha station. The mean concentration of SO_x is 2.18 ppm.




Figure 3. Graphic representation of the concentrations of air pollutants measured in the study area. Data based on Table 2.




Figure 4. Cartographic representation of the concentrations of air pollutants measured in the study area. Data based on Table 1.

5) Hydrogen sulfide is the chemical compound with the formula (H_2S). It is a colorless gas with the characteristic foul odor of rotten eggs; it is heavier than air, very poisonous, corrosive, flammable, and explosive [25].

Table 2 and Figure 3 & Figure 4(e) refer to the highest value of H_2S recorded in this study was 6.24 ppm in the Burchesya sampling station, while the lowest value was 1.21 ppm in the Ashar station. The mean concentration is 2.19 ppm.

6) Hydrocarbon (HCs) is an organic compound consisting entirely of hydrogen and carbon. Hydrocarbon poisoning such as that of benzene and petroleum usually occurs accidentally by inhalation or ingestion of these cytotoxic chemical compounds [22].

HCs gas concentrations in the study area were high, in general. The maximum concentration is 31.23 ppm registered at the Seeba sampling station, whereas the minimum concentration is 1.62 in the Midaina station. The mean concentration of HCs is 14.01 ppm, as shown in **Table 2** and **Figure 3** & **Figure 4(f)**.

7) Methane is a chemical compound with the chemical formula (CH_4). It is the simplest alkane and the main component of natural gas. Methane is not toxic, yet it is extremely flammable and may form explosive mixtures with air [26].

The highest values of CH_4 , as indicated in **Table 2** and **Figure 3** & **Figure 4(g)**, were centered in places with the petroleum industry and exploitation within the study area. The maximum concentration, however, is 25.68 ppm in the Seeba station, while the minimum concentration is 5.54 ppm in the Ras Abasha station. The mean concentration of CH_4 is 12.58 ppm.

8) Formaldehyde (HCHO) is a colorless, highly toxic, and flammable gas at room temperature that is slightly heavier than air. It has a pungent, highly irritating odor that is detectable at low concentrations, but may not provide adequate warning of hazardous concentrations for sensitized persons [27].

Table 2 and **Figure 3** & **Figure 4(h)** show that the concentration of HCHO records the maximum value is 1.86 ppm in the Seeba station, while the minimum value is 0.24 ppm recorded in the Ras Abasha station. The mean concentration of HCHO is 0.74 ppm.

9) Ozone is an inorganic molecule with the chemical formula (O_3). It is a pale blue gas with a distinctively pungent smell. This same high oxidizing potential, however, causes ozone to damage mucous and respiratory tissues in animals, and also tissues in plants, above concentrations of about 100 ppb. This makes ozone a potent respiratory hazard and pollutant near ground level [28].

In the study area, as shown in **Table 2** and **Figure 3** & **Figure 4(i)**, the concentrations of ground level ozone (O_3) were largely varied within the selected sampling stations. The highest value is 0.23 ppb in the Seeba station, whereas the Ras Abasha station register the lowest value is 0.01 ppb. Thus, the mean concentration of O_3 is 0.08 ppb.

In general, the concentrations of air pollutants registered in the study area were spatially varied, it seems somehow a random pattern of distribution in the terms of each pollutant (see **Figure 4**). However, an overall spatial pattern may be drawn by using the index of pollution showing in **Figure 5** based on **Table 2**. This index is a result of summing all values of each element dividing by its numbers, the resulting value is an approximate indicator of the spatial concentration of a pollutant. To simplify explaining the causes of spatial variation in pollutant concentrations, the mentioned **Table 1** lists the major gaseous emission sources affecting air quality in the study area.




Figure 5. Geographical distribution of the levels of air pollution in the study area according to index of pollution. Data based on **Table 2**. Data based on **Table 1**.

As shown in **Figure 5(a)** & **Figure 5(b)**, the elevated levels of air pollution focuses on the western region of the study area, this is due to existing industrial pollution sources. The sampling stations of Qurna, Dayer, and Garmatt Ali also report a significant index values with 34.12, 31.83 and 34.47 respectively, this refers to the petroleum exploitation existing in these places as well as the high values of CO₂, which, in turn, increase the index value in a comparison with the other sampling stations.

It is clear that, however, the highest values of the air pollution index with 69.30 and 41.63 have been recorded in the sampling stations of Burchesya and Seeba, respectively. For the former, this is due to the gaseous pollutants emitted from Abadan refinery offshore this site, in addition to there are a newer exploitation of petroleum within it; for the latter, this increase related to the gaseous emissions from dense works of petroleum exploitation on this site. The sampling station of Ras Abasha was the less polluted with an index value of 21.38, because of its marine nature and relatively distant from the influence of anthropocentric pollution sources.

4. Conclusions

In accordance to the spatial analysis of the geographical distribution of air pollutants, the present study concluded that the levels of air pollution in Basra province (the study area) were spatially varied, and that this spatial variation consequently correlated with the geographical distribution of gaseous emission sources existing in the study area. However, there is an irregular spatial pattern of air pollution concentration, although of some hotspots such as Burchesya and Seeba.

The authors recommend that mitigation to air pollution sources, particularly from stationary emission sources in the study area, is urgent action. Moreover, the installing of fixed stations designated to the air pollution monitoring is so necessary. Nevertheless, the accurate assessment of air pollution status in the study area needs more research and monitoring.

Acknowledgements

The authors acknowledge the following labs for the technical assistants: Environmental Analysis and Research Lab in Department of Geography, College of Arts, University of Basra; Gas Chromatography Lab in Department of Environmental Chemistry, Marine Science Center, University of Basra, and Air Pollution Lab in Department of Environmental and Pollution Engineering, Technical College, Southern Technical University.

References

- [1] Admassu, M. and Wubeshet, M. (2006) Air Pollution: Lecture Notes for Environmental Health Science Students. University of Gondar Publications, Ethiopia, 5-6.
http://www.cartercenter.org/resources/pdfs/health/ephti/library/lecture_notes/env_health_science_students/AirPollution.pdf
- [2] Matthias, A.D., Comrie, A.C. and Musil, S.A. (2006) Atmospheric Pollution. In: Peeper, I.L., Gerba, C.P. and Brusseau, M.L., Eds., *Environmental and Pollution Science*, 2nd Edition, Elsevier, San Diego, 377-394.
- [3] Vallero, D.A. (2008) Fundamentals of Air Pollution. 4th Edition, Elsevier Inc., London, 3.
- [4] Spellman, F.R. (1999) The Science of Environmental Pollution. Taylor & Francis Routledge, Pennsylvania, 245.
- [5] Harrison, R.M. (2001) Air Pollution: Sources, Concentrations and Measurements. In: Harrison, R.M., Ed., *Pollution: Causes, Effects and Control*, 4th Edition, RSC, Cambridge, 169-192. <http://dx.doi.org/10.1039/9781847551719-00169>
- [6] Gittins, M.J. (1999) Air Pollution. In: Bassett, W.H., Ed., *Clay's Handbook of Environmental Health*, 18th Edition, E & FN Spon, London, 729-776. <http://dx.doi.org/10.4324/9780203016312.ch42>
- [7] Christoforou, C. (2004) Air Pollution. In: Stapleton, R.M., Ed., *Pollution A to Z*, Vol. 1, Macmillan Reference, New York, 30-38.
- [8] Al-Asadi, K.A.W. (1998) The Influence of Climatic Factors on the Major Industries in Basra and Their Reflections on the Environmental Pollution. Ph.D. Thesis, College of Arts, University of Basra, 197. (In Arabic)
- [9] Al-Mayahi, I.K. (2005) An Environmental Analysis on the Factors Affecting the Air Pollutants Quality at Basra Province. M.A. Thesis, College of Education, University of Basra, 240. (In Arabic)
- [10] Al-Imarah, F.J.M., Al-Mohameed, R.S.J. and Ibraheem, S.I. (2007) Extent of Atmospheric Pollution by Some Industrial Emissions Released from Petrochemicals and Gas Liquefier Industries in Khor Al-Zubair. *Journal of Kerbala, Special Issue on the Annual Environment of Meeting of Babylon University*, 1-6.

- [11] Al-Saad, H.T., Al-Imarah, F.J.M., Hassan, W.F., Jasim, A.H. and Hassan, I.F. (2010) Determination of Some Trace Elements in the Fallen Dust on Basra Governorate. *Basrah Journal of Science*, **28**, 243-252. <http://www.iasj.net/iasj?func=fulltext&aId=55085>
- [12] Garabedian, S.A.K. (2010) Study of the Main Pollutants of Air Caused by Transportation in Basra. *Proceedings of the Scientific Conference of Marine Science Center and the National Specialized Workshop on Oceanography*, 23-24 December 2008, 125-136. (In Arabic)
- [13] Al-Hassen, Sh.I. (2011) Environmental Pollution in Basra City. PhD Thesis, College of Arts, University of Basra, Basra, 232. (In Arabic)
- [14] Qassim, M.H. (2011) A Geographic Analysis for Air Pollution Problem in Al-Zubayr City and Its Healthy Effects. M.A. Thesis, College of Arts, University of Basra, Basra, 180. (In Arabic)
- [15] Douabul, A.A.Z., Al-Maarofi, S.S., Al-Saad, H.T. and Al-Hassen, Sh.I. (2013) Gaseous Pollutants in Basra City, Iraq. *Air, Soil and Water Research*, **6**, 15-21. <http://dx.doi.org/10.4137/ASWR.S10835>
- [16] Sultan, A.W.A., Al-Hassen, Sh.I., Ateeq, A.A. and Al-Saad, H.T. (2013) Ambient Air Quality in the Industrial Area of Khor Al-Zubayr, Southern Iraq. *Journal of Petroleum Research & Studies*, **4**, 1-11. <http://www.iasj.net/iasj?func=fulltext&aId=87876>
- [17] Karmalla, H.A., Al-Hassen, Sh.I., Adam, R.S. and Qassim, M.H. (2013) A Cartographic Representation to Levels and Impacts of Carbon Monoxide Pollutant in Basra City, Southern Iraq. *Journal of Thi-Qar Science*, **4**, 105-115. (In Arabic) <http://www.iasj.net/iasj?func=fulltext&aId=83346>
- [18] Abdullah, A.S. and Hussien, H.H. (2014) Estimation of Gaseous Pollutants emitted from Al-Najbyia Power Station in Basra. *Journal of Thi-Qar Science*, **4**, 68-71. <http://www.iasj.net/iasj?func=fulltext&aId=92673>
- [19] Al-Hassen, Sh.I., Al-Qarroni, E.H., Qassim, M.H., Al-Saad, H.T. and Alhello, A.A. (2015) An Experimental Study on the Determination of Air Pollutant Concentrations released from Selected Outdoor Gaseous Emission Sources in Basra City (Southern Iraq). *JiARM*, **3**, 88-98. <http://www.jiarm.com/FEB2015/paper19552.pdf>
- [20] Wikipedia-Basra. <https://en.wikipedia.org/wiki/Basra>
- [21] ATSDR (Agency for Toxic Substances and Disease Registry), Toxic Substances Portal-Carbon Monoxide. <http://www.atsdr.cdc.gov/toxfaqs/TF.asp?id=1163&tid=253>
- [22] WHO (World Health Organization) (2010) WHO Guidelines for Indoor Air Quality: Selected Pollutants. WHO Regional Office for Europe, Bonn, 454. http://www.euro.who.int/_data/assets/pdf_file/0009/128169/e94535.pdf
- [23] ATSDR (Agency for Toxic Substances and Disease Registry), Toxic Substances Portal-Nitrogen Oxide. <http://www.atsdr.cdc.gov/toxfaqs/TF.asp?id=396&tid=69>
- [24] ATSDR (Agency for Toxic Substances and Disease Registry), Toxic Substances Portal-Sulfate Oxide. <http://www.atsdr.cdc.gov/toxfaqs/TF.asp?id=252&tid=46>
- [25] ATSDR (Agency for Toxic Substances and Disease Registry), Toxic Substances Portal-Hydrogen Sulfide. <http://www.atsdr.cdc.gov/substances/toxsubstance.asp?toxid=67>
- [26] ATSDR (Agency for Toxic Substances and Disease Registry), Toxic Substances Portal-Methane. <http://www.atsdr.cdc.gov/HAC/landfill/html/ch4.html>
- [27] ATSDR (Agency for Toxic Substances and Disease Registry), Toxic Substances Portal-Formaldehyde. <http://www.atsdr.cdc.gov/mmg/mmg.asp?id=216&tid=39>
- [28] WHO (World Health Organization) (2005) WHO Air Quality Guidelines for Particulate Matter, Ozone, Nitrogen Dioxide and Sulfur Dioxide: Global Update. Geneva, 20. http://whqlibdoc.who.int/hq/2006/who_sde_phe_oh_06.02_eng.pdf