



**ORIGINAL ARTICLE**

## RESPONSE OF FABA BEAN CULTIVARS (*VICIA FABA* L.) TO PHOSPHORUS APPLICATION

Anhar M. Alshummary, Lamiaa M. Alfreeh\* and Kareem H. Mohssen

Department of Field Crops, College of Agriculture, University of Basrah, Basrah, Iraq.

E-mail: lamiaaalfreeh610@gmail.com

**Abstract:** A field experiment was conducted to investigate the impact of phosphorus fertilizer rates ( $P_0=0$ ,  $P_1=35$ ,  $P_2=70$  and  $P_3=105$  Kg/ha) on some growth characteristics, yield component and grain yield of four faba bean cultivars ( $V_1$ =Luzde utono,  $V_2$ =Iquadolis,  $V_3$ =Iquadolge and  $V_4$ =local) during the winter season 2019-2020 at a private field in Al- Zubair district, Basrah, Iraq. A split plot design of three replications was used, the levels of phosphorus were allocated in the main plots while cultivars were in sub-plots. The results showed that application of 70kg/ha resulted in higher number of pods/ plant (10.624), 100 seed weight (126.933g), seed yield about (847.891kg/h) over the unfertilized control treatments. Cultivar Luzde utono resulted in highest growth characteristics, and seed yield (734.307 kg/ha). All interactions were significant. Cultivars Local and Luzde utono with  $P_2=70$  Kg/ha gave the highest seed yield 877.180 kg/ h and 874.370kg/ha, respectively.

**Key words:** Faba Bean, *Vicia faba* L., Phosphorus Application, Split plot design.

### Cite this article

Anhar M. Alshummary, Lamiaa M. Alfreeh and Kareem H. Mohssen (2021). Response of Faba Bean Cultivars (*Vicia faba* L.) to Phosphorus Application. *International Journal of Agricultural and Statistical Sciences*. DocID: <https://connectjournals.com/03899.2021.17.329>

## 1. Introduction

In Iraq, faba bean (*Vicia faba* L.) is one of the main crops of legumes which is grown in many regions, sometimes referred to as broad bean, horse bean, field bean, or tic bean [Singh *et al.* (2010)]. It is used as human food, also for most animal feeds. Its seeds contain a high percentage of protein, which is estimated to be 25-40% [Natalia *et al.* (2008)] in addition to the carbohydrate content of seeds, of which percentage in most cultivars reached 56% [Carmen *et al.* (2005)]. It also improves soil qualities by fixing it to atmospheric nitrogen in the soil if its roots contain root nodes [Mona *et al.* (2011)]. Generally, the lower yield is due to several factors, that contribute to increasing the productivity of this crop, including the good varieties that have the ability to invest the available resources, including the factors related to the soil, such as the abundance of nutrients in it. Phosphorous is one of the critical and determining nutrient. The optimum level of phosphorous in plant tissues leads to an increase in the activity and growth

of the root system and its branching, as well as an increase in the growth of the vegetative parts, and it also helps in increasing the yield, early in the formation and ripening of crops, especially seeds, thus reducing the period of ripening [Havlin *et al.* (1999)]. The lack of available phosphorus in plant is reflected in the yield, as the quantity and quality decrease, the maturity stage is delayed and the chances of plant exposure to diseases increase. Ahmed (2017) found that the application of Phosphorus fertilizer increased seed yields of faba bean. Also, Negasa *et al.* (2019) showed that Phosphorus fertilizers increased growth parameter, yield and yield component. Therefore, the present study was conducted to evaluate the response of bean cultivars to different rates of phosphorous fertilizer in the southern of Iraq.

## 2. Materials and Methods

A field experiment was conducted during the winter season 2019-2020 at a private field in Al- Zubair district, Basrah, Iraq to study the effect of four P fertilizer

(control(P<sub>0</sub>), 35(P<sub>1</sub>), 70(P<sub>2</sub>) and 105 (P<sub>4</sub>) kg P/ha) as calcium super phosphate (21% P) on growth, yield and yield components for four cultivars of faba bean (Luz de otono (V<sub>1</sub>), Icuadolis (V<sub>2</sub>), Icuadolge(V<sub>3</sub>) and local (V<sub>4</sub>)). The soil was sandy loam with a pH value of 7.66 and an organic matter content of 0.11% and available N, P, K of 77.23, 5.37 and 122 mg/Kg, respectively. The experimental design was randomized complete block design (RCBD) using split-plots arrangement with three replications in which P fertilizer treatments were placed in the main plots, while cultivars as sub-plots. Samples soil were analyzed according to the methods outlined by Black (1965) and Page *et al.* (1982). Each experimental plot had an area of 12 m<sup>2</sup> consisting of 5 rows each was 70 cm apart and 3 meter long. Faba bean seeds were planted at 1st of November putting 5 seeds in hills of 20 cm apart. The form of nitrogen fertilization as urea (46% N) applied with a quantity of 60 kg N/ha at three equal doses 15, 30 and 45 days after sowing and the Phosphorus was applied during the oil preparation and before sowing soil before sowing. Twenty days-later after sowing, plants were thinned to one plant/hill. All appropriate management practices were carried out. At harvest, five plants were randomly selected from the center row of each plot to measure as follows: plant height, number of branches/plant, number of pods/plant, 100-seed weight, seeds yield and protein percentage. GenStat program was used to analyse data statistically using L.S.D. test at the probability level of 0.05.

### 3. Results

Data in Table 1 showed that all of the applied treatments of phosphorus fertilizer significantly increased vegetative growth characteristics of faba bean plants. It was observed that plants when treated with P<sub>2</sub> (70 Kg P/ha) gave the highest plant height (91.398 cm), branches/plant (8.707) while the control treatment P<sub>0</sub> gave the lowest 86.317cm, 7.763 branch/plant, respectively. Data in Table 1 clarified that all of phosphor rates significantly increased yield and its components of faba bean plants, where P<sub>2</sub> was the superior for number of pods/Plant (10.624), 100 seed weight(126.933g) seed yield (847.891 Kg/ha) and protein content (20.339%) as compared to the control, which gave the lowest were 8.595 pods/plant, 90.265 g and 445.795 Kg/ha and 16.132%, respectively, whereas the highest number of seeds per pod was obtained by P<sub>3</sub> with 8.215, while the lowest number of seeds with 7.186 seed/pod was found from control (7.186 seed/pod). Several authors have reported significant increase in growth, yield components and yield of faba bean with increase in the rate of application of mineral phosphorus [Mahmoud (2010), Mousa and El-Sayed (2016), El-Sayed *et al.* (2017)]. Moreover, data given in Table 2 indicated that the highest plant height, number of branch/plant and number of pods/plant were obtained by V<sub>1</sub> compared with the other cultivars with 101.275 cm, 9.165 branch/plant and 11.112 pod/plant, respectively, while V<sub>4</sub> gave the lowest values of plant height (81.090 cm) and number of pods/plant (9.375 pod/plant). The highest number of seeds/pods was found in V<sub>2</sub> (8.267

**Table 1:** Effect of different rates of phosphorus fertilizer on growth, yield and yield component of bean.

Phosphorus rates kg/ha	Plant height cm	Number of Branches/plant	Number of Pods/plant	Seeds/pod	seed-100 weight(g)	Seed yield Kg/ha	protein%
P <sub>0</sub>	86.317	7.763	8.595	7.186	90.265	445.795	16.132
P <sub>1</sub>	88.492	8.243	10.461	7.858	120.763	790.339	18.694
P <sub>2</sub>	91.398	8.707	10.624	7.893	126.933	847.891	20.339
P <sub>3</sub>	88.107	8.113	10.343	8.215	117.057	792.691	18.678
RLSD	0.925	0.363	0.114	0.169	1.472	20.211	0.328

**Table 2:** Effect of different rates of phosphorus fertilizer on growth, yield and yield component of bean.

Phosphorus rates kg/ha	Plant height cm	Number of Branches/plant	Number of Pods/plant	Seeds/pod	seed-100 weight(g)	Seed yield Kg/ha	protein%
V <sub>1</sub>	101.275	9.165	11.112	7.233	113.765	734.307	18.174
V <sub>2</sub>	88.258	8.028	9.951	8.267	106.282	713.751	18.605
V <sub>3</sub>	83.692	7.786	9.586	7.807	117.866	725.595	18.664
V <sub>4</sub>	81.090	7.847	9.375	7.845	117.106	703.063	18.400
RLSD	0.925	0.363	0.114	0.169	1.472	20.211	N.S.

**Table 3:** Effect of interaction of phosphor levels and cultivars on growth, yield and yield component of bean.

Treatment	Plant high(cm)	Number of branches/plant	Number of Pods/plant	Number of Seeds/pod	seed weight(g)	Seed yieldKg/ha	protein%	
P <sub>0</sub>	V <sub>1</sub>	100.347	8.687	10.183	7.450	88.253	535.868	16.039
	V <sub>2</sub>	85.363	7.763	8.443	7.353	86.113	427.735	15.811
	V <sub>3</sub>	81.760	7.530	7.520	6.973	93.370	391.594	16.444
	V <sub>4</sub>	77.800	7.073	8.233	6.967	93.323	427.983	16.233
P <sub>1</sub>	V <sub>1</sub>	101.513	9.167	11.343	7.043	116.530	744.748	17.695
	V <sub>2</sub>	88.270	7.977	10.693	8.420	111.357	801.511	18.598
	V <sub>3</sub>	83.547	7.833	10.107	8.133	129.337	850.388	19.447
	V <sub>4</sub>	80.640	7.993	9.700	7.833	125.830	764.708	19.037
P <sub>2</sub>	V <sub>1</sub>	105.960	10.070	11.623	7.140	131.703	874.370	19.587
	V <sub>2</sub>	90.940	8.110	10.453	8.160	114.723	782.800	21.218
	V <sub>3</sub>	85.137	8.120	10.567	7.873	128.810	857.214	20.214
	V <sub>4</sub>	83.557	8.527	9.853	8.400	132.497	877.180	20.336
P <sub>3</sub>	V <sub>1</sub>	97.280	8.737	11.297	7.300	118.573	782.241	19.375
	V <sub>2</sub>	88.460	8.263	10.213	9.133	112.933	842.958	18.792
	V <sub>3</sub>	84.323	7.660	10.150	8.247	119.947	803.183	18.550
	V <sub>4</sub>	82.363	7.793	9.713	8.180	116.773	742.381	17.995
RLSD 0.05	1.600	0.486	0.228	0.337	2.943	40.422	0.657	

seed/pod), while the lowest was found in V<sub>1</sub> (7.233 seed/pod), whereas V<sub>3</sub> was the superior for 100 seed weight (117.866 g). The highest seed yield was recorded from V<sub>1</sub> variety (734.307 Kg/ha) at par with V<sub>3</sub> (725.595 Kg/h), while V<sub>4</sub> gave the lowest value (703.063 t/ha). There were no significant differences among the cultivars in the protein percentage.

Table 3 shows that the effect of the interaction between phosphor rates and cultivars was statistically significant for all traits. The combination of P<sub>2</sub>V<sub>1</sub> was superior with plant high (105.96 cm) number of branch/plant (11.623), while the combination of P<sub>3</sub>V<sub>2</sub> was superior with seeds/pod (9.133), also the interaction between P<sub>2</sub>V<sub>4</sub> gave highest 100 seed weight (132.497 gm) and seed yield (877.180 Kg/ha) with no difference with P<sub>2</sub>V<sub>1</sub> (131.703 Kg/ha). The highest value of protein (21.218 was obtained from P<sub>2</sub>V<sub>2</sub>.

#### 4. Discussion

Phosphorus application significantly influenced some growth parameters of faba bean. Application of 70 kg P/ha gave statistically an increase in plant height, number of branches. This increase may be attributed to the role of phosphorus in the growth of roots and the number of their branches, which led to an increase in the efficiency of the plant in absorbing water and nutrients and its reflection on plant growth [Tisdale *et al.* (1997)]. Also there was an increase in number of pods, 100- seed. This increase is attributed to the positive

effect of Phosphorous which is one of the basic and necessary nutrients for plant growth. It can be attributed to the availability of nutrients and their contribution to increasing vegetative growth. Thus, improving the efficiency of photosynthesis and splitting of carbohydrates into a pod Yield. Also, its direct role to play in most vital biological processes. Thus, it is a major source of cell division and increasing plant growth, and as energy storage and transfer like ADP and ATP, photosynthesis respiration, protein and nucleic acid synthesis, and ion transport across cell membranes [Fageria (2009)]. Increasing the rate of phosphorus from 0 to 70 kg P/ha significantly increased seed yield. It might be as a result of the role of phosphorus in transporting sugars from the places of its formation to where the plants need it, which helps the transfer of Photosynthesis products to the seeds, and thus an increase in the seeds yield. Similar results were reported by Yohannes *et al.* (2014), Negasa *et al.* (2019). Also results showed that cultivars had significant effect on all parameter except protein percentage. The highest seed yield was obtained from Luzde utono cultivar (734.307 Kg/ha), while Local cultivar gave the lowest value (703.063 t/ ha). Thus, this cultivar is recommended for commercial and extensive faba bean farming in the region as a result of its capacity to give high seed yield. The higher seed yield can be attributed to an excess of some of its components

(Number of pods/plant, number of seed / pod and

100-seed weight). These results agreed with those of Ibrahim (2011), Kakahy *et al.* (2012), Al-Freeh *et al.* (2014), Dhary and AL-Baldawi (2017), Almosawy (2018) and Al-Shummary (2020)

Phosphor at the level of 70 Kg/ ha along with Luz de otono and Local cultivars was found to be the optimum level for yield and its component of faba bean. Weldua *et al.* (2012) confirmed the positive effect of application of phosphor levels on faba bean growth, and the variation of cultivars response to applied P fertilization levels.

## 5. Conclusion

The addition of phosphate fertilizer to the faba bean cultivars showed a significant increase in the production of seeds. Adding 70 kgP/ha was proved the productivity of faba bean indicating the importance of Phosphorus macronutrient in achieving high seed yield from faba bean and be recommended for faba bean production in the study area. Local cultivar was the best in growth, yield components and seeds yield compared to the other cultivars. Therefore, the production and productivity of faba beans could be enhanced by using cultivars with better yields as Luzde utono. However, it could be concluded that under the conditions of the experiment, planting Luz de otono and Local cultivars under 70 Kg P/ ha is recommended.

## Acknowledgement

First author would like to express his thanks to Dr. Sundus A.M. Al-Abdulla, Department of Field Crops, Basrah Univ. for her valuable opinions at manuscript reading.

## References

- Ahmed, I. (2017). The Effect of Phosphorus Fertilizers on Faba bean plant growth in soils were polluted by heavy metals. *Eurasian Journal of Science & Engineering*, **3(2)**, 175-180.
- Al-Freeh, L.M., R.H. Al-Obeid and Z.A. Abdul Razak (2014). Response to two types of beans *Vicia faba* to different levels of Potassium in two locations of the province of Basra, *Basrah J. Agric. Sci.*, **27(2)**, 314-329.
- Almosawy, A.N., A.A. Alamery, F.S. Al-Kinany, H.M. Mohammed, L.Q. Alkinani and N.N. Jawad (2018). Effect of optimum nanoparticles on growth and yield of some broad bean cultivars (*Vicia faba*). *International Journal of Agricultural and Statistical Sciences*, **14(2)**, 525-528.
- Al-Shummary, A.M.J. (2020). The role of foliar zinc application on growth and yield of faba bean varieties. *International Journal of Agricultural and Statistical Sciences*, **16(Supplement 1)**, 1157-1161. DocID:https://connectjournals.com/03899.2020.16.1157
- Black, C.A. (1965). *Methods of Soil analysis*, Part I. Soc. Agron. Inc. Publ. Madison Wisconsin, U.S.A.
- Carmen, M.A., Z.J. Carmen, S. Salvador, N. Diego, R.M. Maria Teresa and T. Maria (2005). Detection for agronomic traits in faba bean (*Vicia faba* L.). *Agric. Conspec. Sci.*, **70(3)**, 17-20.
- Dhary, S.I. and M.H.K. AL-Baldawi (2017). Response of different varieties of faba bean to plant source organic fertilizers. *The Iraqi Journal of Agricultural Sciences*, **48(4)**, 1141-1147.
- El-Sayed E.A. El-Sobky and M.A.T. Yasin (2017). Phosphorus and micronutrients fertilization impact on faba bean (*Vicia faba* L.) productivity. *Zagazig J. Agric. Res.*, **44(3)**, 853-863.
- Fageria, N.K. (2009). *The Use of Nutrients in Crop Plants*. 1st Ed. CRC press. New York.
- Havlin, J.L., J.D. Beaton, S.L. Tisdale and W.L. Nelson (1999). *Soil Fertility and Fertilizers*: 6th Edition. Prentice Hall. Upper Saddle River, NJ.
- Ibrahim, R.H. (2011). Response of two faba bean cultivars to foliar application. *The Kufa Journal of Agriculture Sciences*, **3(2)**, 85-92.
- Kakahy, A.N., N.D. Ahmad and A.S. Abdullahi (2012). The effect of planting distance on yield of bean (*Vicia faba* L.) under drip irrigation system. *African Journal of Agricultural Research*, **46(7)**, 6110-6114.
- Mahmoud, A.N. (2010). Economic analysis of the response of broad beans to levels of n and p fertilizers. *Agric. Sci.*, **41(5)**, 125-132.
- Mona, A.M., M.A. Sabah and A.M. Rehab (2011). Influence of potassium sulfate on faba bean yield and quality. *Australian Journal of Basic and Applied Sciences*, **5(3)**, 87-95.
- Mousa, A.M. and S.A. El-Sayed (2016). Effect of intercropping and phosphorus fertilizer treatments on incidence of rhizoctonia root rot disease of faba bean. *Int. J. Current Microbiol. and Appl. Sci.*, **5(4)**, 850-863.
- Natalia Gutierrez, C.M. Avila., M.T. Moreno and A.M. Torres (2008). Development of SCAR markers linked to zt-2, one of the genes controlling absence of tannins in faba bean. *Aust. J. of Agric. Res.*, **59**, 62-68.
- Negasa, G, B. Bedadi and T. Abera (2019). Influence of Phosphorus Fertilizer Rates on Yield and Yield Components of Faba Bean (*Vicia faba* L.) Varieties in Lemu Bilbilo District of Arsi Zone, Southeastern Ethiopia. *International Journal of Plant & Soil Science*, **28(3)**, 1-11,
- Page, A.L., R.H. Miller and D.R. Keeney (1982). *Methods of Soil Analysis*, Part 2, 2nd Edn. ASA Inc. Madison,

- Wisconsin, U.S.A.
- Singh, A.K., N. Chandra, R.C. Bharati and S.K. Dimree (2010) Effect of seed size and seeding depth on Fava bean (*Vicia faba* L.) productivity. *Environmental Ecology*, **28(3A)**, 1722-1527.
- Tisdale, S.L., W.L. Nelson, J.D. Beaton and J.L. Harlin (1997). *Soil fertility and fertilizer*. Prentice. Hall of India, New Delhi.
- Yohannes, D., H. Kiros and W. Yirga (2014) Inoculation, Phosphorus and Zinc fertilization effects on nodulation, Yield and nutrient uptake of faba bean (*Vicia faba* L.) grown on Calcaric Cambisol of Semiarid Ethiopia. *Journal of Soil Science and Environmental Management*, **6(1)**, 9-15.
- Weldua, Y., M. Haileb and K. Habtegebrielb (2012). Effect of zinc and phosphorus fertilizers application on yield and yield components of faba bean (*Vicia faba* L.) grown in calcaric cambisol of semi-arid northern. *Ethiopia. J. Soil Sci. Environ. Manag.*, **3(12)**, 320-326.