

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/325514450>

# Original Research Article Virulence factors of Methicillin Resistant Staphylococcus aureus (MRSA) isolated from burn patients

Article · January 2015

CITATIONS

7

READS

242

3 authors:



Zainab degaim

University of Thi-Qar

10 PUBLICATIONS 11 CITATIONS

SEE PROFILE



Wafaa Sadoon

University of Basrah

14 PUBLICATIONS 76 CITATIONS

SEE PROFILE



Saad S. Hamim

University of Thi-Qar

22 PUBLICATIONS 20 CITATIONS

SEE PROFILE


Some of the authors of this publication are also working on these related projects:



Bacteriology [View project](#)



IMAN ALFAYYADH [View project](#)



## Original Research Article

# Virulence factors of Methicillin Resistant *Staphylococcus aureus* (MRSA) isolated from burn patients

Zainab Dakhil Degaim<sup>1\*</sup>, Wafaa Sadoon Shani<sup>2</sup> and Saad Salman Hamim<sup>3</sup>

<sup>1</sup>Microbiology Department-College of Medicine -University of Thi-Qar, Iraq

<sup>2</sup>Biology Department-College of Science-University of Basrah, Iraq

<sup>3</sup>Pathological analysis department-College of Science-University of Thi-Qar, Iraq

\*Corresponding author

## ABSTRACT

### Keywords

MRSA,  
Burn,  
Virulence  
factors,  
Molecular  
detection

The present study aimed to investigate some of virulence factors among Methicillin Resistant *Staphylococcus aureus* (MRSA) isolated from burn wound. From a total of 126 isolates of *S. aureus*, only eighty five(67.46%) isolates of MRSA were obtained from burn patients at Al- Hussain teaching hospital during the period from July to November, 2014 in Thi-Qar province, Iraq. All MRSA isolates were examined using Polymerase Chain Reaction (PCR) for detection 16SrRNA, *mec A* gene and some virulence factors of this bacteria include *sea*, *hla*, *hly*, and *cap 8*. The results revealed that all isolates have 16SrRNA and *mec A* genes that were used to confirm these bacteria as Staphylococci and MRSA respectively. The virulence factors detection results showed percentages of (72.941%), (82.352%) and (85.882 %) of isolates have *sea*, *hla* and *hly* genes respectively, while only 69 (81.176%) of isolates have *cap8* gene.

## Introduction

Thermal injury destroys the skin barriers that normally prevent invasion by microorganisms (Singh *et al.*, 2003). Burn patients become susceptible to various infections due to the loss of this protective barrier and decreased cellular and humoral immunity (Wong *et al.*, 2002). In these patients, burn wound infections can easily escalate into sepsis (Church *et al.*, 2006).

The common pathogens isolated from burn wound are *S.aureus*(75%), *Pseudomonas aeruginosa*(25%), *Streptococcus pyogenes* (20%) and various coliform bacilli (5%) (Ahmad and Iranzo, 2003).

MRSA is the most important pathogen among Staphylococci (Lee *et al.*, 2007). MRSA strains are isolated in more than half of all community and hospital infections (Klevens *et al.*, 2007). MRSA has become a major public health problem worldwide, and the problem of MRSA continues to rise (Nimmo *et al.*, 2006 and Jarvis *et al.*, 2007). MRSA has been the most commonly recognized multidrug-resistant pathogen in the universe and the emergence of MRSA strains found in increasing number of infections and often multi drug resistant in nature now pose serious therapeutic problems to clinicians (Groundmann *et*

*al.*,2006; Marais *et al.*,2009).

Most MRSA strains carry *mecA* encoding low affinity penicillin-binding protein PBP2a (or PBP2') (Hiramatsu *et al.*, 2001). The MRSA characteristic phenotype is due to the presence of *mecA* which encodes a PBP2a, with degraded affinity for  $\beta$ -lactams (Oliveira and De Lencastre, 2011; Moellering, 2012).

The virulence factors of *Staphylococcus* include surface components, such as the capsule, peptidoglycans, teichoic acid, protein A, enzymes such as (esterases, lipases, fatty-acid modifying enzymes, various proteases, hydrolytic enzymes, catalase, betalactamase), and various toxins, such as (leukocidins, enterotoxins, TSST-1 and alpha, beta, gamma and delta hemolysins) (Vasconcelos and Cunha, 2010).

One of the virulence factors of *S. aureus* is cytolytic, pore-forming toxin (Diep and Otto, 2008), such as alpha-hemolysin (*Hla*) has been implicated in the pathogenesis of *S. aureus* (Labandeira-Rey *et al.*, 2007). *Hla* has cytolytic activity toward a variety of host cell types, including human keratinocytes, epithelial cells and lymphocytes (Hocke *et al.*, 2006; Wardenburget *al.*,2008). Other types of hemolysin is Beta-hemolysin (*Hlb*) is a magnesium-dependent sphingomyelinase C that induces lysis of sheep erythrocytes and human monocytes (Walev *et al.*, 1996).

*S. aureus* is produced one of the extracellular protein toxins, staphylococcal heat stable enterotoxin (SE) is the most important virulent factors belonging to the superantigen family (Pinchuk *et al.*, 2010), and many strain of *S. aureus*, especially MRSA, secreted one or more specific staphylococcal exotoxins, including

staphylococcal enterotoxins (SEs), (Llewelyn and Cohen, 2002).

Among those factors considered for typing, capsular polysaccharides expressed by *S. aureus* are one of them, since they are also important in the pathogenesis of staphylococcal infections, most *S. aureus* isolates are encapsulated and so far eleven capsular serotypes have been described, of these, types *cap 5* and *cap 8* predominate in approximately 75% of the clinical isolates (Murphy *et al.*, 2011).

## Materials and Methods

### Samples collection

Two hundred and seventy six samples were collected from burn patients in burn unit of AL-Hussain Teaching Hospital of Thi-Qar province in the period from July to November, 2014 by moistened sterile swabs with normal saline, then these swabs directly inoculated on Mannitol salt agar (LAB/ United Kingdom) and incubated at 37°C for 24 hours.

### Identification of *S. aureus*

*S. aureus* was identified depending on the morphological properties on culture media and biochemical tests which done according to Bergeys manual (MacFaddin, 2000). API Staph system was used as identification system for *Staphylococcus* and *Micrococcus*. This test was done according to the company instructions (BioMerieux, France).

### StaphyloMonotec test kit Plus

This kit is a new rapid agglutination test for differentiation between *S. aureus* and other *Staphylococcus*. The test was performed according to the directions of manufacturing

company (Fluka Analytical, Switzerland).

### **Antibiotic sensitivity test**

The antibiotic sensitivity test was done by the agar disc diffusion method as described by (Kirby and Bauer, 1966).

### **Molecular Detection**

#### **DNA extraction**

DNA from all MRSA isolates were extracted using Genomic DNA Extractions spin kit (Bosphore, Anatolia genewors). 16S rRNA, *mecA*, *sea*, *hla*, *hly*, and *cap8* genes were identified by using primers described in Table (1).

Amplification of the *mecA* gene was done using primer described by (Jonas *et al.*, 2002). The final volume of reaction tubes is 20µl, consist of 10 µl Master Mix., 1.25µl of both Forward (F) and Reverse (R) of the primer specific for the *mecA* gene, 5µl of template DNA and complete the volume by adding free water to 20µl. Amplification of the 16SrRNA, *sea*, *hla*, *hly*, *hla*, and *cap8* genes were done using primers described as above (Table 1). The final volume of reaction tubes is 20µl, consist of 10 µl Master Mix., 1µl of both F. and R. of the primers specific for these genes, 5µl of template DNA and complete the volume by adding deionizing water to 20µl.

### **Results and Discussion**

Eighty five isolates (67.460 %) of MRSA were obtained from *S. aureus* isolates that collected from burn patients. All strains identified was done by cultural, biochemical and serological tests to confirm exact identification as *S. aureus*. MRSA outbreaks are estimated in about 40-60% of *S. aureus* outbreaks, which are mainly affected by the

infection control program and medical treatments leading to a wide range of hospital infections (Fatholahzadeh *et al.*, 2008). Alfatemi *et al.*, (2014) showed the prevalence of MRSA among *S. aureus* isolates was (42.3%), which indicates little difference in terms of frequency with studies by Fatholahzadeh *et al.*, (2008) who reported MRSA prevalence of (36%) in Tehran.

The prevalence of MRSA in present study was slightly, in agreement with other studies in Iraq, which recorded percentages of (65.3%), (88%) and (75%) respectively (Al-Mussawi, 2014; Yaseen *et al.*, 2013 and Al-Azawi, 2013).

The molecular diagnostic of this bacteria was performing to all MRSA isolates through the amplification of 16SrRNA and *mec A* genes to confirm that the tested isolates are staphylococci and MRSA respectively. Other genes were used to detect many virulence factors of MRSA that included *hla*, *hly*, *sea* and *cap8* genes.

All isolates were showed positive results of both 16SrRNA and *mec A* genes (100%) (Table 2). The bands were 756, 310 bp size corresponds to amplification of 16SrRNA and *mecA* genes respectively, Fig(1 and 2). Al-Talib *et al.*, (2009) reported that all isolates had 16SrRNA, 82 contained *mecA* genes. Current study results agreed with Makgotlho, (2009) who showed that all isolates 97/97 (100%) have 16SrRNA gene while *mec A* gene was detected in 96/97 (99%) of the MRSA isolates, which did not show the presence of *mec A* gene was, however phenotypically identified as MRSA.

The results showed that 69 (81.1 %) of isolates have *cap8* gene (Table 2), the bands were (450 bp) size corresponds to amplification of *cap8* gene, Fig (6). Udo and

Sarkhoo (2010) reported that capsular polysaccharides and types prevalence was (77.3%) and only three isolates (2.2%) yielded negative result for both *cap5* and *cap8*.

Seventy (82.35%) of isolates have *hla* gene only, Table (2). The bands were (209 bp) size corresponds to amplification of *hla* gene, Fig (4). Most of *S. aureus* isolated from human have usually an alpha haemolytic character, because the human platelets and monocytes are more sensitive to the alpha toxin (Todar, 2005). Kateete *et al.*(2011) showed the frequency of *hla* gene was 100%. Likewise, in a study from the United States the *hla* gene frequency was reported at 100% (Shukla *et al.*,2010). The percentage of *hla* gene in MRSA isolates was 73 (85.88%), Table (2). The bands were

(833 bp) size corresponds to amplification of *hla* gene, Fig (5). The study performed by Rusenova *et al.*,(2013) showed that 31 MRSA isolates (42.5%) for beta toxin, 41 (56.2%) of isolates showed double hemolysis (alpha + beta hemolysins), and 1 (1.4%) was non-hemolytic. MRSA isolates have 62 (72.94%) of *sea* gene Table (2). The bands were (120 bp) size corresponds to amplification of *sea* gene, Fig (3). Alfatemi *et al.*,(2014) the frequency of the *sea* gene was 27.39%. Our study was in agreement with Udo and Sarkhoo, (2010) whom reported that 103 (76.3%) isolates yielded positive results for *sea*. However, the role of *S. aureus* superantigenic toxins in the severity of septicemia patients should not be discounted as *sea* is significantly associated with severity of sepsis caused by *S. aureus* (Ferry *et al*, 2005).


**Table.1** Oligonucleotide primers sequences for PCR amplified of 16SrRNA, *mecA*, *sea*, *hla*, *hla* and *cap 8* genes

Genename	Primer Sequences (5'-3')	Length	References
16SrRNA	F: AAC TCT GTT ATT AGG GAA GAA CA R: CCA CCT TCC TCC GGT TTG TCA CC	756 bp	(McClure <i>et al.</i> , 2006)
<i>mecA</i>	F: GTA GAA ATG ACT GAA CGT CCG ATA A R: CCA ATT CCA CAT TGT TTC GGT CTA A	310 bp	(Gehaet <i>et al.</i> , 1994)
<i>hla</i>	F: GCC AAA GCC GAA TCT AAG R: GCG ATA TAC ATC CCA TGG C	833 bp	(Booth <i>et al.</i> , 2001)
<i>hla</i>	F: CTG ATT ACT ATC CAA GAA ATT CGA TTG R: CTT TCC AGC CTA CTT TTT TAT CAG T	209 bp	(Mehrotraet <i>et al.</i> ,2000)
<i>cap8</i>	F: GCG CTA CAA ACA TTA AGC AT R: TTC TTA GCC TGC TGG CAT C	450 bp	(Sauet <i>et al.</i> ,1997)
<i>sea</i>	F: TTGAAACGGTTAAAACGAA R: GAACCTTCCCATCAAAAACA	120 bp	(Betley and Mekalanos, 1988) with modified


**Table.2** The percentage of genes in MRSA isolates

Genes	Positive %	Negative %
<b>16SrRNA</b>	85 (100 %)	-
<i>mecA</i>	85 (100 %)	-
<i>hla</i>	70 (82.352 %)	15 (17.647 %)
<i>hlb</i>	73 (85.882 %)	12 (14.117 %)
<i>sea</i>	62 (72.941 %)	23 (27.058 %)
<i>cap 8</i>	69 (81.176 %)	16 (18.823 %)


**Figure.1** Agarose gel electrophoresis of 16S rRNA gene amplification




**Figure.2** agarose gel electrophoresis of *mec A* gene amplification



**Figure.3** Agarose gel electrophoresis of *sea* gene amplification



**Figure.4** Agarose gel electrophoresis of *hla* gene amplification



**Figure.5** Agarose gel electrophoresis of *hly* hemolysin gene amplification





Figure.6 Agarose gel electrophoresis of cap8 gene amplification



## References

- Ahmad, S.I and Iranzo, O. G. (2003). Treatment of post burns bacterial infections by Fenton reagent, particularly the ubiquitous multiple drug resistant *Pseudomonas* species. *Medical Hypotheses*, 61(4):431–434.
- Al-Azawi, I.H.S. (2013). Antibiotic susceptibility pattern and *mec A* gene detection in methicillin resistant *Staphylococcus aureus* (MRSA) isolated from burn and wound in Al-Diwaniya city. *Journal of Babylon university, Pure and applied sciences*; 21(3): 917-926.
- Al-Mussawi, A.A. (2014). Detection of *Staphylococcus aureus* and Methicillin Resistant *Staphylococcus aureus* (MRSA) from human clinical specimens using conventional biochemical tests and chromogenic media. *Indian Journal of applied research* ;4 (2):7-9.
- Al-Talib, A.; Yean, C.Y.; Al-Khateeb, A.; Hassan, H.; Singh, K.B.; Al-Jashamy, K. and Ravichandran, M. (2009). A pentaplex PCR assay for the rapid detection of Methicillin-Resistant *Staphylococcus aureus* and Pantone-Valentine Leucocidin. *BMC Microbiology*, 9:113 doi:10.1186/1471-2180-9-113.
- Betley, M.J. and Mekalanos, J.J. (1988). Nucleotide sequence of the type A staphylococcal enterotoxin gene. *Journal of Bacteriology* 170, 34–41.
- Booth, M.C.; Pence, L.M. ; Mahasresthi, P.; Callegan, M.C. and Gilmore, M.S. (2001). Clonal association among *Staphylococcus aureus* isolates from various sites of infection. *Inf. and Immun*, 69: 345-352.
- Church, D.; Elsayed, S.; Reid, O.; Winston, B. and Lindsay, R. (2006). Burn wound infections. *Clin Microbiol Rev*, 19: 403–434.
- Diep, B.A. and Otto, M. (2008). The role of virulence determinants in community associated MRSA pathogenesis. *Trends Microbiol*, 16: 361–369.
- Fatholahzadeh, B.; Emaneini, M.; Gilbert, G.; Udo, E.; Aligholi, M.; Modarressi, M.H. et al. (2008). Staphylococcal cassette chromosome *mec* (SCC*mec*) analysis and antimicrobial susceptibility patterns of Methicillin Resistant *Staphylococcus aureus* (MRSA) isolates in Tehran, Iran. *Microb Drug Resist.*;14(3):217–20.
- Ferry, T.; Thomas, D.; Genestier, A.L., et al. (2005). Comparative prevalence of superantigen genes in *Staphylococcus aureus* isolates causing sepsis with and without septic shock. *Clin Infect Dis*, 41: 771-7.
- Geha, D. J.; Uhl, J. R.; Gustafarro, C. A. and Persing, D. H. (1994). Multiplex PCR



- for identification of Methicillin Resistant staphylococci in the clinical laboratory. *J. Clin. Microbiol.* 32:1768–1772.
- Groundmann, H.; Sousa, M.; Boyce, J. and Tiemesma, E. (2006). Emergence and resurgences of MRSA as a public health threat. *Lancet.*; 368(9538):874-85.
- Hiramatsu, K.; Cui, L.; Kuroda, M. and Ito, T. (2001). The emergence and evolution of Methicillin Resistant *Staphylococcus aureus*. *Trends Microbiol*; 9: 486-93.
- Hocke, AC.; Temmesfeld-Wollbrueck, B.; Schmeck, B. *et al.* (2006). Perturbation of endothelial junction proteins by *Staphylococcus aureus* alpha-toxin: inhibition of endothelial gap formation by adrenomedullin. *J Histochem Cell Biol*; 126:305–316.
- Alfatemi, H.; S.M.H.; Motamedifar, M.; Hadi, N. and Saraie, H.S.E. (2014). Analysis of Virulence Genes Among Methicillin Resistant *Staphylococcus aureus* (MRSA) Strains. *Jundishapur J Microbiol.* 7(6): e10741.
- Jarvis, WR.; Schlosser, J.; Chinn, RY.; Tweeten, S. and Jackson, M. (2007). National prevalence of methicillin-resistant *Staphylococcus aureus* in inpatients at US health care facilities, *Am. J. Infect. Control.*, 35: 631-637.
- Jonas, D.; Speck, M.; Daschner, F. D. and Grundmann, H. (2002). Rapid PCR-Based Identification of Methicillin-Resistant *Staphylococcus aureus* from Screening Swabs. *J. Clin. Microbiol.*, 40(5):1821- 1823.
- Kateete, DP.; Namazzi, S.; Okee, M.; Okeng, A.; Baluku, H.; Musisi, NL. *et al.* (2011). High prevalence of methicillin resistant *Staphylococcus aureus* in the surgical units of Mulago hospital in Kampala, Uganda. *BMC Res Notes*; 4:326.
- Kirby, W.M.; Baur, A.W.; Scherris, J.C.; and Torch, M. (1966). Antibiotic susceptibility testing by standardized single methods. *AM. J.eli. Path* 45:493-496.
- Klevens, RM.; Morrison, MA.; Nadle, J. *et al.* (2007). Invasive Methicillin Resistant *Staphylococcus aureus* infections in the United States. *JAMA*, 298: 1763–1771.
- Labandeira-Rey, M.; Couzon, F.; Boisset, S.; Brown, EL.; Bes, M. *et al.* (2007). *Staphylococcus aureus* Pantone-Valentine leukocidin causes necrotizing pneumonia. *J Science* 315: 1130–1133.
- Lee, SS.; Kim, HS.; Kang, HJ.; Kim, JK. and Chung, DR. (2007). Rapid spread of Methicillin Resistant *Staphylococcus aureus* in a new hospital in the broad-spectrum antibiotic era. *J Infect*, 55: 358-62.
- Llewelyn, M. and Cohen, J. (2002). Superantigens: microbial agents that corrupt immunity. *Lancet Infect Dis* 2, 156–162.
- MacFaddin, J.F. (2000). Biochemical tests for identification of medical bacteria, 3rd ed.; Lippincott Williams & Wilkins: USA.
- Makgotlho, P. E. (2009). Molecular Characterization of Methicillin Resistant *Staphylococcus aureus* strains, MSC. University of Pretoria, South Africa.
- Marais, E.; Aithma, N.; Perovic, O.; Oosthuysen, WF.; Musenge, E.; Dusé, AG. (2009). Antimicrobial susceptibility of Methicillin Resistant *Staphylococcus aureus* isolates from South Africa. *J S Afr Med*, 99(3):170-3.
- McClure, J. *et al.*, (2006). Novel multiplex PCR assay for detection of the Staphylococcal virulence marker pantone valentine leukocidin genes and simultaneous discrimination of methicillin- susceptible from resistant staphylococci. *J of clinic Micro*, 44:1141-4.
- Mehrotra M.; Wang, G. and Johnson, WM. (2000). Multiplex PCR for detection of genes for *Staphylococcus aureus* enterotoxins, exfoliative toxins, toxic shock syndrome toxin 1, and methicillin resistance. *J Clin Microbiol.*; 38(3):1032–5.
- Moellering, Jr. (2012). RC. MRSA: the first

- half century. *J Antimicrob.Chemother*, 67:4-11.
- Murphy, E.; Lin, SL.; Nunez,L.; Andrew, L.; Fink, PS.; Dilts, DA.*et al.*,(2011). Challenges for the evaluation of *Staphylococcus aureus* protein based vaccines: monitoring antigenic diversity. *J of Hum Vaccin*; 7:51-9.
- Nimmo, GR.; Coombs, GW.; Pearson, JC.; O'Brien, FG. and Christiansen, KJ. (2006). Methicillin Resistant *Staphylococcus aureus* in the Australian community: an evolving epidemic. *Med. J. Aust.* 184: 384-388. 17.
- Oliveira, DC. and De Lencastre, H. (2011). Methicillin-Resistance in *Staphylococcus aureus* is not affected by the Overexpression in Trans of the *mecA* Gene Repressor: A- Surprising Observation. *J. PLOS ONE*; 6: 1-9.
- Pinchuk IV, Beswick EJ, Reyes VE. (2010). Staphylococcal enterotoxins. *Toxins*, 2:2177-97.
- Rusenova, N.; Gebreyes, W.; Koleva, M.; Mitev, J.; Penev, T. Vasilev, N. and Miteva, T. ( 2013). Comparison of Three Methods for Routine Detection of *Staphylococcus aureus* Isolated from Bovine Mastitis. *J of KafkasUniv Vet FakDerg*, 19 (4): 709-712.
- Sau, S.;Bhasin, N. ; Wann,E.R. ; C. Lee, J.C. Foster, T.J. and Lee, C.W. (1997). The *Staphylococcus aureus* allelic genetic loci for serotype 5 and 8 capsule expression contain the type-specific genes flanked by common genes. *Microbiology*, 143,2395-2405.
- Shukla, SK.; Karow, ME.; Brady, JM.; Stemper, ME.;Kislow, J.; Moore, N.*et al.* (2010). Virulence genes and genotypic associations in nasal carriage, community-associated methicillin-susceptible and methicillin-resistant USA400 *Staphylococcus aureus* isolates. *J Clin Microbiol.*;48(10):3582–92.
- Singh, N.P.;Goyal, R.;Manchanda, V.; Das, S.;Kaur, I. and Talwar, V. (2003). Changing trends in bacteriology of burns in the burns unit, Delhi, India. *Burns*, 29(2): 129-132.
- Todar, K. (2005). Todar Online Textbook of Bacteriology. *Staphylococcus*. University of Wincosin-Madison Department of Bacteriology. [www.textbookofbacteriology.net/staph.html](http://www.textbookofbacteriology.net/staph.html).
- Udo, E.E. and Sarkhoo,E. (2010). The dissemination of ST80-SCCmec-IV community-associated methicillin resistant *Staphylococcus aureus* clone in Kuwait hospitals. Udo and Sarkhoo *Annals of Clinical Microbiology and Antimicrobials*, 9:31.
- Vasconcelos, NG. and Cunha, MLRS. (2010). Staphylococcal enterotoxins: Molecular aspects and detection methods. *J Public Health Epidemiol*; 2: 29-42.
- Walev, I.; Weller, U.;Strauch, S.; Foster, T. and Bhakdi, S. (1996). Selective killing of human monocytes and cytokine release provoked by sphingomyelinase (beta-toxin) of *Staphylococcus aureus*. *Infect Immun* 64: 2974-2979.
- Wardenburg, B. J. and Schneewind, O.(2008). Vaccine protection against *Staphylococcus aureus* pneumonia. *J Exp Med*; 205:287–294.
- Wong, T.H.; Tan, B.J.; Ling, M.L. and Song, C. (2002). Multi-resistant *Acinetobacter baumannii* on a burns unit – clinical risk factors and prognosis. *Burns*, 28(4): 349-357.
- Yaseen, I.H. *et al.*, (2013). high prevalence of multidrug resistant MRSA and VRSA of different infection from Al-Jumhuory Teaching Hospital patients in Mosul. *J of Life Sciences*, 7(12)1255-9.