



## Research Article

The work is licensed under



## Physico Chemical Characterization for Water of Shatt Al- Basrah Channel, 2009-2010

Al-Imarah, F. J. M.<sup>1\*</sup>, Al-Jorany, Y. SH. J.<sup>2</sup>, Mizhir, A. A.<sup>3</sup>

<sup>1</sup>Department of Chemistry and Marine Environ Pollution, Marine Science Center, Basrah, Iraq

<sup>2</sup>Dept of Natural Marine Sciences, Faculty of Marine Sciences, Basrah, Iraq

<sup>3</sup>Department of Chemistry, College of Science, Basrah University Basrah, Iraq

\*Corresponding Author: Al-Imarah, F. J. M., Department of Chemistry and Marine Environ Pollution, Marine Science Center, Basrah, Iraq

Received: 18 May 2017

Revised: 22 August 2017

Accepted: 28 August 2017

### ABSTRACT

During the period 2009-2010, the concentrations of certain physical pH, EC, and Turbidity; chemical: Salinity, TDS, Chloride, Calcium, Magnesium, Bicarbonate, Sulphate, Sodium, Potassium as well as Nutrients (Phosphate, Nitrite, Nitrate, and Silicate) and trace elements (Fe, Cu, Pb, Co, and Ni) were estimated in water samples from three stations along Shatt Al-Basrah Channel, 1) at the beginning of the channel which is connected to Garmat Ali River, 2) The middle of the Channel representing the discharging point of waste water from The city of Basrah, and 3) the end of the channel at which Khawr Al-Zubair lagoon begins. Recorded levels for most of the chemical parameters were alternative throughout the study seasons and assigned mainly due to the pollution by either domestic, Agricultural or domestic or industrial pollutions. For heavy metals, higher levels of Fe, Co, Cd, and Pb were recorded at station 2 as a result of waste water pollution, while Ni was recorded at station 3 due to the effect of petroleum pollution from nearby refinery.

**Keyword:** Physical; chemical; Basrah; water; parameter

### INTRODUCTION

The quality of water is defined in terms of physical, chemical, and biological characterizations and ascertaining its quality is important before use for various intended purposes such as potable, agricultural, and industrial water usages [1]. Shatt Al-Basrah canal is an extended drainage canal for the Main Outlet Drainage (MOD) to draw saline return water from irrigated agricultural lands which turned later to

marshland reclamation. This was to be achieved by diverting flow from Al-Hammar marsh and flushing it out directly to the Arabian Gulf via Khawr Al-Zubair and Khawr Abdullah [2]. Nitrate and phosphate are essential nutrients for plant growth. These nutrients are derived from variety of sources including decomposition of plant and animal materials, river runoff from agriculture field sand industrial wastes. There concentrations in water can vary significantly with time [3].


Naturally, N and P concentrations in rivers vary with basin geology, vegetation and climate [4]. Natural  $\text{NO}_3^-$  and  $\text{PO}_4^{3-}$  concentrations in rivers range 0.05-0.2 and 0.002-0.025 mg/l respectively [5]. Alexander and Smith [6] (2006) reported that during the last 30 years, total nitrogen and total phosphorous concentrations in the U.S. were

## METHODOLOGY

Three stations were selected along Shatt Al-Basrah channel, Station 1 ( $S_1$ ) represents the entrance to the channel, station 2 ( $S_2$ ) represents the middle part of the channel after the joined point of Shatt Al-Basrah channel with the Main Outlet Drainage (MOD), and station 3 ( $S_3$ ) is nearly at the exit point of Shatt Al-Basrah channel to Khawr Al-Zubair, as shown in figure 1. These locations represent different ecological and environmental changes. Based on seasonal water quality measurements, water samples were collected seasonally from the selected sites, and adopting the following methodology, pH, conductivity, salinity, and TDS were measured in the field by using calibrated portable instruments.

decreased slightly following a reduction in agricultural intensity. On the other hand, the concentrations of nutrients experienced tremendous increase during 50 years in the waters of Yangtze river in China [7] due to the intensification of Agriculture following reformation [8].

Chemical parameters were measured in the lab by according to standard methods [9], Sodium and Potassium by flame photometer, Sulphate by turbidimetry, bicarbonate by titration. For the determination of nitrite and nitrate, water sample is mixed with ammonium chloride at  $\text{pH} = 8.5$ , nitrate is reduced to nitrite after passing the sample across a column of Cadmium metal, the resulting sample is treated with a solution of sulphanic amide then with 1-naphthyl ethylene diamine to produce a solution with pink color, its intensity is related directly to the concentration of both nitrite and nitrate in the sample, the absorbance is measured at 543 nm. For this measurement, sodium nitrite and sodium nitrate are used as standard solutions.



**Fig. 1. Location map of southern Iraq showing the sampling stations 1-3 along Shatt Al-Basrah**

## Channel

Phosphate was measured according to Murphy and Riley [10] (1962) in which the blue colour of phospho – molybdenum complex is formed after the reduction of the yellow complex. The intensity of the complex is related directly to the concentration of phosphate, its absorbance is measured at 880 nm. Silicate in the studied stations was measured by mixed reagent of

ammonium molybdate and oxalic acid and the intensity of the complex formed was measured at 810 nm.

For trace elements, water samples were collected in 1 liter polyethylene bottles, few drops of nitric acid is added to each sample, cooled, and transferred to the lab in cooling box. In the lab, each sample was digested in Teflon beaker until

dryness with acid mixture of nitric and hydrochloric acids, then each sample was dissolved in 25 ml distilled water, and samples were kept in store prior to analysis. The concentrations of trace elements Fe, Cu, Pb, Co, and Ni were determined by atomic absorption spectrometer model PyeUnicum SP9.


## RESULTS AND DISCUSSION

Within monitoring programs, generally relevant physical, chemical, and biological parameters are monthly, seasonally or annually sampled and analyzed for the water quality variations. This monitoring may allow proper identification of contaminants and their sources [11].


In this study, station 1 is affected heavily by fresh water coming from Al-Hammar marsh, station 3 is heavily affected by polluted water from certain

industrial estate such as Al-Shuaibah petroleum refinery, petrochemical, iron casting and fertilizer of Khawr Al-Zubair, as well as saline water from the Arabian Gulf during high tide.


The seasonally measured values of selected parameters in the waters of Shatt Al-Basrah channel at the selected stations S<sub>1</sub>, S<sub>2</sub> and S<sub>3</sub> are shown in figures 2-22. Seasonal changes in pH showed high difference between Aut. 2009 and Spr. 2010 at S<sub>3</sub>, Fig.2. Highest value of salinity was reported at S<sub>3</sub>, and they increased with relatively fixed rate from Aut. 2009 to Summer 2010, Fig. 3. Electrical conductivity has the same behavior of salinity, Fig.4. Reported turbidity in this study was high at S<sub>3</sub> and the highest values were recorded during Spr. 2010, Fig.5, due to blooming and phytoplankton's abundance, [13].




**Fig.2:** Seasonal variation of pH in the water of Shatt Al-Basrah Channel during the study period



**Fig.3:** Seasonal variation of Salinity in the water of Shatt Al-Basrah Channel during the study period




**Fig.4: Seasonal variation of Electrical conductivity in the water of Shatt Al-Basrah Channal during the study period**




**Fig.5: Seasonal variation of Turbidity in the water of Shatt Al-Basrah Channal during the study period**

Total Dissolved Solids (TDS) are high in S<sub>3</sub>, Fig.6, and the lowest values at all studied stations were during Winter 2010 probably due to dilution as a result of raining. Chloride is behave in the same trend of the TDS, Fig.7.




**Fig.6: Seasonal variation of Total Dissolved Solids in the water of Shatt Al-Basrah Channal during the study period**



**Fig.7: Seasonal variation of Chloride in the water of Shatt Al-Basrah Channal during the study period**

For calcium, there were some alternatives at stations 2 and 3 while it is decreased heavily at station 3 from Autumn 2009 towards Winter 2010, Fig.8, due to the transfer of soil and sewage water treated by air from Hamdan waste treatment station which through its waste towards Shatt Al-Basrah Channal. Magnesium on the other hand seemed to be fixed at all stations during the study period, Fig.9. Concentrations of bicarbonates are oscillating at station 3 as they decreased during Winter 2010 and increased during Spring 2010 then they decreased during

Summer 2010, while the oscillation of bicarbonates are limited and undetectable at stations 1&2 during all seasons, Fig.10. At all studied stations, sulphates are decreased during Winter 2010 and gradually increased during Spring and Summer 2010, Fig.11. Sodium and Potassium have the same behavior at all stations, their concentrations decreased during Winter 2010 then increased during Spring and Summer 2010, and the lowest values reported for both ions were at S<sub>3</sub> during Winter 2010, Figs 12&13 respectively.



**Fig.8: Seasonal variation of Calcium in the water of Shatt Al-Basrah Channal during the study period**




Fig.9: Seasonal variation of Magnesium in the water of Shatt Al-Basrah Channal during the study period




Fig.10: Seasonal variation of Bicarbonate in the water of Shatt Al-Basrah Channal during the study period






Fig.11: Seasonal variation of Sulphate in the water of Shatt Al-Basrah Channal during the study period



**Fig.12: Seasonal variation of Sodium in the water of Shatt Al-Basrah Channal during the study period**



**Fig.13: Seasonal variation of Potassium in the water of Shatt Al-Basrah Channal during the study period**

The seasonal variations of nutrients (phosphates, nitrites, nitrates and silicates) in the waters of Shatt Al Basrah channel during the study period are shown in figures 14 – 17 respectively. Lowest values reported for phosphates were during Spring 2010 at all stations, moreover, S2 was reported to be higher in phosphate contents during all seasons, Fig.14. For nitrate, there was an interferences in concentrations among the seasons as well as stations. During Spring nitrite was higher at Stations 1&2 and during Autumn 2009

nitrite was high at all stations, while during Summer, nitrite was low at all stations, Fig.15. At stations 2&34 nitrate showed the same behavior of nitrite, it exchange alternatively during Autumn and Spring seasons, while at S1 values of nitrates were low during all seasons of study, Fig.16. Silicates were reported as higher concentrations at all stations of Shatt Al-Basrah Channel during all seasons of study and ranged 70 – 100 ug Si-SiO<sub>2</sub>/ml, Fig.17.




Fig.14: Seasonal variation of Phosphate in the water of Shatt Al-Basrah Channel during the study period




Fig.15: Seasonal variation of Nitrite in the water of Shatt Al-Basrah Channel during the study period





Fig.16. Seasonal variation of Nitrate in the water of Shatt Al-Basrah Channel during the study period






**Fig.17. Seasonal variation of Silicate in the water of Shatt Al-Basrah Channel during the study period**

The seasonal variations of trace metals (Fe, Cu, Pb, Co, and Ni) in the waters of Shatt Al-Basrah channel during the study period are shown in figures 18 – 22 respectively. The levels of trace elements in the water of Shatt Al-Basrah Channel during the period of study were alternative in the studied stations and higher levels for Fe, Pb, Co, and Cu in station 2, while Ni was higher in station 3, this may be due to the exposure of station 2 to

various types of pollutants mostly is the sewage water from Basrah city [13].

The none availability of geochemical studies is the main factor to explain the different compositions of trace elements in the study area[14]. As Shatt Al-Basrah Channel is an extinction of Al Hammar marsh, Southern Iraq, the characterization of Al-Hammar marsh will reflected upon the water properties of Shatt Al-Basrah Channel [15].



**Fig.18. Seasonal variation of (Fe) in the water of Shatt Al-Basrah Channel during the study period**




Fig.19. Seasonal variation of (Cu) in the water of Shatt Al-Basrah Channal during the study period




Fig.20. Seasonal variation of (Pb) in the water of Shatt Al-Basrah Channal during the study period





Fig.21. Seasonal variation of (Co) in the water of Shatt Al-Basrah Channal during the study period



**Fig.22. Seasonal variation of (Ni) in the water of Shatt Al-Basrah Channel during the study period**

### CONCLUSION

As a conclusion the distribution of different types of pollutants in the waters of Shatt Al-Basrah Channel is liable to alternate due to discharging of waste water directly to the channel with out any treatment or even reduction in concentrations[16].The major factor for pollution in the Shatt Al-Basrah Channel is due to waste water from nearby sources, because water flow from marsh land through out the channel has no effect [17].

### CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

### REFERENCES

- Sargaonkar, A, Deshpands, V. Development of an overall index of pollution for surface water based on a general classification scheme in Indian contex. *Environ Monitor Asses* 2003; 89(1): 43-67.
- UNEP. The Mesopotamian Marshland: Demise of an Ecosystem. UNEP/DEWA/GRID-Genava; 2001, p1.
- Carayo C C, Culango KC, Estraro D, Largo DB, Dy DT. Correlation of inorganic nutrients to the standing group of "Green Tide" Algae during the peak growth season in Eastern Mactan Island, Central Philippines. *Philipp: Scient*; 2005, p1.
- Meyback M. Carbon, Nitrogen and Phosphorous transport by Worl river. *American J Sci* 1996; 282:401-450.
- Alexanderm R B, Smith. Trends in the nutrient enrichment of U.S.rivers during the late 20<sup>th</sup> century and their relation to changes in probable stream trophic conditions. *Limnol Oceanograph* 2006; 51:639-654.
- Meyback M, Helmerm R. The quality of rivers – from pristine stage to global pollution. *Global Plaet Chang* 1989; 75:283-309.
- Duan SW, Xu F, Wang LJ. Long term changes in nutrient concentrations of the changing river and principle tributaries. *Biogeochem* 2007; 85:215-234.
- Muller B, Berg M, Yao ZP, Zhang XF, Wang D, Pfluger A. How polluted is the Yangtze river? Water quality downstream from the three Gorges Dam. *Sci Total Environ* 2008; 402: 232-247.
- Standard Methods. Standard methods for the examination of water and waste waters. Michigan, USA: APHA/WPCF/AWWA; 1995.
- Murphy J, Riley JP. A modified single solution method for the determination of phosphate in natural water. *Anal Chem Acta* 1962; 27:31-36.
- Al-Obaidy AHM, Maulod BK, Kadhem AJ. Evaluating of raw and treated water quality of Tigris river within Baghdad by Index Analysis. *J Water Resource Protec* 2010; 2: 629-635.

12. Al-Imarah FJM, Al-Shawi AJM, Al-Shawi IJM. Some physic-chemical environmental parameters, chlorophyll a and primary productivity in Shatt Al-Basrah and Khawr Al-Zubair, Southern Iraq NW Arabian Gulf: A Survey Study 1980-2005. *Uruk Sci J* 2011; 4(2):30-38.
13. Al-Khafaji BY, Dawood YT, Maktoof AA. Trace metals distribution in fish tissues (*Cyprinus carpio* and *Barbus luteus*) and sediments Al-Marsh south of Iraq. Conference on Environmental Sciences 4th 2012 – Babylon University; 2012, p 180.
14. Lenova GA, Anoshin GN, Bychinskii VA, Shcherbov BL, Strakhovenko VO. Landscap-geochemical patterns of distribution of heavy metals in biota and bottom sediments of Altai lakes. *Russian Geol Geolog Geophys Geophysika* 2002; 43(12):1080-1092.
15. Richardson C J, Hussain NA. Restoring the Garden of Aden: An Ecological Assessment of the Marshes of Iraq. *Bioscience* 2006; 56(6): 477-489.
16. Aziz NM. Studying of pollution by petroleum hydrocarbons and some trace metals in water, sediments, and some plants from Shatt Al-Basrah Channal, Iraq. Ph. D. Thesis, College of Science, Basrah University, Basrah; 2004,p 97.
17. Adam RS, Al-Shawi IJM, Al-Imarah FJM. Distribution of some chemical elements in the marsh land of Southern Iraq after rehabilitation. *Marsh Bulletin* 2007; 2(1):11-17.

**Cite this article as:**

Al-Imarah, F. J. M., Al-Jorany, Y. SH. J., Mizhir, A. A. Murugesan. Physico Chemical Characterization for Water of Shatt Al- Basrah Channel, 2009-2010. *J Pharm Chem Biol Sci* 2017; 5(3):165-176