

This article was downloaded by: [The Aga Khan University]
On: 13 March 2015, At: 22:31
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



[Click for updates](#)

Molecular Crystals and Liquid Crystals

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gmcl20>

Synthesis and Mesomorphic Properties of New Methylene-Linked Linear Symmetrical Liquid Crystal Dimers

Uhood J. Al-Hamdani^a, Aiman G. Al-Ibrahim^a, Hanna S. Abbo^b & Salam J. J. Titinchi^b

^a Department of Chemistry, College of Education, University of Basrah, Basrah, Iraq

^b Department of Chemistry, University of the Western Cape, Bellville, Cape Town, South Africa

Published online: 26 Feb 2015.

To cite this article: Uhood J. Al-Hamdani, Aiman G. Al-Ibrahim, Hanna S. Abbo & Salam J. J. Titinchi (2015) Synthesis and Mesomorphic Properties of New Methylene-Linked Linear Symmetrical Liquid Crystal Dimers, *Molecular Crystals and Liquid Crystals*, 607:1, 13-22, DOI: [10.1080/15421406.2014.927549](https://doi.org/10.1080/15421406.2014.927549)

To link to this article: <http://dx.doi.org/10.1080/15421406.2014.927549>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>



Influence of different linkages on the mesomorphic properties of aromatic ring system liquid crystals

Uhood J. Al-Hamdani, Nezar L. Shihabaldain, Duaa S. Al-Mayah, and Ahmed M. Jassem 

Department of Chemistry, College of Education for Pure Sciences, Basrah University, Basrah, Iraq

ABSTRACT

In order to study the effect of the central linkage on the mesomorphic behaviour of aromatic ring liquid crystals we have synthesized two homologues groups, group A includes compounds **1a-d** and group B includes compounds **2a-e**. Group A has azo, azomethine, azo-ester, and azomethine-ester central linkages whereas the group B has azomethine, ether-azomethine, heterocyclic, and methylene-azomethine central linkages. The synthesized compounds were identified and characterized by elemental analyses and spectroscopic methods (proton nuclear magnetic resonance and Fourier transform infrared). The mesomorphic properties were investigated by polarizing optical microscopy (POM) with a heating stage and differential scanning calorimetry (DSC) thermograms with an aim to investigate the influence of different types of linkages on the mesomorphic properties in such aromatic ring systems. The results revealed that the change of the central linkages has a crucial effect on the mesomorphic properties when compared to the influence of different types of linkages. We found that the synthesised compounds (**1a-d**) showed smectic phase with different thermal ranges, while the synthesised compounds (**2a-e**) showed nematic phase except the compounds **2c-e** which did not exhibit any liquid crystalline phase.

KEYWORDS

Aromatic ring liquid crystals; azo; azomethine; ester; linking groups

Introduction

Nowadays, thermotropic liquid crystal materials offer foreseeable applications as they are the basis of displays in mobile telecommunication, storage units, biomedical diagnosis, and computing devices [1–4]. The thermotropic liquid crystal molecules that are used in these devices almost have a rod-like shape [5]. The rod- or disk-like shaped molecules used to be a fundamental prerequisite for conventional formation of thermotropic liquid crystal molecules due to the steric packing considerations which play an important role in this interesting state of soft matter [6]. In the rod-like liquid crystal molecules with linear structure, these molecules are required to have certain rigid structures [7].

Generally, a proper selection of different building blocks such as core moieties, linking groups and terminal substituents is an effective strategy for designing liquid crystal

CONTACT Ahmed M. Jassem  ahmed.majedd@uobasrah.edu.iq  Department of Chemistry, College of Education for Pure Sciences, Basrah University, Basrah, Iraq.

© 2021 Taylor & Francis Group, LLC