

## New records algal species from the Shatt Al-Arab River, Southern Iraq

A.A.A. Alwaeli\* and A.M. Athbi<sup>2</sup>

<sup>1</sup>Marine Science Center, <sup>2</sup>College of Education for Pure Sciences,  
University of Basrah, Iraq

\*e-mail: ali69\_basra@yahoo.com

(Received: 4 March 2021 - Accepted: 16 June 2021)

**Abstract** - Shatt Al-Arab River is an important inland river in Iraq. It is considered an important source of fresh water for the city of Basrah, as it is used for drinking and orchards irrigation also used for industrial and commercial purposes. Sampling was carried out monthly from November 2019 to December 2020 from Shatt Al-Arab River. The present study made records nine of algae species for the first time in Iraq. Represented by six species of Chlorophyta (*Eremosphaera tanganyikae*, *Microspora wittrockii*, *Monoraphidium komarkova*, *Mychonastes pusshpae*, *Scenedesmus similagineus* and *Spongiococcum tetrasporum*), one species of Cyanophyta (*Pseudanabaena catenata*), one species of Dinophyta (*Impagidinium paradoxum*), and one species of Euglenophyta (*Strombomonas verrucosa*).

**Keywords:** Algal species, New record, Shatt Al-Arab River.

### Introduction:

Shatt Al-Arab River is one of the most important hydrographic features in the northwest of the Arabian Gulf, as this River supplies waters to the Arabian Gulf with an amount of 5 x 910 m<sup>3</sup> of freshwater annually, and the effect of this water is observed at a distance of 5 km from the estuary of the river in the Arabian Gulf. Shatt Al-Arab's water is usually rich in organic matter and nutritious salts, which increases the fertility of the waters of the northern part of the Arabian Gulf (Al-Lami 1986). The first environmental study conducted on the Shatt Al-Arab River was that of Mohammad (1965) which is followed by a few limited studies (Al-Kaisi, 1970; Antoine, 1977; Hameed, 1977; Pankow *et al.*, 1979; Al-Saadi *et al.*, 1979 and; Maulood *et al.*, 1981). Our knowledge of the phytoplankton especially in the Shatt Al-Arab River is scarce and incomplete.

Therefore, this study forms an attempt to diagnose some unregistered species of algae that had not been previously recorded to support the diversity of this important waterway and a complement to previous studies.

### Materials and Methods:

Shatt Al-Arab River is originated from the junction of the Tigris and Euphrates River at Al-Quarna town north of Basra City and drain into the Arabian Gulf at latitude 57° 99' North and longitude 35° 48' South with a total length of about 204 km (Fig. 1) (Al-Mahmood, 2012). Shatt Al-Arab River is affected by the climatic or natural conditions that prevail in the region, and the width of the Shatt Al-Arab River ranges from 400 meters to more than two kilometers (Al-Assadi, 2015).

Spread on both sides of the River many palm groves and field crops which affect the quality of the water through the flow of many fertilizers and pesticides into the River in addition to the sewage water that directly flows into it.

Five stations were chosen from different areas of the River Al-Deer, Al-Ashar, Abu-Alkhaseeb, Al-Seeba and Al-Fao. Sampling collection was done monthly from November 2019 to December 2020 from Shatt Al-Arab River, southern Iraq by using a phytoplankton net (mesh size 20  $\mu\text{m}$ ), with 20 cm mouth openings made by the American company Wild. The net was towed behind a boat steam at an average speed for 10-15 minutes at the middle of the River.

The contents of the net were collected in a Plastic bottle of a capacity of 100 ml and fixed with 4% formalin and labeled. Upon conducting the examination process, the sample is concentrated with a centrifuge to reduce its size, then one or two drops are taken from it and placed on a slide (Stein, 1973), then a cover slip is placed on it and left for two minutes for settling, then examined under a compound light microscope type Zeiss with camera Axio Cam ICc 3 type Zeiss (x 400, Scale 10  $\mu\text{m}$ ). Identification of algae are carried out with the aid of the following taxonomic literatures: Desikachary, 1959; Prescott, 1968, Prescott, 1970; Al-Saboonchi *et al.*, 1990.




Figure 1. A map showing the locations of sampling collection sites.



It was also noticed that most of the species were recorded in the first, second and third stations, while the last two stations had a few numbers of species, and this is because the speed of the current and the availability of light are among the most influencing factors in the composition of society in fast-flowing environments (Keithan and Lowe, 1985), because such environments are usually of a lower density in terms of the number of cells and most of the species are of the adherent type compared to the slow flowing environments. During the study period, some species showed their dominance over others the dominance of any species in any environment depends on the quality of the species in its competition with other species (Tilman, 1977).

Many taxonomic studies dealing with phytoplankton were conducted on the Shatt Al-Arab, but those with that gave a description of the species and drawings by means of Camera Lucida or by photography are very rare, such as that of Al-Handal *et al.* (1989) who studied the occurrence of some filamentous algae on the sediments of Shatt Al-Arab River and one of its side canals.

The second study was carried out by Al-Saboonchi *et al.* (1990) in which samples were collected monthly from January 1985 to June 1987 from Shatt Al-Arab River and some of its canals. Fifty-three species were recorded. Twenty four of which were found to be new records for the Shatt Al-Arab region, sixteen species were new recorded for Iraq. In the present study we are dealing with those species which were not previously described. Chlorophyta is represented by 9 species, Cyanophyta by 5 species and Xanthophyta by 2 species.

Representative of the Trebouxiophyceae genus *Eremosphaera* (with the exception of *E. viridis*) showed great variation in their morphology, and some of the described species, varieties and forms are still of uncertain taxonomic position (Stoyneva, 2006). In the present study, *Eremosphaera tanganyikae* was recorded at all months except in June (Table 1), and was recorded in the first 4 stations during January and February, while they were recorded in the second, third and fourth stations during March and May. However *Microspora wittrockii* was recorded only once throughout the study period, during February at the second station only.

Another rare species, *Microspora floccosa* (Vauch), was only reported by Al-Kaisi (1970, 1976) from some water systems in the middle and south of Iraq, but not in the Shatt Al-Arab River, and reported by Al-Handal *et al.* (1989) from Shatt Al-Arab River. Moreover, in the current study *Monoraphidium komarkova* was recorded at all stations during January, while the other species were very rare, and were recorded once or twice throughout the study period.

Nine algal species of phytoplankton were identified from Shatt Al-Arab River for the first time and they were not recorded before in the Iraqi aquatic environment, and represented by:

*Eremosphaera tanganyikae*. Stoyneva Gartner & Vyverman, 2006.

Stoyneva, vol. 139, p. 3-13.

Description:

This is a freshwater species, its shape is elongated oval, solitary cells, surrounded by a thick cell wall, lobed from the inside, and has no flagella, chloroplasts are surrounded by a membrane, which separates them from the cell wall, cell 13.5-14.5  $\mu\text{m}$  in width and 20-30  $\mu\text{m}$  long, wall thickness from the top about 3-4  $\mu\text{m}$ , and from the sides 1.5-2.5  $\mu\text{m}$ .

Division: Chlorophyta

Class: Trebouxiophyceae

Order: Chlorellales

Family: Oocystaceae

Genus: *Eremosphaera*

Species: *Eremosphaera tanganyikae* Stoyneva, 2006.

*Microspora witrockii* (Wille) Lagerheim, 1887.

Skuja, 7; Figs. 119, p. 25-85.

Description: Chlorophyta

Multicellular filamentous algae, consisting of large cells with clear borders, this species is distinguished by a golden color, the filament-forming cells are rectangular in shape 15-18  $\mu\text{m}$  in length, and 12-14  $\mu\text{m}$  in width, the cytoplasm is dense and fills the cell except for a vacuole near the transverse wall, live in freshwater.

Division: Chlorophyta

Class: Chlorophyceae

Order: Ulotrichales

Family: Microsporaceae

Genus: *Microspora*

Species: *Microspora wittrockii* (Wille) Lagerheim, 1887.

*Monoraphidium komarkova* Nygaard, 1979.

Hindak, F. (1988).34(9) 1-264.

Description:

Single cells, with tapering ends, the middle of the cell is wider than at the ends, usually its shape is straight, the plastid extends along the cell and has a bright green color, the length of the filament is approximately 44-50  $\mu\text{m}$ , and its width in the middle reaches 2.5-3  $\mu\text{m}$ .

Division: Chlorophyta

Class: Chlorophyceae

Order: Sphaeropleales

Family: Selenastraceae

Genus: *Monoraphidium*

Species: *Monoraphidium komarkova* Nygaard, 1979.

*Mychonastes pusshpae* Krientiz, C. & Proschold, 2011

Krientiz, 50(1), p: 89-106.

Description:

The type species (holotype) of the genus is *Mychonastes ruminates* P.D. Simpson & S.D. Van Valkenburg. This is a freshwater species, No synonyms are currently included in algaebase, The alga is a colony of six spherical cells that are linked together by plasma ligands, cell diameter 3-4  $\mu\text{m}$ .

Division: Chlorophyta

Class: Chlorophyceae

Order: Sphaeropleales

Family: Mychonastaceae

Genus: *Mychonastes*

Species: *Mychonastes pusshpae* Krientiz, 2011

*Scenedesmus similagineus* Hortobagyi, 1960 (Fig. 5)

Hortobagyi, 1-345-381, pl. 64-75.

**Description:**

The type species (holotype) of the genus is *Scenedesmus obtusus* Meyen. No synonyms are currently included in algaebase, this is a freshwater species, colony 2-celled. Individual cells elongated ovoid to spindle shaped poles rounded, with small spines of about 4.68  $\mu\text{m}$  long it's cell body long about 17 to 19  $\mu\text{m}$ , and 7.5 to 8.5  $\mu\text{m}$  wide.

Division: Chlorophyta

Class: Chlorophyceae

Order: Sphaeropleales

Family: Scendesmaceae

Genus: *Scenedesmus*

Species: *Scenedesmus similagineus* Hortobagyi, 1960.

*Spongiococcum tetrasporum* Deason, 1959.

Deason; 574, Figs. 1-9

**Description:**

This is the type species (holotype) of the genus *Spongiococcum*, No synonyms are currently included in algaebase. Colonies Spherical, cells, 10-20  $\mu\text{m}$  in diameter. A sponge like chloroplast, with a pyrenoid usually situated in the center, large and fragmented pyrenoid of which are polygonal and somewhat compact.

Division: Chlorophyta

Class: Chlorophyceae

Order: Chlamydomonadales

Family: Chlorococaceae

Genus: *Spongiococcum*

Species: *Spongiococcum tetrasporum* Deason, 1959

*Pseudanabaena catenata* Lauterborn, 1915.

Lauterborn, 1915: 432, pl. III, Fig. 27

**Description:**

Short unbranched filaments consisting of 10 to 20 elongated cells, flexible, close to each other, the color of the filaments is dark blue to olive, the length of the cell is 4-6  $\mu\text{m}$ , and the width is 2.5-3.5  $\mu\text{m}$ .

Division: Cyanophyta

Class: Cyanophyceae

Order: Synechococcales

Family: Pseudanabaenaceae

Genus: *Pseudanabaena*

Species: *Pseudanabaena catenata* Lauterborn, 1915.

*Impagidinium paradoxum* Wall, 1967.

Wall, pl. 15, Figs. 5-8, p. 106-107; Zonneveld, 39(3), p. 387-407.

**Description:**

Solitary cell, ovoid in shape, protruding from its outer surface several short spines, the length of the cell body 35-45  $\mu\text{m}$  and 25-35 its width, The front of the body is slightly protruding, the surface of the body is covered with granules, marine and freshwater species.

Division: Dinophyta

Class: Dinophyceae

Order: Gonyaulaceae

Family: Gonyaulacales

Genus: *Impagidinium*

Species: *Impagidinium paradoxum* Wall, 1967.

*Strombomonas verrucosa* (E. Daday) Deflandre, 1930

Araujo, vol.6, pl.(1)-203, Figs. 306.; Hu, H. pp.(4 pls. of 16 Figs.), (i-iv), i-ix, 1-1023

Description:

Solitary cells, conical in shape, the anterior pole is usually round and prominent 2.5-5  $\mu\text{m}$  in length, and the back pole is slightly extended 2-3.5  $\mu\text{m}$ , the cytoplasm is lobular and fills the cell, length of the body is 41-43  $\mu\text{m}$ , and it's width 25-27  $\mu\text{m}$ . Some zigzags you can be seen in the outer wall, this is a freshwater species.

Division: Euglenophyta

Class: Euglenophyceae

Order: Euglenida

Family: Euglenidae

Genus: *Strombomonas*

Species: *Strombomonas verrucosa* (E. Daday) Deflandre, 1930.



*Eremosphaera tanganyikae*



*Microspora wittrockii*



*Monoraphidium komarkova*



*Mychonastes pusshpae*



*Scenedesmus similagineus*



*Spongiococcum tetrasporum*



*Pseudanabaena catenata*



*Impagidinium paradoxum*



*Strombomonas verrucosa*

Figure 1. Photographs of the Planktonic algal species recorded for the first time from the Shatt Al-Arab River, Basrah, Iraq.

**Reference:**

- Al-Assadi, S.A., Al-Mahmood, H.K. and Abdulla, S.C. 2015. Estimation of the minimum net water discharge in the Shatt Al-Arab (Southern Iraq). Basrah Literature M., Issue 72: 285-292 p.
- Al-Handal, A.Y., Al-Assdi, S.A. and Al-Mukhtar, M.A. 1989. Occurrence of some filamentous Algae in the River Shatt Al-Arab, Iraq. *Marina Mesopotamica*, 4(1): 67-81.
- Al-Kaisi, K.A. 1970. Introductory study on the algae of mid and south Iraq. *Bulletin of the College of Science, University of Baghdad*, 11: 45-80.
- Al-Kaisi, K.A. 1976. Contribution to the algal flora of the rice fields of southern Iraq. *Nova Hedwigia*, 27: 813-827.
- Al-Lami, A.A. 1986. Ecological study on phytoplankton on some marsh regions Southern Iraq. M.Sc. Thesis, College of Science, University of Basrah, 98p (In Arabic).
- Al-Mahmood, H.K.H. 2012. Study the nature of the volume of discharge and the concentration of total soluble substances of Shatt Al-Arab, Iraq. *Mesopotamian Journal of Marine Science*, 23(1): 34-40.
- Al-Saadi, H.A., Pankow, H. and Huq, M.F. 1979. Algological investigation in the polluted Ashar canal and Shatt Al-Arab in Basrah. Iraq. *Int. Revue Ges. Hydrobiol.*, 64(4): 527-540.
- Al-Saboonchi, A.A., Ghani, A.A. and Abdul Hussein, M.M. 1990. Non-Diatom Algae from the Shatt Al-Arab River, Iraq. *Marina Mesopotamica*, 5(1): 89-126.
- Antoine, S.E. 1977. Seasonal variations of the environmental characteristics and phytoplankton blooms of Tigris, Iraq. M.Sc. Thesis, University of Basrah, 91p.
- Araujo, G.J.M. and Bicudo, C.E.M. 2018. Flora ficologica deestado desao Paulo. Vol.6 Euglenophyceae. Pp(i)-viii, (1)-203, plus 306 Figs. Sao Paulo: Trima: Fapesp.
- Deason, T.R. 1959. Three new Chlorophyceae from Alabama soil. *American Journal of Botany*, 46: 572-578.
- Desikachary, T.V. 1959. Cyanophyta Indian. *Concil of Agricultural Research*. New Delhi, India, 686p.
- Hameed, H.A. 1977. Studies of the ecology of phytoplankton of Shatt Al-Arab River at Basrah, Iraq. M.Sc. Thesis, University of Basrah, Iraq, 102p.
- Hidak, F. 1988. Studies on the chlorococcal algae (chlorophyceae) lv. *Biologicke prace* 34(1/2): 1 264.
- Hortobagyi, T. 1960. Algen aus den fishteichen von buzsak. ii. *Scendesmus* arten . *Nova hedwigia* 1: 345-382, Pl. 64-75.
- Hu, H. and Wei, Y. 2006. The freshwater algae of China. *Systematics, taxonomy and ecology*. (4 pls. of 16 Figs.), (i-iv), i-xv, 1-1023, China.
- Keithan, E.D. and Lowe, R.L. 1985. Primary productivity and spatial structure of phytolith growth in stream in the Great Smoky mountains National park, Tennessee. *Hydrobiol.*, 123: 59-67.
- Krienit, T. 2011. Taxonomic reassessment of the genus *Mychonastes* (Chlorophyta) including the description of eight new species. *Phycologia*, 50(1): 89-106.
- Lauterborn, R. 1915. Saproelische lebewelt Ein Beitrag zur biologia des faulschlammes natuerlicher gewasser. *Verhandlungen des naturhistorisch-medizinischen vereins zu Heidelberg*, Ser. 2(13): 395-481, pl. III.
- Mohammad, M.B. 1965. Preliminary observations on some chemico-physical features of the Shatt Al-Arab estuary. *Proc. Iraq Scient. Soc.*, 6: 34-40.



- Maulood, B.K., Hinton, G.C.F., Whitton, B.A. and Al-Saadi, H.A. 1981. On the Algal Ecology of the Lowland Iraqi Marshes. *Hydrobiologia*, 80: 269-276.
- Pankow, H., Al-Saadi, H.A., Huq, M.F., Hadi, R.A.M. 1979. On the algal flora of the marshes near Qurnah (Southern Iraq). *Willdenowia*, 8(3): 493-506.
- Prescott, G.W. 1968. Algae of the Western great lake area. Ellion C., Brown Co. Pub., Dugugue, Iowa, USA, 977p.
- Prescott, G.W. 1970. How to know the freshwater algae 2<sup>nd</sup> ed. McGraw-Hill Book Co. New York, 719p.
- Skuja, H. 1934-1933. Beitrag zur Algenflora lettlands. I. *Acta Horti Botanici Universitatis Latviensis*, 7: 25-85, 119 Figs.
- Stein, J.R. 1973. Hand book of phycological methods. Cambridge Univ. Press., Cambridge., UK.
- Stoyneva, M., Gartner, G., Coquyt, C. and Vyverman, W. 2006. *Eremosphaera tanganyikae* sp. Nov. (Trebouxiophyceae), a new species from Lake Tanganyika. *Belgian Journal of Botany*, 139: 3-13.
- Tilman, D.L. 1977. Resource competition between phytoplanktonic algae: An experimental and theoretical approach. *Ecology*, 58: 338-348.
- Wall, D. 1967. Fossil microplankton in deep-sea cores from the Caribbean Sea. *Palaeontology*, 10(1): pl. 14-16, p. 95-123.

## تسجيل تسعة أنواع طحلبية جديدة في نهر شط العرب، جنوب العراق

عقيل عبد الصاحب عبد الحسين<sup>1</sup> و أحمد محسن عذبي<sup>2</sup>

<sup>1</sup> مركز علوم البحار، <sup>2</sup> كلية التربية للعلوم الصرفة، جامعة البصرة، العراق

**المستخلص** - جمعت العينات شهريا من كانون الثاني الى كانون الاول 2020 من نهر شط العرب، وسجلت الدراسة الحالية تسعة انواع من الهائمات النباتية لأول مرة في العراق، وتم اعطاء وصف وابعاد كل نوع من الانواع المشخصة، وتمثلت الطحالب الخضراء بـ 6 انواع ونوع لكل من الطحالب الخضراء المزرقية والطحالب الدوارة والطحالب اليوغلينية، كما تناولت الدراسة ظهور ووجود الانواع قيد الدراسة خلال اشهر السنة المختلفة في مياه النهر.