

**First Study** As part of a study on lung adenocarcinomas in older laying hens, 150 commercial egg-laying chickens were necropsied at 3 years of age. Tumors were identified in 67 of the hens. Grossly visible metastatic tumors in the lungs of a few birds prompted collection of lungs from 50 affected hens for histologic examination. Metastatic adenocarcinomas were identified histologically in the lungs of 9 of the 50 hens. The number of tumors and extent of involvement of the lung parenchyma varied from minimal to extensive. Tumor emboli were occasionally seen in pulmonary vessels. Metastatic foci showing expansive or invasive growth were identified. These findings indicate that vascular spread of adenocarcinomas is more common than previously thought in chickens.

**Second Study** Brown egg-laying hens are highly susceptible to amyloid arthropathy as a result of *Enterococcus faecalis* and *Mycoplasma synoviae* infections. Other frequently associated bacteria include *E. coli*, *Salmonella enteridis*, and *Staphylococcus aureus* have been associated with amyloidosis in chickens.



Dr. Bahaa A. Alsereah, Professor (Assistant) in the Dep. of Veterinary Public Health, Faculty of Veterinary Medicine at the University of Basrah, born in 1978. He work in field of Poultry Nutrition, Published 36 Research Paper in a number of International Journals and 13 Book Published in Germany.



Bahaa Al-Sereah  
Hanan Al-Rashid  
Dhuha Al-Lami

## Pulmonary Tumor and Amyloidosis in Commercial Laying Hen

light microscopic study

