

Composition of fish assemblages in the southern part of the Main Outfall Drains, Southern Iraq

Abdul Hussein Jaafer Abdullah^{1*}, Sajad Abdul Gani Abdullah², Ali Taha Yaseen¹

1. Department of Marine Vertebrate, Marine Science Centre, University of Basrah, Iraq

2. Department of Biology, College of Education-Qurna, University of Basrah, Iraq

* Corresponding author's Email: abdulhassain.abdulah@uobasrah.edu.iq

ABSTRACT

Fish assemblage was investigated in the southern part of the Main outfall drains, southern Iraq from January to December 2019. Monthly variations were recorded in water quality parameters including water temperature, dissolved oxygen, salinity, electric conductivity, turbidity, hydrogen ion, and total dissolved solids. A total of 10981 fish specimens belonging to 27 fish species, 25 genera, 17 families, and eight orders were collected from the southern part of the Main outfall drains. Marine fish species were 19 and eight species from freshwater. Three species were presiding the numerical relative abundance of species formed 80.96% of the total number of species. In this region, *Acetrogobius dayi* represented 32.00%, *Boleophthalmus dussumieri* 28.92% and *Aphanius dispar* 20.04% of the total number of species. The status of ecological indices were poor for diversity index (H), richness index (D) was troubled, and semi-balanced for evenness index (J). The present study showed bad water quality after analyses of physicochemical variables, and also sensory indicators (colour and smell) of the water which play a major role in the blending composition of the fish community structure. The results revealed that salinity has a large contributor to determine the nature spread and distribution of the fish populations. The study exhibited that the river habitat is unstable, affected by three factors: 1) Quantity and quality of freshwater coming from Al-Siphone regulator in Thi-Qar Province; 2) The tidal process spread organic pollution that are dumped in Hamdan region, Southern Basrah City; and 3) The chemical and organic pollutions resulting from reclamation of agricultural lands.

Keywords: Fish assemblage, Indicators, Water Quality, Main outfall drain

Article type: Research Article.

INTRODUCTION

Fish populations have been always utilized as ecological indicators for evaluating and measure of water deterioration in the diverse spatial and temporal variations of water bodies (de Mello *et al.* 2018; Jiang *et al.* 2020). All specialists in fish population studies realize that there is a dialectical correlation between the quality and nature of fish assemblages as well as the water quality (Huang *et al.* 2019). Through the presence of specific species and the nature of the fish community structure, we can predict water quality and vice versa (Zeni & Casatti 2014). Fish assemblages are widely used to know the alterations in the fish community structure, although their ecosystems were modified (Ngor *et al.* 2018). The long-term variations which are correlated with stress from slight incremental alterations in environmental circumstances, lead to alterations in the abundance of fish fauna, and provide an important practical assessment of the impacts of water quality on the fish assemblages of fish (Parker *et al.* 2015). Freshwater ecosystems are among the most changeable systems due to anthropogenic activities and include the hotspot area of biological diversity. These ecosystems encompass a large number of endemic species which are seriously affected by urban pollution, originate from populations growth. Most of their problems are represented by water ecosystem modification, industrial waste, agricultural process, habitat alteration, modifications of hydrology and alien species (Huang *et al.* 2019; Pyron *et al.* 2019). Water

contamination and hydrological modification are recognized as the most real threats to freshwater fauna (Pyron *et al.* 2019). Freshwater habitats are more vulnerable to human threats than terrestrial and marine ecosystems, and maintain water resources require knowledge of the effects and historical trends of water, as well as using land to plan suitable solutions for the conservation of the waters (Craig *et al.* 2017; Poff *et al.* 2018). Pyron *et al.* (2019) mentioned that the composition of an assemblage of fish (taxonomic structure) was related to variations in land use. However, assemblages' species composition is correlated with modification of hydrological factors, reduces minimum discharge limits, declines the filling rate, and increased level rates. Generally, the highest diversity values were recorded in the tropical region, whereas in Asia, the level of diversity for families (121 families) was higher than in Africa, and 50 to 55 families were found in inland waters of Latin America. The estimated number of fish species identified is about 34300 fish species over the world, of these species nearly 15,666 species are recorded in freshwater, while other 15,562 in marine water and 3,115 in brackish or are diadromous (Froese & Pauly 2022). Systems of fish assemblage's approximation and divergence (genotype) are based on environmental circumstance variations passed across numerous spatial and temporal scales (Bower & Winemiller 2019). The large harmful impacts on fish populations in the rivers and streams in the desert or semi-desert regions come from the changes in hydrology that alter the ecosystem directly or indirectly and stimulate the changes in propagation and distributions of exotic species (Yang *et al.* 2019). The amount of shifting in riverine fish populations may depends on spatial parameters and the initial resemblance in the fish population structure happened before anthropogenic alterations. It is also related to the extent of habitats which is important to understand the impacts of variations in accumulation on fish assemblages (Taylor *et al.* 2018; Mazareiy *et al.* 2019; Sattari *et al.* 2019; Forouhar Vajargah *et al.* 2020a,b). There is no study discussing the relationship between fish populations and water quality in southern Iraq, but there were several investigations implemented on the Main outfall drains and Shatt Al-Basrah canal, which are taken into account extension to the Main outfall drains include fish diversity (Yousif, 1986; Al-Daham & Yousif 1990; Jasim 2003; Younis & Al-Shamary 2011; Younis & Al-Shamary 2012; Abdullah *et al.* 2018). The aim of the present study is to investigate the effect of water quality represented by physicochemical parameters and demonstration of anthropogenic activities' impacts on the nature of building fish assemblage on the southern part of the main outfall drain in southern Iraq.

MATERIALS AND METHODS

The main outfall drain is called the third river in Iraq, which is stretched to a distance of 565 km from the northwest of Baghdad City to the north of the Persian Gulf (Abdullah *et al.* 2018). The present study area extends about 30 km from the Al-Gttra region N 30° 40' 12" E 47° 34' 12" to the south of Al-Khora village N 30° 34' 12" E 47° 41' 24" about five km. The position of study area was determined by Global position system (GPS). The river width is 110 m, but the submerged area can be increased due to the status of the tide and the amount of water provided from the Siphone regulator in Thi-Qar Province to reach one or two km in some places (Al-Munshed 1998; Fig. 1). Some environmental variables were verified simultaneously with the fishing processes. Water temperature (°C) was measured by a mercurial thermometer (-10 to 100 °C), salinity (g L⁻¹) and hydrogen ion was estimated by Lovibond-Sensor Direct 150 manufactured in Germany; total dissolved solids (TDS) in mg L⁻¹ and electrical conductivity (EC) in mS cm⁻¹ were estimated directly by Hanna instruments (a waterproof HI-9146). Turbidity was calculated in turbid meter HI- 93703C in Nephelometric turbidity (NTU). Dissolved oxygen (mg L⁻¹) was determined according to Welch (1964). Samples were monthly collected from the region during the study period from January to December 2019 by several fishing methods, drift gillnets, fixed gillnets, seine nets, cast nets as well as observations from fishermen. Fishes were identified according to Fricke *et al.* (2021) and the rest of the scientific names were updated according to Froese & Pauly (2022). Analyses of fish assemblage were performed as follows:

The measurement of relative abundance was performed according to Walag *et al.* (2016). Fish diversity was calculated by Huang *et al.* (2019); evenness and richness based on the methodology of Nyitrai *et al.* (2012). To analyse the relationships among physicochemical factors and species, we used Principal Component Analysis (PCA) program, Canonical Correspondences Analyses (CCA), and Statistical Package for Social Science, Ver. 20 (SPSS).

RESULTS AND DISCUSSION

Physicochemical parameters have an essential role in forming and building a diversity of fish assemblage in riverine ecosystems. The monthly variations in the properties of the ecological variables were illustrated in Fig.

2. The minimum value of water temperature was 15 °C in January and started to increase gradually to reach the maximum of 35 °C in July with an average of 24.67 °C ± 6.83, thereafter declined to December. Dissolved oxygen values set out to be slightly high in the cold months then descended in the hot months to record 6.2 mg L⁻¹ in July and elevated to 7.9 mg L⁻¹ in January, with the mean value of 7.13 ± 0.52 mg L⁻¹. The ranges of salinity were oscillating and average values were observed at 8.09 g L⁻¹ in April, while 45.80 g L⁻¹ in March (mean = 25.06 ± 10.84 g L⁻¹). Electrical conductivity was strongly related to salinity values recording 13.96 mS cm⁻¹ in April, while 71.70 mS cm⁻¹ in March (mean = 39.4596 ± 16.74 mS cm⁻¹). Turbidity started in tempered values to decline at the minimum point of 24.40 NTU in April then the maximum was 88.00 NTU in June and an average of 44.30 NTU ± 19.74. The hydrogen ion recorded relatively high values at the beginning of the study then the values fluctuated to record the lowest at 7.00 in September and the highest at 8.10 in January (mean = 7.42 ± 0.42). A total dissolved solid was observed in the first three months of the study and declined to 8934 mg L⁻¹ in April, while 45846 mg L⁻¹ in March (mean = 25151.83 ± 10751.19 g L⁻¹) and then the values fluctuated to December (Table 1).




Fig. 1. Map of the southern part of the main outfall drain, southern Iraq.




Fig. 2. Monthly variations in the physicochemical variables in the southern part of the Main Outfall Drain, southern Iraq.

The physicochemical properties of lotic ecosystems play an important role in the building of fish populations. Water temperature is the most important parameter due to directly effect on the level of dissolved oxygen in the ecosystems (Rajwa-Kuligiewicz & Rowiński 2015). The variations in temperature recorded in the present study area were due to different sampling times and monthly variance in temperature that influences the biological, biochemical, trophic, and reproduction of fishes (Camargo *et al.* 2021). The amount of DO is a determinant of water quality which is strongly related to temperature. The pollution of organic matter in the Main outfall drains

was high due to the river representing a carrier for heavy water, agricultural residues, and land-washing products. Therefore the degree of oxygen saturation of water is at its lowest status, as a result of the increased organic matter and other pollutants (Wen & Van De Giesen 2017).

Table 1. The variations in physicochemical parameters with range and mean values with standard deviation (\pm SD).

Parameter	Unit	Range	Mean \pm SD
Water temperature	$^{\circ}\text{C}$	15-35	24.67 \pm 6.83
Dissolved oxygen	mg L^{-1}	6.2-7.9	7.13 \pm 0.52
Salinity	g L^{-1}	8.09-45.80	25.06 \pm 10.84
Electric conductivity	mS cm^{-1}	13.96-71.70	39.45 \pm 16.74
Turbidity	NTU	25.50-76.11	44.30 \pm 19.74
Hydrogen ion	-	7.00-8.10	7.42 \pm 0.42
Total dissolved solids	mg L^{-1}	8934-45846	25151 \pm 10751

Salinity has played a major role in the distribution of fish fauna in aquatic ecosystems, its effects on primary production, geographical distribution, dispersal, recruitment of larval, and the marine species behaviour (Herbert *et al.* 2015; Santos *et al.* 2017). The Main outfall drains are human-managed, so the river is subject to constant shafting in the quantities of and qualities of freshwater released towards the south, from the Al-Siphon regulator in the Al-Nasiriyah City, as well as affected by marine tidewaters from the Persian Gulf. These changes created a distinct dynamic system, which has a great implication for the fish population's composition (Abdullah *et al.* 2018). The results of electric conductivity were in continuous dynamic changing, when the freshwater discharge from the Al-Siphon regulator decreases, the effect of tidal phenomenon on the salinity ion concentration in the water elevates, leading to the upraised conductivity. Our result is consistent with Corwin & Yemoto (2017). The results showed a high level of turbidity in all months in the present section of the river due to raising the ratio of suspended solid matters as consequently of anthropogenic activities which increased insoluble organic or inorganic materials (Butler & Ford 2018). The present results of pH values rates showed within the ranges of Iraqi inland waters (Ewaid & Abed 2017). The present results of total dissolved solids were high in the studied regions due to high levels of salinity and discharges of effluents from anthropogenic activities especially from January to March, which are in line with Corwin & Yemoto (2017). Fig. (2-A) illustrates the increased influence of salinity, electric conductivity and total dissolved solid respectively compared to the other factors, followed by water temperature and dissolved oxygen, whereas a slight effect appears due to turbidity in the study area. The figure showed an inverse correlation between water temperatures and the other factors (DO, salinity, electric conductivity, hydrogen ion, and total dissolved solids) except the water turbidity which was positively correlated with temperature. The results of environmental variables were subjected to Principal Component Analysis Technique (PCA; Figs. 3A and B), exhibiting that the eigenvalue of the studied factors was 6.39, which is a good value based on the seven factors measured, while the rates of variables were 56.38, 20.79 and 14.20% respectively. However, the impact of accumulation composition of the three components (F_1 , F_2 and F_3) was 56.38, 77.17 and 91.37% respectively. Fig. 4. revealed that January, February, March, April, June, July, and September formed 94.66% of the total impacts on the habitat of the study area. November exhibited the lowest variation in the variables (0.007%), while the highest values of impacts were 20.35% in January. Canonical correspondences analysis (CCA; Fig. 5) displayed that the fish species were divided into groups according to species tendencies. Fishes that prefer salinity, turbidity, temperature, electric conductivity, and total dissolved solids were the most influential variables in the study area contributing to the distribution and dispersal of fish species. Turbidity and water temperature also played an important role in the occurrence and distribution of fish species and were inversely correlated with salinity, DO, electric conductivity, and total dissolved solids. Fig. 5. revealed two fish groups according to their preferred first group coexistence associated with salinity represented by species inside the blue circle such as *Ilisha compressa*, *Hyporhamphus limbatus*, *Nuclequula gerreoides*, *Amblyaster sirum*, *Platycephalus indicus*, *Sardine albella*, *Pseudosynanceia melanostigma* and *Eupleurogrammus muticus*. The small group condensation turbidity (species inside a green circle) included *Aphanius dispar* and *Trichiurus lepturus*.




Fig. 3. The correlation among the physicochemical variables in the study area.

Another group including species preferred temperature represented by the species inside orange circle *C. zillii*, *Acanthopagrus bifasciatus* *T. ilish*, *Thryssa whiteheadi*, *Acanthopagrus arabicus*, *P. latipinna*, *N. nasus*, *Planiliza subviridis*, and *Brachirus orientalis*. These results were in line with Plavan *et al.* (2016) who studied the seasonal composition and abundance of the sub-estuarine fish assemblage on the coast of the Rio de la Plata estuary and with Mohamed & Abood (2017) who reported the compositional alteration in the fish assemblage structure in Shatt Al-Arab River, Southern Iraq.




Fig. 4. Principal Component Analysis Technique (PCA). The impacts of present physicochemical variables on study area among the months.

Relative abundance

Three species preside the numerical relative abundance of the study area, forming 80.96% of the total number of species. *Acetrogobius dayi* constituted 32.00%, *Boleophthalmus dussumieri* 28.92%, and *Aphanius dispar* 20.04% of the total caught. Gobiidae, the most abundant family in the investigated area, formed 60.92%, followed by Cyprinodontidae 20.04% and Engraulidae 8.70%, whereas among the eight orders, Perciformes was the most abundant order encompassing 65.26%, Cyprinodontiformes 20.94%, and Clupeiformes 10.47%. The three most abundant species were a non-commercial small size that was not subjected to fishing pressure. The present results were in accordance with Younis & Al-Shamary (2011) and Abdullah *et al.* (2018) from the point view of occurrence and abundance trends (Table 3).




Fig. 5. Canonical Correspondences Analysis (CCA) between the environmental variables and fish species.

Table 2. Fish species with their families and orders southern part of Main Outfall Drains.

Species	Family	Order	Habitat
<i>Amblyaster sirum</i>			M
<i>Nematalosa nasus</i>			M
<i>Sardine albella</i>	Clupeidae		M
<i>Tenelousa ilisha</i>		Clupeiformes	M
<i>Thryssa whiteheadi</i>	Engraulidae		M
<i>Ilisha compressa</i>	Pristigasteridae		M
<i>Carassius gibelio</i>			F
<i>Carasobarbus luteus</i>	Cyprinidae	Cypriniformes	F
<i>Hemiculter leucisculus</i>			F
<i>Aphanius dispar</i>	Cyprinodontidae		F
<i>Pocillia latipinna</i>	Pociliidae	Cyprinodontiformes	F
<i>Hyporhamphus limbatus</i>	Hemiramphidae	Beloniformes	M
<i>Planiliza abu</i>			F
<i>Planiliza subviridis</i>	Mugilidae	Mugiliformes	M
<i>Platycephalus indicus</i>	Platycephalidae		M
<i>Pseudosynanceia melanostigma</i>	Synanceiidae	Scorpaeniformes	M
<i>Acetrogobius dayi</i>			M
<i>Boleophthalmus dussumieri</i>	Gobiidae		M
<i>Coptodon zillii</i>			F
<i>Oreochromis aureus</i>	Cichlidae		F
<i>Eupleurogrammus muticus</i>			M
<i>Trichiurus lepturus</i>	Trichiuridae	Perciformes	M
<i>Silago sihama</i>	Sillaginidae		M
<i>Nuchequula gerreoides</i>	Leiognathidae		M
<i>Acanthopagrus arabicus</i>			M
<i>Acanthopagrus bifasciatus</i>	Sparidae		M
<i>Brachirus orientalis</i>	Soleidae	Pleuronectiformes	M

Ecological indices

The study revealed evident fluctuation in the values of ecological indices during the study period. The diversity index values ranged from 0.98 in March to 2.07 in April. The richness index varied from 0.96 in November to 1.94 in April. The evenness index values fluctuated from 0.38 in March to 0.89 in February (Fig. 6).




Fig. 6. Monthly variations in the ecological indices southern part of Main Outfall Drains, southern Iraq.

Table 3. The relative abundance of fish species with their families and orders southern part of Main Outfall Drains, southern Iraq.

Species	No.	%	Family	No	%	Order	No	%
<i>Amblyaster sirum</i>	12	0.10						
<i>Nematalosa nasus</i>	18	0.164	Clupeidae	192	1.75	Clupeiformes	1150	10.47
<i>Sardine albella</i>	16	0.146						
<i>Tenelousa ilisha</i>	146	1.33						
<i>Thryssa whiteheadi</i>	955	8.69	Engraulidae	955	8.70			
<i>Ilisha compressa</i>	3	0.02	Pristigasteridae	3	0.03			
<i>Carassius gibelio</i>	42	0.38	Cyprinidae	57	0.52	Cypriniformes	57	0.52
<i>Carasobarbus luteus</i>	11	0.1						
<i>Hemiculter leucisculus</i>	4	0.03						
<i>Aphanius dispar</i>	2200	20.04	Cyprinodontidae	2200	20.04	Cyprinodontiformes	2299	20.94
<i>Pocillia latipinna</i>	99	0.90	Poeciliidae	99	0.90			
<i>Hyporhamphus limbatus</i>	9	0.08	Hemiramphidae	9	0.082	Beloniformes	9	0.08
<i>Planiliza abu</i>	65	0.59	Mugilidae	232	2.11	Mugiliformes	232	2.11
<i>Planiliza subviridis</i>	167	1.52						
<i>Platycephalus indicus</i>	6	0.055	Platycephalidae	6	0.055	Scorpaeniformes	9	0.08
<i>Pseudosynanceia melanostigma</i>	3	0.027	Synanceiidae	3	0.027			
<i>Acetrogobius dayi</i>	3514	32.00	Gobiidae	6690	60.92			
<i>Boleophthalmus dussumieri</i>	3176	28.92						
<i>Coptodon zillii</i>	6	0.05	Cichlidae	32	0.29			
<i>Oreochromis aureus</i>	26	0.23						
<i>Eupleurogrammus muticus</i>	238	2.16	Trichiuridae	262	2.39	Perciformes	7166	65.26
<i>Trichiurus lepturus</i>	24	0.21						
<i>Silago sihama</i>	37	0.337	Sillaginidae	37	0.34			
<i>Nuquequula gerreoides</i>	9	0.08	Leiognathidae	9	0.08			
<i>Acanthopagrus arabicus</i>	134	1.22	Sparidae	136	1.24			
<i>Acanthopagrus bifasciatus</i>	2	0.01						
<i>Brachirus orientalis</i>	59	0.53	Soleidae	59	0.54	Pleuronectiformes	59	0.54

The present results revealed that the diversity index is shown as poor according to Jorgensen *et al.* (2005). The richness index tends to be troubled and the evenness index seems semi-balanced. The values of ecological indices were within the general trend of the previous studies (Yousif 1986; Jasim 2003; Younis & Al-Shamary 2011; Abdullah *et al.* 2018), with a slight increase in the range of some indices due to that studies implemented to the south adjacent to the northwest area of the Persian Gulf which allows entering a large number of marine species leading to the elevated values of the diversity index (Table 4).

Table 4. Comparison of ecological indices with the previous studies southern part of Main outfall drains, southern Iraq.

The study	The study region	Diversity index	Evenness index	Richness index
Yousif (1986)	Shatt Al-Basrah canal	0.49-2.84	0.07-0.50	1.33-5.32
Jasim (2003)	Shatt Al-Basrah Canal and Shatt Al-Arab River	0.17-2.13	0.24-0.98	0.28-2.87
Younis and Al-Shamary (2011)	Station 1	1.01-2.32	0.37-0.86	0.95-3.05
	Shatt Al-Basrah Canal			
Abdullah <i>et al.</i> 2018	Station 2	0.88 -2.24	0.37-0.79	1.45-3.06
	Main outfall drains (Al-Kii bridge area)	1.23-2.17	0.52-0.88	1.14-2.02
The present study	The southern part of Main Outfall Drains, Southern Iraq	0.98-2.07	0.38-0.89	0.96-1.94

CONCLUSION

The present study concluded that water quality represented by physicochemical variables plays an important role in the composition of the fish community structure that mostly constitutes the tolerant marine and freshwater fish species. Moreover, salinity contributes to a large degree in determining the nature of fish assemblages. In addition, the present section of the river represented an unstable habitat with poor water quality due to organic and chemical pollutions resulting from the process of reclamation of agricultural lands, and as a result of human management of freshwater quantities coming from Al-Siphone regulate in the Thi-Qar Province. The tidal processes contribute to the spread of organic pollutants which are thrown into the river from the Hamdan region in Southern Basrah City.

ACKNOWLEDGMENT

The author's grateful Department of Marine Vertebrate, Marine Science Centre to provide the facilities and opening the laboratories to perform the present work.

REFERENCES

- Al Daham, NK & Yousif, AY 1990, Composition, seasonality and abundance of fishes in Shatt Al-Basrah channel, an estuary southern Iraq. *Estuarine, Coastal and Shelf Science*, 30: 1-11.
- Al Munshed, HN 1998, Primary productivity in the Shatt Al-Basra canal after the completion of the Saddam River. MSc. Dissertation, College of Agriculture, University of Basrah, Iraq, 77 p.
- Abdullah, AHJ, Faris, RA & Abdullah, SA 2018, Structural diversity of fish assemblage in the southern sector of Main Outfall Drains northwest of Basrah, Iraq. *Basrah Journal of Agricultural Sciences*, 31: 1-11.
- Bower, LM & Winemiller, KO 2019, Fish assemblage convergence along stream environmental gradients: an intercontinental analysis. *Ecography*, 42: 1691-1702.
- Butler, BA & Ford, RG 2018, Evaluating relationships between total dissolved solids (TDS) and total suspended solids (TSS) in a mining-influenced watershed. *Mine Water and the Environment*, 37: 18-30.

- Camargo, TR, Ramos, P, Monserrat, JM, Prentice, C, Fernandes, CJ, Zambuzzi, WF & Valenti, WC 2021, Biological activities of the protein hydrolysate obtained from two fishes common in the fisheries bycatch. *Food Chemistry*, 342: 128361 p.
- Corwin, DL & Yemoto, K 2017, Salinity: Electrical conductivity and Total dissolved solids. *Soil Science Society of America Journal*, 2: 1-16. DOI:10.2136/amsa2015.0039.
- Craig, LS, Olden, JD, Arthington, AH, Entekin, S, Hawkins, CP, Kelly, JJ, Kennedy, TA, Maitland, BM, Rosi, EJ, Roy, AH & Strayer, DL 2017, Meeting the challenge of interacting threats in freshwater ecosystems: A call to scientists and managers. *Elementa: Science of the Anthropocene*, 5: 1-15.
- de Mello, K, Valente, RA, Randhir, TO & Vettorazzi, C.A 2018, Impacts of tropical forest cover on water quality in agricultural watersheds in southeastern Brazil. *Ecological Indicators*, 93: 1293-1301.
- Ewaid, SH & Abed, SA 2017, Water quality index for Al-Gharraf river, southern Iraq. *The Egyptian Journal of Aquatic Research*, 43: 117-122.
- Fricke, R, Eschmeyer, WN, Van der Laan, R 2021, Eschmeyer's catalogue of fishes: genera, species, references. Electronic version accessed.
- Froese, R & Pauly, D 2022, Fish Base. World Wide Web electronic publication. www.fishbase.org, version (02/2022).
- Forouhar Vajargah, M, Sattari, M, Imanpur, J & Bibak, M 2020a. Length-weight relationship and some growth parameters of *Rutilus kutum* (Kaminski 1901) in the South Caspian Sea. *Experimental Animal Biology*, 9:11-20.
- Forouhar Vajargah, M, Sattari, M, Imanpour Namin, J & Bibak, M 2020b. Length-weight, length-length relationships and condition factor of *Rutilus kutum* (Actinopterygii: Cyprinidae) from the southern Caspian Sea, Iran. *Journal of Animal Diversity*, 2:56-61.
- Herbert, ER, Boon, P, Burgin, AJ, Neubauer, SC, Franklin, RB, Ardón, M, Hopfensperger, KN, Lamers, LP & Gell, P 2015, A global perspective on wetland salinization: ecological consequences of a growing threat to freshwater wetlands. *Ecosphere*, 6: 1-43.
- Huang, J, Huang, L, Wu, Z, Mo, Y, Zou, Q, Wu, N & Chen, Z 2019, Correlation of fish assemblages with habitat and environmental variables in a headwater stream section of Lijiang River, China. *Sustainability*, 11: 1135.
- Jasim, AA 2003, Some biological aspect of fish juveniles in Basrah canal and Shatt Al-Arab River. PhD Dissertation, College of Agriculture, University of Basrah, 72 p.
- Jiang, X, Pan, B, Sun, Z, Cao, L & Lu, Y 2020, Application of taxonomic distinctness indices of fish assemblages for assessing effects of river-lake disconnection and eutrophication in floodplain lakes. *Ecological Indicators*, 110: 105955.
- Jørgensen, SE, Xu, FL, Salas, F & Marques, JC 2005, Application of indicators for the assessment of ecosystem health. Handbook of ecological indicators for assessment of ecosystem health, Taylor & Francis, 25 p.
- Mazareiy, MH, Sattari, M, Imanpour Namin, J, 2019, Length-weight relationship and some biological parameters of *Alosa braschnikowi* (Borodin 1904) from coast of the Guilan Province, *Journal of Aquaculture Sciences*, 7: 9-18 (In Persian).
- Mohamed, ARM & Abood, AN 2017, Compositional change in fish assemblage structure in the Shatt Al-Arab River, Iraq. *Asian Journal of Applied Sciences*, 5: 944-958.
- Ngor, PB, Grenouillet, G, Phem, S, So, N & Lek, S 2018, Spatial and temporal variation in fish community structure and diversity in the largest tropical flood-pulse system of South-East Asia. *Ecology of Freshwater Fish*, 27: 1087-1100.
- Nyitrai, D, Martinho, F, Dolbeth, M, Baptista, J & Pardal, M 2012, Trends in estuarine fish assemblages facing different environmental conditions: combining diversity with functional attributes. *Aquatic Ecology*, 46: 201-214. <https://doi.org/10.1007/s10452-012-9392-1>,
- Parker, J, Epifanio, J, Casper, A & Cao, FY 2015, The effects of improved water quality on fish assemblages in a heavily modified large river system. *River Research and Applications*, 32: 992-1007.
- Plavan, AA, Gurdek, R, Muñoz, N, Gutiérrez, JM, Spósito, M, Correa, P & Caride, A 2016, Seasonal composition, abundance and biomass of the sub-estuarine fish assemblage in Solís Chico (Río de la Plata estuary, Uruguay). *Brazilian Journal of Biology*, 77: 622-631.
- Poff, NL 2018, Beyond the natural flow regime? Broadening the hydro-ecological foundation to meet environmental flows challenges in a non-stationary world. *Freshwater Biology*, 63: 1011-1021

- Pyron, M, Mims, MC, Minder, MM, Shields, RC, Chodkowski, N & Artz, CC 2019, Long-term fish assemblages of the Ohio River: Altered trophic and life history strategies with hydrologic alterations and land use modifications. *PlosOne*, 14: e0211848.
- Rajwa Kuligiewicz, A, Bialik, RJ & Rowiński, PM 2015, Dissolved oxygen and water temperature dynamics in lowland rivers over various timescales. *Journal of Hydrology and Hydromechanics*, 63: 353-363.
- Santos, RVS, Ramos, S & Bonecker, ACT 2017, Environmental control on larval stages of fish subject to specific salinity range in tropical estuaries. *Regional Studies in Marine Science*, 13: 42-53.
- Sattari, M, Namin, JI, Bibak, M, Vajargah, MF, Hedayati, A, Khosravi, A & Mazareiy, MH 2019, Morphological comparison of western and eastern populations of Caspian kutum, *Rutilus kutum* (Kamensky, 1901) (Cyprinidae) in the southern Caspian Sea. *International Journal of Aquatic Biology*, 6: 242-247.
- Taylor, CM, Miyazono, S, Cheek, CA, Edwards, RJ & Patiño, R 2019, The spatial scale of homogenisation and differentiation in Chihuahuan Desert fish assemblages. *Freshwater Biology*, 64: 222-232.
- Walag, AMP & Canencia, MOP 2016, Physico-chemical parameters and macrobenthic invertebrates of the intertidal zone of Gusa, Cagayan de Oro City, Philippines. *Advance Environmental Science International Journal Bioflux Society*, 8: 71-82.
- Welch, PS 1964, *Limnology*. 2nd Ed, Mc Graw Hill Book Co., New York, 538 p.
- Xin, XK, Li, KF, Finlayson, B & Yin, W 2015, Evaluation, prediction and protection of water in Danjiangkou Reservoir, China. *Journal of Water Science and Engineering*, 8: 30-39, DOI:10.1016/J.WSE.2014.11.001
- Wen, Y, Schoups, G & Van De Giesen, N 2017, Organic pollution of rivers: Combined threats of urbanization, livestock farming and global climate change. *Scientific Reports*, 7: 1-9.
- Yang, B, Dou, M, Xia, R, Kuo, YM, Li, G & Shen, L 2020, Effects of hydrological alteration on fish population structure and habitat in river system: A case study in the mid-downstream of the Hanjiang River in China. *Global Ecology and Conservation*, 23: e01090.
- Younis & Al Shamary 2011, Species composition of fishes' assemblage in Shatt Al-Basrah canal, South of Iraq. *Iraqi Journal of Aquaculture*, 8: 137-156.
- Younis & Al Shamary, 2012, Impact of salt front on fishes' assemblage in Shatt Al-Basrah canal. *Basrah Journal of Agricultural Sciences*, 25: 89-103.
- Yousif, AY 1986, Species structure of fishes in the Shatt Al-Basrah canal and their feeding relationships. M. Sc. Thesis, College of Agriculture, University of Basrah, 102 p.
- Zeni, JO & Casatti, L 2014, The influence of habitat homogenization on the trophic structure of fish fauna in tropical streams. *Hydrobiologia*, 726: 259-270.