

Genetic polymorphism of β -Lg gene in Iraqi buffalo using Polymerase Chain Reaction – Restriction Fragment Length Polymorphism

Bashar F. Zaqeer*, Muntaha Y Yousief, Salah Mahdi Alsudany, Jaafar M. Owaid, Falih Hasan Hamad

College of Agriculture, University of Basrah, Basrah, Iraq

* Corresponding author's E-mail: bashar.zaqeer@uobasrah.edu.iq

ABSTRACT

This study was conducted on 50 buffaloes from Basrah Governorate, Iraq to study the genetic polymorphisms of the β -lactoglobulin gene in the local buffaloes. The lab examinations were conducted in the Genetic Engineering Laboratory, College of Agriculture, University of Basrah. DNA was extracted by kit (PureLink Genomic DNA kits, Invetrogen, USA). The results of this technique showed the appearance of the expected gene bundle of 252 base pairs from all Iraqi buffalo samples, as the primer interacted with all DNA isolates taken from the studied buffalo samples. Digestion technique with restriction enzymes (HaeIII) was used to detect the genetic polymorphisms of β -lactoglobulin gene, exhibiting two bands to all studied buffalo samples. Only the homozygous BB genotype has been shown.

Keywords: Genetic polymorphisms, β -lactoglobulin gene, PCR technique, Iraqi buffalo.

Article type: Research Article.

INTRODUCTION

One of the basic elements in determining the genetic forms of milk proteins is the use of electrophoresis (Cattaneo *et al.* 1996). Studies on the genetic polymorphisms of proteins have developed over time. Ng-Kwai-Hang (1998) summarized its most important goals by revealing the chemical properties of proteins, including milk and blood proteins, and some features of the similarity between these proteins, as well as clarifying the different relationships between different types of animals. Electrophoresis is used to detect genetic polymorphisms in milk proteins that result from the substitution of one amino acid for another or from the deletion of one or more amino acids. Gel electrophoresis, which displays several bands of various lengths, allows researchers to see even variation among genotypes (David & Deutch 1992). As it depends on the existence or lack of mutations, DNA analysis you could also examine the DNA. Point mutations and genetic DNA restructuring (Recombinant DNA - rDNA) are two of the most significant mutations that take place at the DNA level (Di Gregorio *et al.* 1991). In the case of point mutations, a single nucleotide is replaced, and in this case, differences between individuals are detected using restriction enzymes that cut DNA in specific regions, depending on the type of enzyme. As for the phenomenon of DNA organization, it includes the insertion or deletion of part of the genetic DNA fragments. The most common is the formation of sequential chains, and these can be individually amplified and observed using the Polymerase Chain Reaction (PCR) technique (Soller 1990). The phenomenon of genetic polymorphism in cattle has been well studied in recent years due to the close relationship between the genotypes of milk proteins and the important economic characteristics in cattle. Milk contains four main types of casein, namely, α S1-casein, α S2-casein, β -casein, κ -casein (Davies & Law 1987). The PCR-RFLP amplification is one of the first techniques used to identify genetic variation in DNA between two or more organisms (Bruns *et al.* 1991 & Strange 2003). It is also one of the most important techniques used in studies of the evolution of breeds with many organisms, as it has been used

extensively in taxonomy. In this technique, a gene conserved on the target region is amplified and then the amplification product is digested by a specialized digestive enzyme to obtain the digested segments expressing that species (Chen 1992; Appel & Gordon 1995). The present study aimed at the use of milk protein polymorphisms in association with milk yield and milk component of Iraqi buffalo.

MATERIALS AND METHODS

DNA extraction

DNA was extracted from 50 buffaloes from Basrah Governorate in the Laboratory of Genetic Engineering, College of Agriculture, University of Basrah, Iraq. DNA was extracted by kit (PureLink Genomic DNA kits, Invetrogen, USA).

Measurement of the amount of DNA by the Nano Drop device

The amount of DNA was measured by a Nano Drop device prepared by the American company Thermo Scientific (Fig. 1) to identify the size of the genome ($\text{ng } \mu\text{L}^{-1}$). The presence of DNA was detected by electrophoresis technique in the presence of agarose gel at a concentration of 1%.



Fig. 1. The Nano Drop device, which is equipped with the American Thermo Scientific Company.

The materials for polymerase chain reaction technology were prepared and placed in a bowl containing pieces of ice for protecting from heat. The work was done in a sterile and clean place in a special PCR cabin, which contained UV rays to sterilize the micropipettes, tubes and tabs, taking into account wearing sterile medical gloves when working. The PCR reaction mixture was prepared in a 100 μL Eppendorf tube, the final concentration of the components was 25 μL , then the tubes were placed in a Snip centrifuge for 30 seconds. Table 1 shows the amounts of materials used in the reaction.

Table 1. Amounts of materials used in the PCR-RFLP techniques (microliters).

Chemical material	Master mix	Primer	DNA template	Distill water	Final volume
Volume	12.5	1*	5	6.5	25

* In the PCR-RFLP technique, two primers (Forward and Reverse) are used, and the amount of distilled water is reduced to 5.5 μL .

PCR-RFLP technique

Iraqi buffalo DNA was amplified using a specific primer for the β -lactoglobulin gene 5-GTC CTT GTG CTG GAC ACC GAC TAC A-3 and (forward) 5-CAG GAC ACC GGC TCC CGG TAT ATG A-3. The tubes were then placed in a thermo cycler, and the device's program was set according to Di Xi Ren *et al.* (2011) using the program for the initiator and according to the following steps, which are mentioned in Table 2.

Table 2. Phases of PCR reactions time, cycle numbers and temperature.

Phases	Temperature ($^{\circ}\text{C}$)	Time (minutes)	No. of cycles
Primary denaturation	94	5.00	1
Denaturation	94	1.00	
Annealing	61	1.00	35
Elongation	72	1.00	
Final elongation	72	10.00	

Electrophoresis technique for amplification product

The electrophoresis technique was used on an agarose gel to determine the success of the DNA amplification process. The DNA Marker (100 base pairs) was used by mixing 1.5 μ L of the molecular weight marker with 3.5 μ L methyl orange. As for the rest of the pits, a PCR product of 5 μ L was placed in it, the PCR product was passed on an agarose gel with a concentration of 2%, i.e., 0.5 g agarose was dissolved in 25 mL 1X TBE solution, with the addition of 1 μ L ethidium bromide dye. Then, the electric current was fixed at 85 volts and 65 mA for 45 minutes. After completing migration, the gel was examined using a UV device to know the sizes of the bands in the gel.

Digestion by restriction enzyme

The digestion process was carried out by restriction enzymes, by cutting the gene of size (252 bp) with the help of (HaeIII) enzyme. The digestion mixture was prepared for the samples and mixed with a micropipette, then the enzyme was added to the mixture. So that, the final mixture became 20 μ L, and then the samples were placed in a centrifuge for 5 seconds. Table 3 shows the concentrations and quantities of the materials used in the reaction.

Table 3. Amounts of materials used in the digestion process by the restriction enzyme (μ L).

Id	Item	Quantity
1	DNA	7
2	Restriction Enzyme	1.5
3	10x Tango Buffer	2
4	BSA	0.5
5	Sterilized distill water	9

The samples were incubated in a water bath at a temperature of 37 °C for 3 hours. After the incubation period ended and the digestion process was completed, 3 μ L of the product was taken and electrophoresis was carried out on a 2% agarose gel with a DNA marker (50 bp) placed to check the digestion process. After completing the electrophoresis process, the gel was examined with a UV device and the DNA bands were identified.

RESULTS AND DISCUSSION

Since the primers used in the experiment interacted with all of the DNA samples taken from the blood of the studied buffaloes, the results of this technique demonstrated the appearance of the expected bands of the desired gene, amounting to 252 base pairs, in all of the buffalo samples under study (Fig. 2). This confirmed that all of the tested isolates are from the buffalo.




Fig. 2. Results of electrophoresis on a 2% agarose gel using the primer for the β -lactoglobulin gene, M. Molecular weight indicator (100 bp), (sample No. 1 and 5 are control samples and samples 2-4 and 6-10 represent the β -lactoglobulin gene).

PCR- RFLP technique

Digestion technique with categorical enzymes was used to detect the genetic polymorphisms of the gene β -lactoglobulin. Therefore, the amplification product was digested by the restriction enzyme HaeIII, which was

given in two packages to all studied buffalo samples (Fig. 3). This indicates that all studied samples belong to one genotype, which is BB. This result was consistent with Da Xi Ren *et al.* (2011), who indicated that the expected gene size of the Indian buffalo, using the same primers that was used in this study, as the gene size was 252 bp. If a genetic polymorphism of the β -lactoglobulin gene was found in the blood of local buffaloes, as electrophoresis examinations showed the presence of two clear bands indicating the presence of the genetic polymorphisms, however, all samples were similar genotype. The findings of the current study conflicted with those of Patel *et al.* (2007) and Kirimi *et al.* (2012), who discovered several alleles and three genotypes, AA, BB, and AB, respectively, with AB being the dominant genotype in a different investigation on the same gene but with a different primer (398 bp). Badola *et al.* (2003) and Soumi *et al.* (2006) obtained a single homozygous genotype. These results supported the present findings where there was only a single genotype using the same primer.




Fig. 3. Results of electrophoresis on a 2% agarose gel using HaeIII enzyme, M molecular weight marker (10 bp), sample No. 1 and 2 control, samples 3-10 homozygous genotype BB.

REFERENCES

- Appel, DJ & Gordon, TR 1995, Intraspecific variation within populations of *Fusarium oxysporum* based on RFLP analysis of the Intergenic spacer region of the rDNA. *Experimental Mycology*, 19: 120-128.
- Badola, S, Bhattacharya, TK, Biswas, TK, Shivakumar, BM, Kumar, P, Mishra, SS & Sharma, A 2003, Bubaline beta-lactoglobulin gene polymorphism: A contrast from its bovine counterpart. *Journal of Buffalo Science*, 19: 291-298.
- Bruns, TD, White, TJ & Taylor, JW 1991, Fungal Molecular Systematics. *Annual Review of Ecology and Systematics*, 22: 525-564.
- Cattaneo, TMP, Nigro, F & Greppi, GF 1996, Analysis of cow, goat and ewe milk mixtures by capillary electrophoresis: Primary approach. *Milchwissenschaft-milk Science International*, 51: 616-619, Corpus ID: 99517542.
- Chen, W 1992, Restriction fragment length polymorphism in enzymatically amplified ribosomal DNAs of three heterothallic *Pythium* species. *Journal of Phytopathology*, 82: 1467-1472.
- Chianese, L, Garro, G, Addeo, F, Lopez Galvez, G & Ramos, M 1993, Discovery of an ovine α -s2 casein variant. *Journal of Dairy Research*, 60: 485-493.
- David, VA & Deutch, AH 1992, Detection of bovine α -s1 casein genomic variants using the allele - specific polymerase chain reaction. *Animal Genetics*, 23: 425-429.
- Davies, DT & Law, AJ 1987, Quantitative fraction of casein mixtures by fast protein liquid chromatography. *Journal of Dairy Research*, 54: 369-376.
- Da Xi, R, Shu Ying, M, You liang, Ch, Cai Xia, Z, Xian Wei, I & Jian Xin, L 2011, Genotyping of the k-casein and β -lactoglobulin genes in Chinese Holstein, Jersey and water buffalo by PCR-RFLP. Indian Academy of Sciences, *Journal of Genet*, 19; 90(1): e1-5.
- Di Gregorio, P, Rando, A, Pieragostini, E & Masina, P 1991, DNA polymorphism at the casein loci in sheep. *Animal Genetics*, 22: 21-30.

- Jaayid, TA 2013, Rapid and sensitive identification of horse and donkey meat in Iraqi markets using SSR and PCR-RFLP based on mitochondrial DNA cytochrome b gene. *Journal of Agricultural Science and Technology*, 3: 11. USA.
- Jaayid, TA & Al Allak, ZS 2013, Successful diagnostic of fetal gender using polymerase chain reaction (PCR) of ovine amniotic fluid and embryos produced in vitro. *International Journal of Molecular Genetics*. 3: 29-37.
- Jaayid, TA & Dragh, MA 2013, Genetic biodiversity in buffalo population of Iraq using microsatellites marker. *Journal of Agricultural Science and Technology*, 4: 297-301.
- Jaayid, TA & Saker, DK 2012, Detection of species adulteration in imported meat from farm animals and pork using polymerase chain reaction with species specific repeat (PCR-SSR). *Basrah Journal of Agriculture*, 25: 151-183.
- Jaayid, TA & Saker, DK 2013, Using restriction fragment length polymorphism technique of cytochrome b gene for the identification of meat species and meat products in farm animals. 2nd International Food Safety Conference (IFSAC2013) Food Safety: Critical Dimension of Food Security in Emerging Economies. University of Putra, Malaysia.
- Jaayid, TA, Fahid, TA & Jasim, NY 2013, Oocytes extraction from Iraqi local goat ovaries by using aspiration and in vitro fertilization techniques. *Basrah Journal of Agriculture*, 26: 250-264.
- Jaayid, TA, Yousief, Zaqeer, BF & Owaid, JM 2013, Genetic polymorphism of kappa casein (κ -CN) in Iraqi buffalo using polymerase chain reaction-restriction fragment length polymorphism. *Journal of Agricultural Science and Technology*, 3: 10.
- Jaayid, TA, Al Waily, DS & Almubarak, ZHA 2013, Genetic diversity of *Fusarium oxysporum* f.sp *cucumerinum* isolates from cucumber: Differentiation by using random amplified polymorphic DNA. *Basrah Journal of Agricultural Sciences*, 26: 141-158.
- Jaayid, TA, Ayied, AY & Zaqeer, BF 2012, Study of β -lactoglobulin genotypes of arabi sheep milk and their relations with some genetic and production parameters. *Basrah Journal of Agricultural Sciences*, 25: 1-11.
- Jaayid, TA, Fahid, TA & Jasim, NY 2013, Oocytes extraction from Iraqi local goat ovaries by using aspiration and in vitro fertilization techniques. *Basrah Journal of Agricultural Sciences*, 26: 250-264.
- Karimi, K, Beigi Nasiri, MT, Fayyazi, J, Mirzadeh, KH & Roushanfekr, H 2009, Allele and genotype frequencies of β -lactoglobulin gene in Iranian Najdi cattle and buffalo populations using PCR-RFLP. *African Journal of Biotechnology*, 8: 3654-3657, <http://www.academicjournals.org/AJB>.
- Ng Kwai Hang, KF 1998, Genetic polymorphism of milk proteins: relationships with production traits, milk composition and technological properties. *Canadian Journal of Animal Science*, 78: 131-147.
- Nualchuen, W, Srisakwattana, K, Chethasing, S, Tasripoo, K, Usawang, S, Hengtrakulsin, R & Kamonpatana, K 2012, RFLP Analysis of beta-actoglobulin gene in swamp and murrah buffaloes using a single restriction enzyme. *Iranian Journal of Applied Animal Science*, 2: 301-303.
- Patel, AK & Tripathi, VN 1998, Effect of non-genetic factors on economic traits of Surti buffaloes. *The Indian Journal of Animal Sciences*, 68: 566-569.
- Soller, M 1990, Genetic mapping of the bovine genome using deoxyribonucleic acid - level markers to identify loci affecting quantitative traits of economic importance. *Journal of Dairy Science*, 73: 2628-2646.
- Soumi, S, Tarun, KB, Venkatachalaphthy, RT, Pushpendra, K & Sharma, A 2006, Cloning and characterization of α S₂-casein gene of Reverine buffalo DNA Seq. 17: 458-464. PMID: 17654008.
- Strange, RN 2003, Introduction to plant pathology. Great Britain by TJ International Ltd., Padstow, Cornwall, University College London., 497 p.

Bibliographic information of this paper for citing:

Zaqeer, B, F, Yousief, M, Y, Alsudany, S, M, Owaid, J, M, Hamad, F, H 2023, Genetic polymorphism of β -Lg gene in Iraqi buffalo using Polymerase Chain Reaction – Restriction Fragment Length Polymorphism. *Caspian Journal of Environmental Sciences*, 21: 343-347.
