

PAPER • OPEN ACCESS

Study of *Unio* shells in the Quaternary deposits, southern Iraq

To cite this article: Mohanad Hamid Al-Jaberi *et al* 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **1087** 012020

View the [article online](#) for updates and enhancements.

You may also like

- [Control Technology of Overlying Roadway in 7m Goaf](#)
Cong Liu, Zhennan Liu, Ji Li et al.
- [Biological Activity of Quinolinium Salts](#)
B M Gutsulyak
- [Species diversity of unionid mussels \(Mollusca: Bivalvia: Unionidae\) as a bioindicator of the state water environment of river basins of Ukraine](#)
L Shevchuk, L Vasileva, R Romaniuk et al.



The Electrochemical Society
Advancing solid state & electrochemical science & technology

243rd ECS Meeting with SOFC-XVIII

More than 50 symposia are available!

Present your research and accelerate science

Boston, MA • May 28 – June 2, 2023

[Learn more and submit!](#)

Study of *Unio* shells in the Quaternary deposits, southern Iraq

Mohanad Hamid Al-Jaberi^{1,2,4}, Harith A. Al-Saad¹, Ghazi A. Hussain² and Hiba Khalid Lafta³

¹ University of Basrah- College of Science- Geology department;

² A I Karkh University of Science – College of Energy and Environmental Science;

³ University of Basrah- College of Science- Geology department.

⁴ Corresponding author's e-mail: jaberi76@yahoo.com

Abstract. Shells are useful in evolutionary biology and paleobiology. Freshwater shells compose the main types of mollusks, and can use as a key for ecological variations. Shell samples in the present study were collected on the highway of Nasyriah city- Samawa city, southern Iraq. The study area contains a large accumulation of shells at 6-7m depths, especially at 7m depth. The main types of these shells are *Unio*, *Corbicula*, and *Melanoides tuberculata*. *Unio* was the most species common in the study area. *Unio* species were used to detect the paleoenvironment in Quaternary deposits, in southern Iraq. Calcite and aragonite are the common minerals in the *Unio* shells. According to the distribution and percentage of calcite and aragonite in the *Unio* shells, it was found the calcite at 6m is more than 7m, whereas the aragonite at 7m is highest compared to 6m. Under X-Ray Fluorescence analyzers, found that calcium oxide, silica, magnesium oxide, and phosphor oxide are the major oxides in the *Unio* shells, while Zn, Sr, Pb, and Rb are the main trace elements. Strontium (Sr) has a high concentration in the *Unio* shells at 7m than 6m, while magnesium (Mg) is found in a high concentration at 6m than 7m. According to trace and some of the major elements concentrations, the paleotemperature, paleosalinity, and water contamination of rivers at 6m are more than 7m, and the river water at 7m depth was fresh water with a high abundance of nutrients and climate was suitable to growth mollusks shells like *Unio* shells.

1. Introduction

The bivalve shells are used as environmental indicators [1]. Trace elements are incorporated into the bivalve shells according to the concentration of these elements in the water [2]. This is also affected by water salinity and temperature [3]. The observation of heavy metals in the aquatic environment involves detecting the metal in biota, sediments, and water [4,5,6,7]. The tolerance of mollusks' bivalves has made them an ecosystem monitoring [8]. The level of trace metal contamination of aqueous organisms is a result of a balance between metal loss and metal uptake [9]. The palaeoenvironmental recorders can be detected by the use of mollusks. The three-layered shell of *Unio* species of an outer thick periostracum, a prismatic layer, and a thick nacreous layer, as well as the aragonite mineralogy [10, 11, 12]. A number of authors have studied the mineralogy of *Unio* shells, such as the study of Gregoire (1961) [13] have revealed that *Unio* shells had a few crystals collection of aragonite. Taylor et al. (1969) [14] found two arrangements of the nacreous layers in several species of *Unionidae*. A study by Dauphin et al. (2017) [15] concluded the mineralogy and composition of a nacropismatic bivalve species: *Unio pictorum* and explained that the prismatic layer of *Unio* is aragonite, and the inner structure of the prismatic units strongly differs from those of the calcitic layers. Lyubas et al. (2019) [16] used *Unio* freshwater shells



as indicators of paleoclimate changes compare to the present in the Pleistocene and Pliocene deposits of ancient rivers in Russia. The main objective of this study was the reconstruction of some parameters of the aquatic environment in the Quaternary period, in southern Iraq.

2. Location of the study area

The study area is located in the north of Nasyriah city. The study area consists of Quaternary sediments in southern Iraq (Figure 1).




Figure 1. Map of the study area.

3. Materials and methods

Subfossil shells are the materials for this study, these shells are freshwater bivalve mollusks, belonging to the genera *Unio* [16]. The depths of 6 and 7m for two sites (S1 and S2) of outcrop Quaternary deposits (Figure 2,3,4) are chosen to study the *Unio* shells in the highway of Nasyriah - Samawa cities. The deposits in the study area are terrigenous sediments composed of sand, silt, and clay (Figure 4). The *Unio* samples were put in an acid-washed polyethylene bag. Grinding the shells by a glass mortar. X-ray fluoresces (XRF) and X-ray diffraction (XRD) are applied in this study to reveal of minerals and chemical composition of *Unio* shells.



Figure 2. *Unio* shells Shells at 6m.



Figure 3. *Unio* shells Shells at 7m.

4. Results

4.1. Mineralogy study

Calcite and aragonite are the common minerals in the *Unio* shells. According to the distribution and percentage of calcite and aragonite in the *Unio* shells, found the calcite at 6m is more than 7m, whereas the aragonite at 7m is highest compared to 6m.

4.2. Geochemical study

According to X-Ray Fluorescence (XRF) analyzers for *Unio* shells, it was estimated that calcium oxide, silica, magnesium oxide, and phosphor oxide are the major oxides, while Zn, Sr, Pb, and Rb are the main trace elements (Table 1).




Figure 4. The sequence layers from the top to the bottom in the study area.

Table 1. The XRF analysis results of *Unio* shells for the studied sites.

Elements	S1		S2	
	6m	7m	6m	7m
SiO₂ %	0.2188	0.0016	0.2672	0.0022
MgO %	0.2199	0.1394	0.2596	0.1282
P₂O₅ %	0.2119	0.2065	0.2155	0.2172
CaO %	55.49	55.95	55.6	55.86
MnO %	0.05792	0.02919	0.05438	0.02715
Zn ppm	12.131	11.087	12.86	10.581
Rb ppm	0.641	0.366	0.683	0.384
Ba ppm	0.00858	0.00831	0.00849	0.00824
Pb ppm	1.946	1.265	1.862	1.386
Sr ppm	1152	1380	1173	1376

5. Discussions

Mollusk is the second most phylum on the earth [17]. The shell shapes reflect the environment and their phylogenetic history [18, 19, 20, 21]. The chemical composition of bivalves shells can be used as monitoring contamination and detect the different environmental conditions [22,23,24,25,4,5]. The adaptability of the mollusks has made them a preferred organism as an indicator of the ecosystems [7,8,9]. XRF analysis of hard body parts of bivalves can give a piece of information for biomineralization and the interaction between the surrounding environment and organisms. XRD

analysis revealed two main layers in the *Unio*, the outer layer constitutes calcite and the inner layer constitutes aragonite. The high abundance of *Unio* shells at 6 and 7m depths gave an indication of the freshwater environment. The high accumulation of shells at 7m gave an indication of suitable conditions in that period of Quaternary Formation to grow and to provide the appropriate nutrients for reproduction. [25,26] The rate of *Unio* shells in the river was present in a high accumulation in the nutrient silty clay channels. Major and trace elements in the bulk *Unio* shells are displayed and summarized in Table 1, which emphasized that calcite is higher in the *Unio* shells at 6m depth compared with 7m, which may probably be due to the increase of calcium ions in the river water, these ions uptake by *Unio* shells to build their skeleton. The aragonite layer in the *Unio* shells at 7m is highest than 6m. The elements contained in the mollusk shells are suitable to concentrate in the calcite or aragonite according to their ionic radii [27]. Strontium (Sr) has a high concentration in the *Unio* shells at 7m than 6m, while magnesium (Mg) is found in a high concentration at 6m than 7m. [28,29,30] Indicated that the Sr element is more suitable to concentrate in the aragonite than the calcite layer. [31,32] Mentioned that Sr^{+2} substitutes more readily for Ca^{+2} in the aragonite than calcite. [27] The magnesium showed more preferential concentration in calcite rather than aragonite in shell growth. The Mg^{+2} content in the outer calcite layer is increased with increased environmental temperature, but not so regular as Sr concentration. The Mg content in the skeleton may be used for paleotemperature reconstruction, decrease the calcium elements and increase magnesium giving an indicator of high paleotemperature at 6m depth compare with 7m depth. An increase in the calcite layer with an increase in the Zn, Pb, Si, Mn, and Rb presented in large concentrations in *Unio* shells at 6m than 7m. [29] Mentioned the Zn, Pb, and Si which have small ionic radii and are present in a higher concentration in the calcite than aragonite. [29] Showed the concentration of Pb, Zn, and Mn, in the mollusks shells are higher when increasing the water salinity. In addition, the high accumulation of Mn, Zn, Pb, and Ba may have been attributed to sediments and water contamination at 6m than 7m, all of these showed the river at that time in 7m depth was fresh water with a high abundance of nutrients and climate was suitable to growth mollusks shells like *Unio* shells.

6. Conclusions

- (1) The high abundance of *Unio* shells at 6m and 7m depths gave an indication of a freshwater environment, and the great number of shells at these depths gave a clear evidence to appropriate conditions for growth and generation.
- (2) The aragonite layer in the *Unio* shells at 7m depth is more than 6m, while the calcite presence was reversed.
- (3) Strontium has a high concentration in the *Unio* shells at 7m than 6m, while magnesium is found in a high concentration at 6m than 7m.
- (4) Strontium was found in the aragonite layer, while magnesium is concentrated in the calcite layer of *Unio* shells.

References

- [1] Al-Dabbas, M., and Al-Jaberi, M. 2015 Elemental distribution in the calcite and aragonite layers of *Chione californiensis* from the Iraqi Coasts—North Arabian Gulf. *Arabian Journal of Geosciences* **8(11)** 9433-9440
- [2] Censi, P., Spoto, S. E., Saiano, F., Sprovieri, M., Mazzola, S., Nardone, G., Di Geronimo, S. I., Punturo, R., Ottonello, D. 2006 Heavy metals in coastal water systems: A case study from the northwestern gulf of Thailand. *Chem* **64** 1167–1176
- [3] Al-Dabbas, M., Hubbard, F.H, and McManus, J. 1983 The proportion of aragonite in the shells of *Mytilus edulis* from the Tay estuary *JAWRR* **2(2)** 73–79
- [4] Szefer, P., Frelek, K., Szefer, K., Lee, C.B., Kim, B.S., and Warzocha, J., Zdrojewska I., Ciesielski T. 2002 Distribution and relationships of trace metals in soft tissue, byssus and shells of *Mytilus edulis trossulus* from the southern Baltic *Environ Pollut* **120** 423–444
- [5] Liu, J.H., and Kueh, C. 2005 Biomonitoring of heavy metals and trace organics using the intertidal

- mussel *Perna viridis* in Hong Kong coastal waters. *Mar Pollut Bull* **51** 857–875
- [6] Hamed, A.H. and Emara, A.M. 2006 Marine molluscs as biomonitors for heavy metal levels in the Gulf of Suez *Red sea*. *J Mar Syst* **60** 220–234
- [7] Darvish, A. 2007 The survey of heavy metals (Cd, Cu, Ni and Hg) in water and soft tissue of *Saccostrea cucullata* in intertidal zone of Hormoz islans. Persian Gulf, MS thesis *Environmental Engineering* Azad University of Ahvaz, Iran. 140
- [8] Conti, M., and Cecchetti, G. 2003 A biomonitoring study: trace metals in algae and molluscs from Tyrrhenian coastal areas *Environ Res* **93** 99–112
- [9] Goodfriend, G., Magaritz, M., and Gat, J.R. 1989 Stable isotope composition of land snail body water and its relation to environmental waters and shell carbonate *Geochimica et Cosmochimica Acta* **53** 3215–3221
- [10] Dauphin, Y., Cuif, J.P., Cotte, M. and Salomé, M. 2012 Structure and composition of the boundary zone between aragonitic crossed lamellar and calcitic prism layers in the shell of *Concholepas concholepas* (Mollusca, Gastropoda) *Invertebr. Biol* **131** 165–176
- [11] Dauphin, Y., Cuif, J.P., Castillo-Michel, H., Chevillard, C., Farre, B. and Meibom, A. 2013 Unusual micrometric calcite–aragonite interface in the abalone shell *Haliotis* (Mollusca, Gastropoda). *Microsc. Microanal* **20** 276–284
- [12] Schmidt, W.J. 1924 Die Bausteine des Tierkörpers in polarisiertem. *Lichte. F. Cohen, Bonn* 528
- [13] Grégoire, C. 1961 Sur la structure submicroscopique de la conchioline associée aux prismes des coquilles de mollusques *Bull. K. Belg. Inst. R. Sci. Nat. Wet* **37** 1–34
- [14] Taylor, J.D., Kennedy, W.J. and Hall, A. 1969 The shell structure and mineralogy of the Bivalvia. Introduction. Nuculacae – Trigonacae. *Bull. Brit. Mus. Nat. Hist. Zool* **3** 1–125
- [15] Dauphin, Y., Luquet, G., Salome, M., BELLOT - GURLET, L., and Cuif, J. P. 2017 Structure and composition of *Unio pictorum* shell: arguments for the diversity of the nacropismatic arrangement in molluscs *Journal of microscopy* **270(2)** 156-169
- [16] Lyubas, A. A., Kabakov, M. B., Kriauciunas, V. V., Obada, T. F., Nicoara, I. N., and Tomilova, A. A. 2019 Freshwater mollusks from Neogene-Quaternary Dniester and Prut riverine deposits as indicator paleoenvironments: chemical composition of shells and its palaeoecological interpretation *Arctic Environmental Research* **19(1)** 35-42
- [17] Bank, R. A., Bieler, R., Bouchet, P., Decock, W., Dekeyser, S., Gofas, S., Kroh, A., Marshall, B., Neubauer, T.A., Neubert, E.G., Rosenberg, G., Sartori, A.F., Schneider, S., Trias-Verbeeck, A., Vandepitte, L.A., and Vanhoorne, B. 2014 Mollusca Base – announcing a World Register of all Molluscs. In White T. S. (ed.) Book of Abstracts EUROMAL 2014 The 7th Congress of the European Malacological Societies, 7th - 11th September 2014 *The Malacological Society of London, Cambridge*
- [18] Bogan, A. E. 2008 Global diversity of freshwater mussels (Mollusca, Bivalvia) in freshwater *Hydrobiologia* **595** 139–147
- [19] Sparks, B. W. 1961 The ecological interpretation of Quaternary non-marine Mollusca. *Proceedings of the Linnean Society of London* **172** 71–80
- [20] Graf, D. L. and Cummings, K.S. 2007 Review of the systematics and global diversity of freshwater mussel species (Bivalvia: Unionoida) *Journal of Molluscan Studies* **73** 291–314
- [21] Aldridge, D. C., Fayle, T. M. and Jackson, N. 2007 Freshwater mussel abundance predicts biodiversity in UK lowland rivers *Aquatic Conservation: Marine and Freshwater Ecosystems* **17** 554–564
- [22] Lydeard, C., Cowie, R.H., Ponder, W.F., Bogan, A.E., Bouchet, P., Clark, S.A., Cummings, K.S., Frest, T.E., Gargominy, O., Herbert, D.J., Hershler, R., Perez, K.E., Roth, B., Seddon, M., Strong E.E.F.G., Thompson, F.G. 2004 *The global decline of nonmarine mollusks*. *Bioscience* **54** 321–330
- [23] Dalbeck, P.C. 2008 Crystallography, stable isotope and trace element analysis of *Mytilus edulis* shells in the context of ontogeny *Unpubl. MSc thesis, University of Glasgow* 235
- [24] Hubbard, F., McManus, J., and Al-Dabbas, M. 1981 Environmental influences on the shell

- mineralogy of *Mytilus edulis* *Geo-Marine Letters* **1(3-4)** 267
- [25] Boening, W. 1999 An evaluation of biomonitors of heavy metals pollution in marine waters *Environ Monit Asses* **55** 459–470
- [26] Zając, K., Zając, T., and Ćmiel, A. 2018 What can we infer from the shell dimensions of the thick-shelled river mussel *Unio crassus*? *Hydrobiologia* **810(1)** 415-431
- [27] Al-Jaberi, M. H. 2015 Study of the clastic and shells in selected area at NW of Arabian Gulf-South Iraq *Baghdad university, published thesis* 249P
- [28] Al-Dabbas, M. A. 1980 An examination of shell fragment distribution and geochemical features of *Mytilus Edulis* in the Tay estuary *PhD thesis, University of Dundee* 173
- [29] Broman, D., Lindqvist, L., and Lundbergh, I. 1991 Cadmium and zinc in *Mytilus edulis* L. from the Bothnian Sea and the northern Baltic proper *Environmental Pollution* **74(3)** 227-244.
- [30] Deer, W.A., Howie, R.A., and Zussman, J. 1992 An Introduction to the Rock-Forming Minerals, London (Longman Scientific & Technical), *Mineralogical Magazine* **56(385)** 617-619
- [31] Yan, H., Chen, J., and Xiao, J. 2014 A review on bivalve shell, a tool for reconstruction of paleoclimate and paleo-environment *Chinese Journal of Geochemistry* **33(3)** 310–315 <https://doi.org/10.1007/s11631-014-0692-0>
- [32] McMillan, E.A., Fairchild, I., Frisia, S., Borsato, A., and McDermott, F. 2005 Annual trace element cycles in calcite–aragonite speleothems: evidence of drought in the western Mediterranean 1200–1100 yr BP *J Quat Sci*, **20(5)** 423–433