

PAPER • OPEN ACCESS

## Survey of the Butterflies (Lepidoptera, Rhopalocera) Associated with some Brassicaceae Plants in Basrah Provence, Iraq

To cite this article: Faisal Nasser Jaber *et al* 2023 *IOP Conf. Ser.: Earth Environ. Sci.* **1262** 032003

View the [article online](#) for updates and enhancements.



245th ECS Meeting • May 26-30, 2024 • San Francisco, CA

Submit now!

Don't miss your chance to present!

Connect with the leading electrochemical  
and solid-state science network!

Deadline Extended: December 15, 2023



# Survey of the Butterflies (Lepidoptera, Rhopalocera) Associated with some Brassicaceae Plants in Basrah Provence, Iraq

Faisal Nasser Jaber<sup>1</sup>, Aqeel Alyousuf<sup>2</sup> and Hanaa H. Al-Saffar<sup>3</sup>

<sup>1,2</sup> Department of Plant Protection, Faculty of Agricultural, University of Basra, Basra, Iraq.

<sup>3</sup>Iraq Natural History Research Center and Museum, University of Baghdad, Baghdad, Iraq.

<sup>1</sup>E-mail: faisal.nasser@uobasrah.edu.iq

**Abstract.** Butterflies were collected from two regions of Basra south of Iraq during November 2021 to March 2022. This survey showed ten species belong to ten genera under four families: Nymphalidae, Pieridae, Lycaenidae and Hesperiidae. The species *Vanessa cardui* showed the highest level of distribution while the lowest one was *Pieria rapae*.

**Keywords.** Basra, Brassicaceae, Butterflies, Lepidoptera, Rhopalocera.

## 1. Introduction

The order of Lepidoptera is one of the important economic insects that spreads in various regions of the world, which includes about 180,000 species, 126 families and 46 subfamily's, 10% of the total described species of living organisms [1]. Butterflies are considered adequate ecological representative for the invertebrates [2-5], and clear evidence of changes in the environmental quality. Butterflies also play important ecological roles in natural and agricultural habitats; they perform basic ecosystem services [6,7].

The Brassicaceae or Cruciferae is known as the Mustard family, it consists of 338 genera and about 3709 species throughout the world[8]. The family is an important of human health [9] and source of many food plants such as, (Cabbage, *Brassica rapa* var.*capitata* L, white radish, *Raphanus sativus* L. and others, as well as some species of ornamental plants such as *Shabuy repandum* L., silver rose, *Lobularia maritima* Des [10]. In Iraq the family represented by 80 genera and 177 species [11].

This family includes a large number of vegetable crops including *Brassica oleracea* Var *capitata* and *Brassica oleracea* var. *italica*, *Brassica oleracea* Var. *botrytis*, radish and watercress ,cabbage crop is one of the most important crops of the cruciferous family, which is cultivated in large areas in Iraq and throughout the year almost. This is due to its nutritional value [12].

Host plant play one of critical point which are as shelter and larval host plants. Diversity Lepidoptera was influenced by environmental conditions. Environmental factors that affect the altitude, temperature, humidity, intensity of light, weather and seasons. Thus the Lepidoptera are as bio-indicators that one can be used to measure the effect of climate change on biodiversity.



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Cruciferous plants are angiosperms that require obligatory cross-pollination or facultative cross-pollination [14-16], and that pollinators have an important role in food production and conservation. On plant biodiversity [17,18]. more than 78% of angiosperm species depend on pollinators. Compulsory and optional pollination makes insect pollination necessary, or at least a positive factor in improving The process of fertilization in flowers [19]. Cruciferous plants attract a variety of pollinators, including honey bees such as *Apis mellifera* L. (Hymenoptera: Apidae), and solitary bees, such as *Andrena* spp. (Hymenoptera: Andrenidae), and syrphid flies, such as *Eristalis tenax* L. (Diptera: Syrphidae) [20].

## 2. Materials and Methods

The current survey was conducted at two regions of Basra (Karmat Ali and Al-Zubair ). The total garden area was (12x 60) m<sup>2</sup>. The areas were made up of vegetation , include seven species of the family Brasicaceae crops (Arugula *Eurica sativa*, White radish *Raphanus sativus* subsp. *acanthiformis*, cabbage *Brassica oleracea*, variety *capitata*, rapeseed *Brassica napus*, mustard *Brassica carinata* , cauliflower *Brassica oleracea*, variety *botrytis* and broccoli *Brassica oleracea*, variety *italic*), during September 2021. The plants were identified at the botanical herbarium of the College of Agriculture / University of Basrah. The specimens of butterflies were collected weekly during the flowering times of brasicus crops (November2021 to March 2022) by the aid of air and sweeping nets. Butterflies were killed by freezing about 10-15 minutes. Then the specimens were mounted by insect pins and storage in insect boxes.

The identification of butterflies (Families, genera and species) by using several taxonomic keys such as: [21-29]. The synonyms of species were according to (GBIF) 2022.

## 3. Results and Discussions

The survey showed ten species belonging to ten genera under four families. The specimens were collected from crusiferae plants at Basra provences . from two regions during November 2021 to March 2022. The number of material examined of species showed the specimens which seen and few killed for identification.

### 3.1. Family, Nymphalidae

- Common name: Brush footed Butterflies
- Synonyms: Acraiidiae
- Amathusiidae
- Brassolidae
- Danaidae
- Heliconidae
- Heliconiidae
- Ithomiidae
- Libytheidae
- Morphidae

#### 3.1.1. Genus: *Danaus* Kluk. 1802

- Common name: Tiger Milkweeds Butterflies
- Synonyms: *Danaida* Latreille, 1804
- *Danais* Latreille, 1807
- *Danaomorpha* Kremky, 1925
- *Danaus* Latreille, 1809
- *Danaus* Linnaeus, 1758
- *Danis* Lucas, 1889
- *Danus* Mitchell, 1872
- *Diogas* d'Almeida, 1938
- *Festivus* Crotch, 1872

- *Nasuma* Moore, 1883
- *Panlymnas* Bryk, 1937
- *Salatura* Moore, 1880
- *Tasitia* Moore, 1883

### 3.1.2. *Danaus Chrysippus* (Linnaeus, 1758)

- Common Name: Plain Tiger
- Synonyms: *Danais chrysippus* (Linnaeus, 1758)
- *Danais chrysippus limbata* Matsumura, 1929
- *D. chrysippus subpurpurea* Matsumura, 1929
- *D. chrysippus vigelii* Heylaerts, 1884
- *D. cratippus* Felder, 1860
- *D. fuscippus* van Eecke, 1915
- *D. lemeemagdalena* Lemée, 1950
- *Danaus aegyptius* Schreber, 1759
- *D. albinus* Lanz, 1896
- *D. alcippoides* Moore, 1883
- *D. alcippus* Cramer, 1777
- *D. anomala* Dufrane, 1948
- *D. asclepiadis* Gagliardi, 1811
- *D. auriflave* Eecke, 1914
- *D. axantha* Hayward, 1922
- *D. bataviana* Moore, 1883
- *D. bipunctata* Dufrane, 1948
- *D. bowringi* Moore, 1883
- *D. candidata* Hayward, 1922
- *D. chrysipellus* Strand, 1910
- *D. chrysippus aegyptius* (Linnaeus, 1758)
- *D. clarippus* Weymer, 1884
- *D. completa* Dufrane, 1948
- *D. cratippus* Felder, 1862
- *D. deficiens* Dufrane, 1948
- *D. dorippus* Klug, 1845
- *D. duplicata* Dufrane, 1948
- *D. Dupont* Dufrane, 1948
- *D. fuscippus* Eecke, 1915
- *D. gelderi* Snellen, 1891
- *D. hanoiensis* Dufrane, 1948
- *D. impunctata* Dufrane, 1948
- *D. canariensis* Fruhstorfer, 1898
- *D. klugii* Butler, 1886
- *D. lemeemagdalena* Lemeé, 1950
- *D. liboria* Hulstraet, 1931
- *D. limbata* Matsumura, 1929
- *D. luxurians* Dufrane, 1948
- *D. margarita* Ruber, 1926
- *D. orientis* Aurivillus, 1909
- *D. ornata* Dufrane, 1948
- *D. (Anosia) petilea* Linnaeus, 1758
- *D. petilia* Stoll, 1790

- *D. praealbata* Frorich, 1928
- *D. pseudopetilea* Kalis, 1933
- *D. radiata* Dufrane, 1948
- *D. reducta* Dufrane, 1948
- *D. rubra* Eecke, 1915
- *D. semialbinus* Strand, 1910
- *D. subpurpurea* Matsumura, 1929
- *D. subreducta* Dufrane, 1948
- *D. vigelii* Heylaerts, 1884
- *D. yuchingkini* Murayama & Shimonoya, 1960
- *Limnas klugii* Butler, 1886
- *Papilio asclepiadis* Gagliardi, 1811
- *P. chrysippus* Linnaeus, 1758
- *P. petilia* Stoll, 1790
- *Salatura chrysippus canariensis* Fruhstorfer, 1899

Material examined: 347 specimens: 223(♀♀+♂♂) Karmat Ali;(35. Nov. 2021; 33,Des.2021 and 50. Jan. 2022; 45. Feb. 2022; 60. March. 2022). 124(♀♀+♂♂) Al- Zubair: ;(24. Nov. 2021; 25,Des.2021 and 20. Jan. 2022; 25. Feb. 2022; 30. March. 2022).

Distribution: Iraq [30]; New Zealand [31]; Egypt[32]. Northern Africa Morocco, Algeria and Tunisia, Spain, France, Sardinia, Sicily, Italy, Albania, and Greece[33]; Jordon [34] ; Europe [35];Turkey[36]; Pakistan[37] and Azerbaijan [38].

### 3.1.3. Genus: *Vanessa Fabricius, 1807*

- Synonyms: *Amiralis* Rennie, 1832
- *Cynthia* Fabricius, 1807
- *Fieldia* Nicolescu, 1979
- *Kaniska* Moore, 189
- *Neofieldia* Özdkmen, 2008
- *Neopyrameis* Scudder, 1889
- *Papilio* Hubner, 1821
- *Phanessa* Sodovsky, 1837
- *Pyramides* Hübner, 1826
- *Vanessa* Latrelle, 1804
- *Vanessa* Sodoffsky, 1837

### 3.1.4. *Vanessa Cardui* (Linnaeus, 1758)

- Common Name: Painted Lady
- Synonyms: *Cynthia cardui* (Linnaeus, 1758)
- *C. eimi* Rambur, 1829
- *C. litoralis* de Souza, 1926
- *Papilio belladonna* Linnaeus, 1758
- *P. carduelis* Cramer, 1775
- *P. carduelis* Seba, 1765
- *P. cardui* Linnaeus, 1758
- *Parameis cardui* (Linnaeus, 1758)
- *Pryameis martha-maria* Stephan, 1924
- *Pyrameis brunnea-albimaculata* Reuss, 1915
- *Py. carduelina* Alphéraky, 1908
- *Py. carduelis* Cramer, 1775
- *Py. carduelis* Schultz, 1906

- *Py. cardui* (Linnaeus, 1758)
- *Py. elymnias* Rambur, 1829
- *Py. flava* Bandermann, 1928
- *Py. japonica* Stichel, 1908
- *Py. John Fischer de Waldheim*, 1932
- *Py. leachiana* Doubleday, 1849
- *Py. minor* Cannaviello, 1900
- *Py. pallens* Noel, 1881
- *Py. pallida* Schøyen, 1881
- *Py. rogeri* Meilhan, 1929
- *Py. rosacea* Reuss, 1915
- *Py. rosea* Pionneau, 1926
- *Py. schoenfellneri* Hoffmann, 1925
- *Py. subfracta* Stach, 1925
- *Py. takesakiana* Katô, 1925
- *Py. varini* Meilhan, 1929
- *Vanessa albicans* Verity, 1950
- *V. albipuncta* Lempke, 1956
- *V. belladonna* Godart, 1821
- *V. carduelis* Seba, 1765
- *V. elimi* Rambur, 1829

Material examined: 543 specimens: 403(♀♀+♂♂) Karmat Ali; (50. Nov. 2021; 55, Des. 2021 and 48. Jan. 2022; 100. Feb. 2022; 150. March. 2022). 140(♀♀+♂♂) Al- Zubair: ; (20. Nov. 2021; 25, Des. 2021 and 15. Jan. 2022; 30. Feb. 2022; 50. March. 2022).

Distribution: Iraq[40]; North, Central, and South America[41]; Jordon [42,43]; Turkey [44]; Syria and Armenia [45]. Pakistan[28] Azerbaijan [38] and Kuwait [46].

### 3.1.5. Genus: *Junonia Hubner, 1819*

- Common : Buckeyes & Pansies
- Synonym: *Alcyoneis* Hübner
- *Aresta* Billberg, 1820
- *Aresta* Dalman, 1820
- *Coryphaeola* Butler, 1878
- *Dunonia* Mabille, 1876
- *Callimula* Holland, 1920
- *Precis* Hubner, 1819

### 3.1.6. *Junona Orithya Linnaeus, 1758*

- **Common name:** Blue Pansy
- Synonyms: *Junonia albocincta* Butler, 1875
- *J. orithya* Linnaeus, 1758
- *J. albocincta* Linnaeus, 1758
- *J. jacouleti* Watari, 1941
- *J. leechi* Alphéraky, 1897
- *J. orithyra* Linnaeus, 1758
- *J. orthya leechi* Alphéraky, 1897
- *Papilio orithya* Linnaeus, 1764
- *Precis flava* Wichgraf, 1918
- *Precis Jacouleti* Watari, 1941

Material examined: 4 specimens: Karmat Ali; (2♀♀+2♂♂, Feb. 2022).

Distribution: Iraq [40]. Jordon [42] ; Turkey[49]. Pakistan[48]; Florida[47].; and Kuwait[46].

### 3.2. Family: Lycaenidae

#### 3.2.1. Genus: *Lampides* Hubner, 1819

- Synonyms: *Cosmolyce* Toxopeus, 1927
- *Lampidella* Hemming, 1933
- ***Lampides boeticus*** (Linnaeus, 1767)
- Common name: Bean Butterfly
- Synonyms: *Cosmolyce boeticus* (Linnaeus)
- *Hesperia boetica* Fabricius, 1793
- *Lampides anamariae* Gómez Bustillo, 1973
- *L. armeniensis* Gerhard, 1882
- *L. caerulea* Tutt, 1907
- *L. caeruleafasciata* Tutt, 1907
- *L. caeruleamarginata* Tutt, 1907
- *Lampides clara* Tutt, 1907
- *L. clarafasciata* Tutt, 1907
- *L. claramarginata* Tutt, 1907
- *L.fusca* Tutt, 1907
- *L. fuscafasciata* Tutt, 1907
- *L. fuscomarginata* Tutt, 1907
- *L. grisescens* Tutt, 1907
- *L. kawachensis* Hirose, 1934
- *L. leguminis* Scott, 1890
- *L. major* Tutt, 1907
- *L. minor* Tutt, 1907
- *L. typicafasciata* Tutt, 1907
- *L. typicamarginata* Tutt, 1907
- *Lycaena albovittata* Oberthür, 1910
- *Ly. boetica* (Linnaeus, 1767)
- *Ly.boetica armeniensis* Gerhard, 1882
- *Ly. ecaudata* Oberthür, 1910
- *Ly. minor* Pionneau, 1929
- *Papilio boeticus* Linnaeus, 1767
- *P. damoetes* Fabricius, 1775
- *P. pisorum* Fourcroy, 1785
- *Polyommatus boeticus* (Linnaeus, 1767)
- *Tarucus boeticus boeticoides* Stauder, 1925
- *T. boeticus immarginata* Stauder, 1925
- *T.boeticus nigrita* Stauder, 1925
- *T.s boeticus pygmaea* Stauder, 1925

Material examined: 118 specimens: 66(♀♀+♂♂) Karmat Ali;( 8. Des. 2021; 12. Jan. 2022; 20. Feb. 2022; 26. March. 2022). 52(♀♀+♂♂) Al- Zubair: ( 15,Des.2021 and 7. Jan. 2022; 10. Feb. 2022; 20. March. 2022).

Distribution: Iraq [55]; Jordon[42]; Arminia(Dachenko *et al.*,2011 ); Albaniab(Verovnik and Popovic, 2013); Azerbaijan[38].

#### 3.2.2. Genus: *Tarucus* Moore, 1881

- *Tarucus balkanicus* (Freyer, 1845)

- Common name: Little tiger blue
- Synonyms: *Lycaena balkanica* Freyer, 1843
- *Lycaena psittacus* Herrich-Schäffer, 1844
- *Tarucus balkanica* var. *areshana* Bethune-Baker, (1918)
- *Tarucus callinara* f. *nigra* Bethune-Baker, [1918]
- *Tarucus frivaldszkyi* Aigner-Abafi, 1906
- *Tarucus clorinda* Verity, 1938
- *Tarucus balkanica* (Freyer, 1844)

Material examined: 221 specimens: 109(♀♀+♂♂) Karmat Ali;(42. Nov. 2021; 8,Des.2021 and 12. Jan. 2022; 12. Feb. 2022; 37. March. 2022). 112(♀♀+♂♂) Al- Zubair: (32,Des.2021 and 22. Jan. 2022; 24. Feb. 2022; 37. March. 2022).

Distribution: Iraq[40]; Algeria and Tunisia(Tennet, 1996); Jordon[42] Albania(Verovnik and Popovic, 2013); Bangladesh(Neogi et al.,2014); Azerbaijan[38]; India(Basu et al., 2019); Armenia(Langourov, 2019); Serbia [68]; Croatia[70].

### 3.2.3. Genus: *Zizeeria Chapman, 1910*

- *Zizeria karsandra* (Moore, 1865)
- Common name: Dark grass blue
- Synonyms: *Lampides neis* Walker, 1870
- *Lycaena brahma* C.Felder & R.Felder, 1865
- *Lycaena conformis* Butler, 1877
- *Polyommatus karsandra* Moore, 1865

Material examined: 198 specimens: 118(♀♀+♂♂) Karmat Ali;(35. Nov. 2021; 8,Des.2021 and 20. Jan. 2022; 20. Feb. 2022; 35. March. 2022). 80(♀♀+♂♂) Al- Zubair: (25. Nov; .8Des.2021 and 18. Jan. 2022; 10. Feb. 2022; 12. March. 2022).

Distribution: Iraq [40]; Jordon[42]; Turkey[51], India[67] Sri Lanka[66] ; Iran [38]; Greece [65].

## 3.3. Family: Pieridae

Common name: Orang-Tips Butterflies

### 3.3.1. Genus: *Belenois Hubner, 1819*

- Synonyms: *Anaphaeis* Hubner, 1819
- *Glycestha* Billberg, 1820
- *Pseudohuphina* Stoneham, 1940
- *Pseudanaphaeis* Bernard, 1953
- *Belenois aurota* (Fabricius, 1793)
- Common name: Brown-Veined White
- Synonyms: *Anaphaeis aurotota* (Fabricius, 1793)
- *Papilio aurota* Fabricius, 1793

Material examined: 55 specimens: 33(♀♀+♂♂) Karmat Ali (20. Nov.; 2021 ; 3. Jan. 2022 ; 5. Feb. 2022; 5. Mar. 2022). Al- Zubair ;( 3.Nov.2021; 5. Jan. 2022; 5. Feb. 2022; 7. Mar. 2022).

Distribution: Iraq [64]; South Africa[62], Pakistan [60]; China[63].

### 3.3.2. Genus: *Colotis Hubner, 1819*

- Synonyms:*Calais* Boisduval, 1836
- *Cuneacolotis* Henning, Henning, Joannou & Woodhall, 1997
- *Idmais* Boisduval, 1836
- *Colotis fausta* (Olivier, 1804)
- Common name: Large Salmon Arab

Material examined: 75 specimens: 50(♀♀+♂♂) Karmat Ali (5. Jan. 2022; 20. Feb. 2022; 25. Mar. 2022). 25(♀♀+♂♂) Al- Zubair ;( 5. Jan. 2022;10. Feb. 2022; 10. Mar. 2022).

Iraq [40]; Turkey [59], Pakistan[60], Oman[6]and India [61].

### 3.3.3. Genus: *Pieris* Hübner, 1819

- Synonym: *Tachyptera* Berge, 1842
- Common name: white garden

### 3.3.4. *Pieris rapae* (Linnaeus, 1758)

- Common name: Cabbage Butterfly
- Synonyms: *Artogeia Rapae* (Linnaeus, 1758)
- *Papilio rapae* Linnaeus, 1758
- *Pieris brassiae* Var. *Crucivora* Boisduval, 1836
- *Pieris metra* (Stephens, 1827)

Material examined: 3 specimens (2♀♀+1♂) Karmat Ali, Nov. 2022

Distribution: Iraq [50]; Jordon[42]; Iran[53]; Pakistan [58]. Egypt [54] and Azerbaijan [38].

Family: Hesperiidae

### 3.3.5. Genus: *Pelopidas* Walker, 1870

- Synonym: Chapra Moore, 1881
- Peliophidas Swinhoe, 1913
- Pelopidae thrax(Hübner,1821)
- Common name: Millet skipper
- Synonyms: Gegenes Thrax Hübner, 1821
- *Pelopidas midea* Walker, 1870

Material examined: 43 specimens: 27(♀♀+♂♂) Karmat Ali (12Nov.2021, 4. Des.2021 and 3. Jan. 2022; 1. Feb. 2022; 7. Mar. 2022). 16(♀♀+♂♂) Al- Zubair ;( 4.Des.2021 and 5. Feb. 2022; 7. Mar. 2022).

Distribution: Iraq [50]. Turkey [51]; Oman[55], Greece [56]; Iran [57]; Egypt [52].

## References

- [1] Perveen, F. K., & Khan, A. (2017). Introductory Chapter: Lepidoptera. Lepidoptera. doi: 10.5772/intechopen.70452. In Perveen, F. K. (Ed.). (2017). *Lepidoptera*. BoD–Books on Demand. DOI: 10.5772/intechopen.70452, 154pp.
- [2] Kremen, C. (1992). Assessing the indicator properties of species assemblages for natural areas monitoring. *Ecological applications*, 2(2), 203-217.
- [3] Kremen, C. (1994). Biological inventory using target taxa: a case study of the butterflies of Madagascar. *Ecological applications*, 4(3), 407-422.
- [4] Kumar, S., Simonson, S. E., & Stohlgren, T. J. (2009). Effects of spatial heterogeneity on butterfly species richness in Rocky Mountain National Park, CO, USA. *Biodiversity and Conservation*, 18, 739-763.
- [5] Van Swaay, C. A., Dennis, E. B., Schmucki, R., Sevilleja, C., Balalaikins, M., Botham, M., ... & Roy, D. B. (2019). The EU Butterfly Indicator for Grassland species: 1990-2017. Technical report.23pp.
- [6] Schmidt, P., Simonyi, S. J., Ábrahám, L., Sáfián, S., & Ilniczky, S. (2020). A faunistic contribution to the butterfly fauna of Oman (Lepidoptera: Diurna). *Natura Somogyensis*, (34), 153-187.
- [7] Sharma, M., & Sharma, N. (2017). Suitability of butterflies as indicators of ecosystem condition: a comparison of butterfly diversity across four habitats in Gir Wildlife Sanctuary. *International Journal of Advanced Research in Biological Sciences*, 4(3), 43-53.
- [8] AL-Shehbaz, I. A., Mutlu, B., & Dönmez, A. A. (2007). The Brassicaceae (Cruciferae) of Turkey, updated. *Turkish Journal of Botany*, 31(4), 327-336.
- [9] Kapusta-Duch, J., Kopec, A., Piatkowska, E., Borczak, B., & Leszczynska, T. (2012). The beneficial effects of Brassica vegetables on human health. *Roczniki Państwowego Zakładu Higieny*, 63(4), 389-395.
- [10] Al-Mousawi, A. H. I. 1987. Plant Taxonomy. First Edition. Dar Alkutb for printing and publishing, University of Baghdad.379pp. (in Arabic).

- [11] Al-Abide, N. M. (2019). A morphological comparative study of some species of the Brassicaceae in the governorate of Erbil-Iraq. *Plant Archives*, 19(2), 289-293.
- [12] Khalil, M. A. I. (2004). Vegetable Plants / Propagation, Nurseries, Cell Culture And plant tissues, division, botanical description, varieties. Knowledge facility in Alexandria. Republic of Egypt Arabic, 540 p.
- [13] Sharma, H. C., & Dhillon, M. K. (2020). Climate change effects on arthropod diversity and its implications for pest management and sustainable crop production. *Agroclimatology: Linking Agriculture to Climate*, 60, 595-619.
- [14] Crepet, W. L. (1984). Advanced (constant) insect pollination mechanisms: pattern of evolution and implications vis-à-vis angiosperm diversity. *Annals of the Missouri Botanical Garden*, 607-630.
- [15] Preston, R. E. (1986). Pollen-ovule ratios in the Cruciferae. *American journal of botany*, 73(12), 1732-1740.
- [16] Abrol, D. P. (2012). *Pollination biology: biodiversity conservation and agricultural production* (p. 792). New York: Springer.
- [17] Aizen, M. A., Garibaldi, L. A., Cunningham, S. A., & Klein, A. M. (2008). Long-term global trends in crop yield and production reveal no current pollination shortage but increasing pollinator dependency. *Current biology*, 18(20), 1572-1575.
- [18] Ahuja, I., Rohloff, J., & Bones, A. M. (2011). Defence mechanisms of Brassicaceae: implications for plant-insect interactions and potential for integrated pest management. *Sustainable Agriculture*, 2, 623-670.
- [19] Ollerton, J., Winfree, R., & Tarrant, S. (2011). How many flowering plants are pollinated by animals?. *Oikos*, 120(3), 321-326.
- [20] Badenes-Pérez, F., Bhardwaj, T., & Thakur, R. (2017). Integrated pest management and pollination services in Brassica oilseed crops. *Integrated Management of Insect Pests on Canola and Other Brassica Oilseed Crops*, 341-349.
- [21] Dugdale, J. S. 1988: Lepidoptera - annotated catalogue, and keys to family-group taxa. *Fauna of New Zealand* 14, 264 pp.
- [22] Bashar, M. A., Mamun, M. A. and Rahman, SK. M. 2005. Wing Venation as a factor for the identification of Pierid butterflies in the forest of Bangladesh. *Bangladesh J. Zool.* 33(1): 49-56
- [23] Scudder, G. G. E., & Cannings, R. A. (2007). The Lepidoptera families and associated orders of British Columbia.
- [24] Naz, F., Ilyas, M., Ashfaque, M., & Shahzad, A. (2010). The genus Junonia (Lepidoptera: Nymphalidae) in Pakistan. *Sarhad J. Agric*, 26(2), 267-270
- [25] Abbas, M., M.A. Rafi, M. Inayatullah, M.R. Khan and H. Pavulaan (2002).Taxonomy and distribution of butterflies (Papilionoidea) of the Skardu region, Pakistan. *International Lepidoptera survey*. 3(9): 01-15
- [26] Dantchenko, A. V., Aghababyan, K. E., Khanamiryan, G. G., & Andreev, S. A. (2011). To the Fauna of Butterflies of Armenia. In Proceedings of the International conference:
- [27] Perveen, F., & Fazal, F. (2013). Key for identification of butterflies (Lepidoptera) of Hazara university, garden campus, Mansehra, Pakistan. *International Journal of Agriculture Innovations and Research*. 2013c, 1(5), 156-161.
- [28] Perveen, F. K. (2016). A contribution key for identification of butterflies (Lepidoptera) of Tehsil Tangi, Khyber Pakhtunkhwa, Pakistan. *Arthropods*, 5(3), 97-108.
- [29] Mehra, D., Kirti, J. S., & Sidhu, A. K. (2018). A taxonomic study of six species of the genus Junonia HÃ¼bner,[1819](Insecta: Lepidoptera: Nymphalidae) from the northwestern Himalayan region in India. *Journal of Threatened Taxa*, 10(7), 11934-11947.
- [30] Derwesh, A.I. 1965. A preliminary list of Identified Insects and some Arachnids of Iraq. Direct. Gen. of Res. and Agric. Proj., Minist. Agric. Bull. No. 121: 123 PP.
- [31] Dugdale, J. S. (1988). Lepidoptera-annotated catalogue, and keys to family-group taxa. *Fauna of New Zealand*, 14.
- [32] Larsen, T.B. (1994). Butterflies of Egypt. *Saudi Aramco world*; 45 (5):24-27.
- [33] Fermon H, Schulze CH, Waltert M, Mühlenberg M.(2001). The butterfly fauna of the Noyau Central, Lama Forest (Republic Of Benin), with notes on its ecological composition and geographic distribution. *African Entomology*, 9:17-26.
- [34] Katbeh-Bader, A., Amr, Z. S., Baker, M. A., & Mahasneh, A. (2003). The butterflies of Jordan. *Journal of Research on the Lepidoptera*, 37:11-26
- [35] Van Swaay, C., Cuttelod, A., Collins, S., Maes, D., Lopez Munguira, M., Šašić, M., Settele,

- [36] Karaçetin, E. and Welch, H.J. (2011). Red Book of Butterflies in Turkey. Ankara, Turkey: Doğa Koruma Merkezi. Available from: [www.dkm.org.tr].
- [37] Perveen, F. K. (Ed.). (2017). Lepidoptera. BoD–Books on Demand. DOI: 10.5772/intechopen.70452, 154pp.
- [38] Snegovaya, N. Y., & Petrov, V. A. (2019). A catalogue of butterflies (Lepidoptera, Rhopalocera) of Azerbaijan. *Acta Biologica Sibirica*, 5(3), 62-117.
- [39] Swaay, C. V., Cuttelod, A., Collins, S., Maes, D., López Munguira, M., Šašić, M., ... & Wynhoff, I. (2010). European red list of butterflies.
- [40] Wiltshire, E. P. (1957) . Lepidoptera of Iraq Revised and Enlarged edition . London , Nicholas kaye limited ( For Ministry of Agriculture Government of Iraq . 162 pp.
- [41] Ecuador, G. I. (1992). World distribution of the Vanessa cardui group (Nymphalidae). *Journal of the Lepidopterists' Society*, 46(3), 235-238.
- [42] Katbeh-Bader, A., Amr, Z. S., Baker, M. A., & Mahasneh, A. (2003). The butterflies of Jordan. na.
- [44] Karacatin, E. and Weleh, H. J. (2011).Red book of butterflies in Turkey.Doga koruma Merkezi (DKM).
- [45] Zarikian, N. (2016). A faunistic comparative study of families Hesperiidae and Nymphalidae (Lepidoptera: Rhopalocera) of Syrian Arab Republic and Republic of Armenia. *International Journal of Agricultural and Biosystems Engineering*, 10(8), 518-521.
- [46] Amr, Z. S. (2021). The state of biodiversity in Kuwait. IUCN. 247pp
- [47]. Lalonde, M. M., McCullagh, B. S., & Marcus, J. M. (2018). The taxonomy and population structure of the buckeye butterflies (genus Junonia, Nymphalidae: Nymphalini) of Florida, USA. *The Journal of the Lepidopterists' Society*, 72(2), 97-115.
- [48] Perveen, F. K. (2016). A contribution key for identification of butterflies (Lepidoptera) of Tehsil Tangi, Khyber Pakhtunkhwa, Pakistan. *Arthropods*, 5(3), 97.
- [49] Biricik, M. (2011). First record of Junonia orithya (Linnaeus, 1758)(Lepidoptera: Nymphalidae) in Turkey. *Zoology in the Middle East*, 53(1), 130-132.
- [50] Al- Ali , A. S . 1977 . Phytophagous and entomophagous insects and mites in Iraq . Natural History Research Centre ( Iraq ) Publication No . 33 , 142 pp .
- [51] Wagener, S. (2006). Butterfly diversity and protection in Turkey. *Bonner Zoologische Beiträge*, 54(1/2), 3.
- [52] Hendawy, A. S., Taha, A. S., Ghada, M. R., Ismael, M. M., & Heba, S. A. (2022). Rice skipper, Pelopidas thrax (hubner)(Hesperiidae: lepidoptera): An emerging insect pest in rice fields in Egypt, host plants, population fluctuation, damage and related natural enemies. *Journal of Entomology and Zoology Studies* 2022; 10(5): 16-22.
- [53] Hasanshahi, G., Jahan, F., Abbasipour, H., Karimi, J., & Askarianzadeh, A. (2013). Spatial Tehran. *Zoology and Ecology*, 23(3), 233-239
- [54] Salem, M. M., Mahmoud, Y. H. H., & Al-Azab, S. A. (2019). Survey and faunistic studies on moths and butterflies in New Valley Governorate, Egypt. 1. Butterflies. *Egyptian Journal of Plant Protection Research Institute*, 2(4), 629-640
- [55] Cock, M. J. (2009). Observations on the biology of Pelopidas thrax (Hübner)(Lepidoptera: Hesperiidae: Hesperiinae) in the Hajar Mountains, Oman. *Tribulus*, 18, 42-49.
- [56] Cuvelier, S. (2009). Pelopidas thrax, a new species for the Island of Kós and an update of its distribution in Greece (Lepidoptera: Hesperioidae & Papilionoidea). *Phegea*, 37(3), 90.
- [57] Zerganipour, A., Esfandiari, M., Naderi, A., & Rabieh, M. M. (2019). Identification of the butterflies (Lepidoptera, Papilionoidea) in the IMCA (Insect and Mite Collection of Ahvaz) at Shahid Chamran University of Ahvaz, Khuzestan, Iran. *Phegea*, 47(4), 129-139.
- [58] Mal, B.; Memon, N.; Shah, M. A. & Shah, N. A. (2013). Redescription and new record of small Arab group of Genus Colotis (Lepidoptera Pieridae) from Jamshoro, Sindh, Pakistan. Department of Zoology Univ. of Sindh Jamshoro. Sindh Univ. Res. J. (Sci. ser.) Vol. 45(2357-364.):
- [59] Kemal, M., Koçak, A. Ö., & Seven, E. (2010). Occurrence of Colotis fauna in South East Turkey (Lepidoptera, Pieridae). *Centre for Entomological Studies, Ankara, Cesa News*, 61, 2-8.
- [60] Noor, S., Ahmed, H. A., Mengal, F., Durrani, S., Rasheed, S., Abang, F., & Sani, I. A. (2018). An annotated list of the butterfly fauna of Quetta, Pakistan. *Journal of Entomology Zoology Studies*, 6, 771-777
- [61] Kaur, M., & Kirti, A. K. S. J. S. (2022). Taxonomic study of the genus Colotis Hubner,(1819)(Lepidoptera: Pieridae) from India. *The Pharma Innovation Journal SP-11(5)*: 1334-1349
- [62] Willis, C. K., & Woodhall, S. E. (2010). Butterflies of South Africa's National Botanical Gardens. South African National Biodiversity Institute. 232pp

- [63] Ge, S. X., Hu, S. J., Shi, H. L., Han, F. Y., Li, M. J., & Ren, L. L. (2021). The first record of the genus *Belenois* (Lepidoptera: Pieridae) from China. *Biodiversity Data Journal*, 9.
- [64] Verovnik, R., Verstraél, T., Warren, M., Wiemers, M. and Wynhof, I. 2010. European Red List of Butterflies. Publications Office of the European Union, 58pp.
- [65] Viborg, A. L. (2019). Supplementary information on Zizeeria karsandra (Lepidoptera: Lycaenidae) on Crete. *Phegea*, 47(3), 93-96.
- [66] Harinath, P., Suryanarayana, K., & Venkata Ramana, S. P. (2015). Eco-biology of the dark grass blue butterfly, Zizeeria karsandra (Moore)(Lepidoptera: Rhopalocera: Lycaenidae) from the Eastern Ghats of Southern Andhra Pradesh. *Journal of Entomology and Zoology Studies*, 3(5), 225-231.
- [67] Venkata Ramana, S. P., Harinath, P., & Prasanna Kumar, V. (2014). Life history of the dark grass blue Zizeeria karsandra (Lepidoptera: Rhopalocera: Lycaenidae) From Southern Andhra Pradesh-India. Centre for Info Bio Technology (CIBTech) *Journal of Zoology*, 3(1), 1-6.
- [68] Tot, I., Djurić, M., Vukajlović, F., & Vujić, M. (2021). Little tiger blue, *Tarucus balkanicus* (Freyer, 1845)—new butterfly species in the fauna of Serbia. *Acta entomologica serbica*, 26(2), 85-88.
- [69] Kunte, K. (2019). The genus Tarucus Moore,[1881](Lepidoptera: Lycaenidae) in the Indian Subcontinent. *Tropical Lepidoptera Research*, 87-110.
- [70] Koren, T., Laus, B., Burić, I., Štih, A., Kulijer, D., Dender, D., ... & Verovnik, R. (2022). New records of Tarucus balkanicus at the northern limit of its distribution
- [71] Swaay, C. V., Cuttelod, A., Collins, S., Maes, D., López Munguira, M., Šašić, M., ... & Wynhoff, I. (2010). European red list of butterflies. Published by IUCN (International Union for Conservation of Nature) and Butterfly Conservation Europe in collaboration with the European Union. 58pp.