

Fungal diversity associated with *Bemisia tabaci* (Hemiptera: Aleyrodidae) on cucumber and comparative effectiveness of bioassay methods in identifying the most virulent entomopathogenic fungi

Abdulnabi Abdul Ameer Matrood¹, Abdelhak Rhouma^{2,*}, Lobna Hajji-Hedfi² & Mohammad Imad Khriebea³

¹ Department of Plant Protection, College of Agriculture, University of Basrah, Iraq

² Regional Centre of Agricultural Research of Sidi Bouzid, CRRRA, Gafsa Road Km 6, B.P. 357, 9100, Sidi Bouzid, Tunisia

³ National Center for Biotechnology (NCBT), Researcher Doctor at NCBT, Damascus, Syria

Dr. Abdulnabi Abdul Ameer Matrood and Dr. Abdelhak Rhouma contributed equally in this work and contributed equally as first authors of this manuscript.

* e-mail: abdelhak.rhouma@gmail.com

Matrood A.A.A., Rhouma A., Hajji-Hedfi L. & Khriebea M.I. (2023) Fungal diversity associated with *Bemisia tabaci* (Hemiptera: Aleyrodidae) on cucumber and comparative effectiveness of bioassay methods in identifying the most virulent entomopathogenic fungi. – *Sydowia* 75: 269–282.