

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/339416049>

ON A GENERALIZATION OF LINEAR POSITIVE OPERATORS FOR FUNCTIONS OF GROWTH $\mu(x) + \mu^2(x)$ IN TWO DIMENSIONS ($\mu(x), \mu(x)$)

Article · June 2016

CITATIONS

0

1 author:



Haneen Jaafar

University of Basrah

8 PUBLICATIONS 1 CITATION

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Approximation Theory [View project](#)



Approximation theory [View project](#)

ON A GENERALIZATION OF LINEAR POSITIVE OPERATORS FOR FUNCTIONS OF GROWTH 2^{x+y} IN TWO DIMENSIONS (x, y)

Haneen J. Sadiq

University of Basrah. College of Science. Department of Mathematics

Haneen.mahdi@yahoo.com

Haneen.mah2010@gmail.com

Abstract:

In this paper, we introduce a generalization of linear positive operators $L_n^{\sim}(f; x)$ which we studied in [1], generalized in two dimensions (x, y) . We defined a sequence of linear positive operators to approximate unbounded functions in the domain $(\mathbb{R}_0 \times \mathbb{R}_0)$. We study some approximation properties for these operators $L_{n,m}^{\sim}(f; x, y)$ like Korovkin theorem and we proved Voronoviskaja – type asymptotic formula for the operators $L_{n,m}^{\sim}(f; x, y)$, whom we defined.

Key words: Linear positive operators, Korovkin Theorem, Voronoviskaja Theorem.

1.Introduction

In 1995 Lupas[9], defined and studied the identity;

$$\frac{1}{(1-a)^\alpha} = \sum_{k=0}^{\infty} \frac{(\alpha)_k}{k!} a^k, \quad |a| < 1; \quad \text{Where } (\alpha)_k = \begin{cases} \alpha(\alpha+1) \dots (\alpha+k-1) & k \in \mathbb{N} := \{1, 2, \dots\} \\ 1 & k = 0 \end{cases} \quad (1.1)$$

for more details, see [9],[2],[4] and [5] and he introduced the linear positive operators

$$L_n(f; x) = (1-a)^{nx} \sum_{k=0}^{\infty} \frac{(nx)_k}{k!} a^k f\left(\frac{k}{n}\right).$$

By letting $\alpha = nx$ and $x \geq 0$. Where $f: \mathbb{R}_0 \rightarrow \mathbb{R}$. In 1999 Agratini [2], Supposing that $L_n(1; x) = 1$, he found that $a = \frac{1}{2}$. Hence Lupas defined the linear positive operators;

$$L_n(f; x) = 2^{-nx} \sum_{k=0}^{\infty} \frac{(nx)_k}{2^k k!} f\left(\frac{k}{n}\right) \quad (1.2)$$

In 2007 A. Erencyn and Fatma Tasedelen, [5], define a sequence of positive linear operators and study some approximation properties for it that he fine the generalization of the operators (1.2) above, so he define this operators:

$L_n(f; x) = 2^{-a_n x} \sum_{k=0}^{\infty} \frac{(a_n x)_k}{2^k k!} f\left(\frac{k}{b_n}\right)$, where $x \in \mathbb{R}_0, n \in \mathbb{N}, \{a_n\}, \{b_n\}$ some increasing and unbounded sequence. In 2014 Mohammed and Sadiq [1] defined a new sequence of positive and linear operators as follow:

$$L_n^{\sim}(f; x) = \frac{1}{G_x} 2^{-nx} \sum_{k=0}^{\infty} \frac{(nx)_{k+r}}{2^{k+r} (k+r)!} f\left(\frac{k+r}{n}\right) \quad (1.3)$$

where $G_x = 2^{-nx} \sum_{k=0}^r \frac{(nx)_k}{2^k k!}$ or $G_x = 2^{-nx} \sum_{k=0}^{\infty} \frac{(nx)_{k+r}}{2^{k+r} (k+r)!}$

$G_x := \sum_{k=0}^r d_{n,k}(x)$ or $G_x := \sum_{k=0}^{\infty} d_{n,k+r}(x)$

Then, it can be written the operator (1.3) as follow

$$L_n^{\sim}(f; x) = \frac{1}{G_x} \sum_{k=0}^{\infty} d_{n,k+r}(x) f\left(\frac{k+r}{n}\right) \quad \text{where } x \in \mathbb{R}_0, n \in \mathbb{N}, \mathbb{R}_0 = [0, \infty), \mathbb{N} := \{1, 2, \dots\} \text{ and } r \in \mathbb{N}.$$

In this paper, we define and study the operators $L_{n,m}^{\sim}(f; x, y)$ which represents the generalization of the operators (1.3) in two dimensions (x, y) ([8], [6] and [12]) as follow:

$$L_{n,m}^{\sim}(f; x, y) = \frac{1}{G_x G_y} \sum_{k=0}^{\infty} d_{n,k+r}(x) \sum_{j=0}^{\infty} d_{m,j+s}(y) f\left(\frac{k+r}{n}, \frac{j+s}{m}\right) \quad (1.4)$$

where $d_{n,k+r}(x) = 2^{-nx} \sum_{k=0}^{\infty} \frac{(nx)_{k+r}}{2^{k+r} (k+r)!}$

and $x, y \in \mathbb{R}_0, f \in C_{\rho,q}(\mathbb{R}_0 \times \mathbb{R}_0),$

$C_{\rho,q}$: The space of all continuous and unbounded functions f on the area $\mathbb{R}_0 \times \mathbb{R}_0,$ with growth $(O(2^{\rho+q})),$ for some $\rho, q > 0,$ with the norm

$$\|f(x, y)\|_{\rho,q} = \text{Sup}_{(x,y) \in (\mathbb{R}_0 \times \mathbb{R}_0)} \frac{|f(x,y)|}{\rho(x)q(y)}, \text{ where } \rho(x) = 2^x.$$

2. Main Results:

Firstly, we study the convergent conditions on the operators (1.4), and before that we need to offer some theorems (Korovkin theorem [3] and [11]) Mohammed and Sadiq proved in 2014 [1], on the operators (1.3):

Theorem(2.1) (Korovkin Theorem):[1]

For $x \in \mathbb{R}_0, f \in C_{\rho}$ and by applying Korovkin Theorem on the operator $L_n^{\sim}(f; x),$ we have:

- 1) $L_n^{\sim}(1; x) = 1$
- 2) $L_n^{\sim}(t; x) = x + \frac{2r}{n G_x} d_{n,r}(x).$
- 3) $L_n^{\sim}(t^2; x) = x^2 + \frac{2x}{n} + r d_{n,r}(x) \left[\frac{6nx+2r+6}{3n^2 G_x} \right]$
- 4) $L_3^{\sim} = \frac{8}{7} r d_{n,r}(x) \left[r^2 + \frac{(nx+r)(3r+nx+5)+21+nx(33+7nx)+4r(nx+3)}{4} \right]$
- 5) $L_4^{\sim} = +n^4 x^4 G_x + 12n^3 x^3 G_x + 36n^2 x^2 G_x + 26nx G_x$
 $+ \frac{16}{15} r d_{n,r}(x) \left[\begin{array}{l} 1.8214n^3 x^3 + 15.3362n^2 x^2 + 34.6249nx + 1.4285 + r^3 + 1.6071r^2 \\ + 11.5892r + 0.4999r^2 nx + 0.75rn^2 x^2 + 5.9642rnx \end{array} \right].$

Lemma (2.2): For the operator $L_n^{\sim}(f; x, y),$ where $f(t, u) \in C_{\rho,q}(\mathbb{R}_0 \times \mathbb{R}_0),$ we have:
 $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(f; x, y) - f(x, y)\|_{\rho,q} = 0$ if and only if

- 1) $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(1; x, y) - 1\|_{\rho,q} = 0 \quad ; \quad (2.1)$
- 2) $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(t; x, y) - x\|_{\rho,q} = 0 \quad ; \quad (2.2)$
- 3) $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(u; x, y) - y\|_{\rho,q} = 0 \quad ; \quad (2.3)$
- 4) $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(t^2 + u^2; x, y) - (x^2 + y^2)\|_{\rho,q} = 0 \quad ; \quad (2.4)$

for $x, y \in \mathbb{R}_0$ and $f \in C_{\rho,q}(\mathbb{R}_0 \times \mathbb{R}_0).$

Proof: Clearly,

- 1) $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(1; x, y) - 1\|_{\rho,q} =$
 $\left| \frac{1}{G_x} \frac{1}{G_y} \sum_{k=0}^{\infty} d_{n,k+r}(x) \sum_{j=0}^{\infty} d_{m,j+s}(y) - 1 \right| = 0$
- 2) $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(t; x, y) - x\|_{\rho,q} = \left| \frac{1}{G_x} \frac{1}{G_y} \sum_{k=0}^{\infty} \left(\frac{k+r}{n} \right) d_{n,k+r}(x) \sum_{j=0}^{\infty} d_{m,j+s}(y) - x \right| =$
 $\left| \frac{1}{G_x} \sum_{k=0}^{\infty} \left(\frac{k+r}{n} \right) d_{n,k+r}(x) - x \right|$

By using theorem (2.1) (2) we get;

$$L_n^{\sim}(t; x) = x + \frac{2r}{n G_x} d_{n,r}(x)$$

that means $\frac{1}{G_x} \sum_{k=0}^{\infty} \left(\frac{k+r}{n} \right) d_{n,k+r}(x) = x + \frac{2r}{n G_x} d_{n,r}(x)$ and this a broach to x as $n \rightarrow \infty.$

Then $\left| \frac{1}{G_x} \sum_{k=0}^{\infty} \left(\frac{k+r}{n} \right) d_{n,k+r}(x) - x \right| \rightarrow 0$ as $n \rightarrow \infty.$

Therefore $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(t; x, y) - x\|_{\rho,q} = \frac{2r}{n G_x} d_{n,r}(x)$ which tend to 0 as $n \rightarrow \infty.$

We proved (2.1) and (2.2), by the same way we shall prove (2.3)

$$3) \lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(u; x, y) - y\|_{\rho,q} = \left| \frac{1}{G_x} \frac{1}{G_y} \sum_{k=0}^{\infty} d_{n,k+r}(x) \sum_{j=0}^{\infty} \binom{j+s}{m} d_{m,j+s}(y) - y \right| = \left| \frac{1}{G_y} \sum_{j=0}^{\infty} \binom{j+s}{m} d_{m,j+s}(y) - y \right|$$

By using theorem (2.1) (2) we have;

$$L_m^{\sim}(u; y) = y + \frac{2s}{m G_y} d_{m,s}(y);$$

$$\text{Then } \left| \frac{1}{G_y} \sum_{j=0}^{\infty} \binom{j+s}{m} d_{m,j+s}(y) - y \right| \rightarrow 0 \text{ as } m \rightarrow \infty.$$

Therefore $\lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(u; x, y) - y\|_{\rho,q} = \frac{2s}{m G_y} d_{m,s}(y)$ which tend to 0 as $m \rightarrow \infty$.

Now, we want to prove (2.4) bellow;

$$4) \lim_{n,m \rightarrow \infty} \|L_{n,m}^{\sim}(t^2 + u^2; x, y) - (x^2 + y^2)\|_{\rho,q} \\
= \lim_{n,m \rightarrow \infty} \left\| L_{n,m}^{\sim} \left(\left(\frac{k+r}{n} \right)^2 + \left(\frac{j+s}{m} \right)^2 ; x, y \right) - (x^2 + y^2) \right\|_{\rho,q} \\
= \left| \frac{1}{G_x} \frac{1}{G_y} \sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left(\left(\frac{k+r}{n} \right)^2 + \left(\frac{j+s}{m} \right)^2 \right) d_{n,k+r}(x) d_{m,j+s}(y) - (x^2 + y^2) \right| \\
= \left(\frac{1}{G_x} \sum_{k=0}^{\infty} \left(\frac{k+r}{n} \right)^2 d_{n,k+r}(x) - x^2 \right) \left(\frac{1}{G_y} \sum_{j=0}^{\infty} \binom{j+s}{m} d_{m,j+s}(y) - y^2 \right)$$

By using theorem (2.1) (3) we have;

$$L_n^{\sim}(t^2; x) = x^2 + \frac{2x}{n} + r d_{n,r}(x) \left[\frac{6nx+2r+6}{3n^2 G_x} \right],$$

$$\text{So, } \left(\frac{1}{G_x} \sum_{k=0}^{\infty} \left(\frac{k+r}{n} \right)^2 d_{n,k+r}(x) - x^2 \right) = \frac{2x}{n} + r d_{n,r}(x) \left[\frac{6nx+2r+6}{3n^2 G_x} \right]$$

which tend to 0 when $n \rightarrow \infty$.

$$\text{and } \frac{1}{G_y} \sum_{j=0}^{\infty} \binom{j+s}{m} d_{m,j+s}(y) - y^2 = \frac{2y}{m} + s d_{m,s}(y) \left[\frac{6my+2s+6}{3m^2 G_y} \right]$$

which tend to 0 as $m \rightarrow \infty$.

So the proof is complete. ■

Later, we shall prove Voronovskaja theorem on the operators (1.4) and to prove it we need the next lemma to get the moment to the operators (1.3) which proved by Mohammed and Sadiq in 2014 in [1].

Lemma (2.3):[11] Let $r \in \mathbb{N}$, then for all $x \in \mathbb{R}_0$ and $n \in \mathbb{N}$, we have

- 1) $T_{n,0}^{\sim}(x) = 1.$
- 2) $T_{n,1}^{\sim}(x) = \frac{2r}{n G_x} d_{n,r}(x).$
- 3) $T_{n,2}^{\sim}(x) = r d_{n,r}(x) \left(\frac{6nx+2r+6}{3n^2 G_x} + \frac{4x}{n G_x} \right) + \frac{2x}{n}.$
- 4) $T_{n,3}^{\sim}(x) = r d_{n,r}(x) \left(\frac{8r^2+2(nx+r)(3r+nx+5)+42+2nx(33+7nx)+8r(nx+3)}{7n^3 G_x} - \frac{2rx+6}{n^2 G_x} \right) + \frac{6x}{n^2}.$
- 5) $T_{n,4}^{\sim}(x) = r d_{n,r}(x) \left[5.9428 \frac{x^3}{n G_x} + 28.358 \frac{x^2}{n^2 G_x} + 36.9332 \frac{x}{n^3 G_x} + 4.8 \frac{rx^2}{n^2 G_x} + 0.3618 \frac{rx}{n^3 G_x} + \frac{1.5237}{n^4 G_x} + 1.0666 \frac{r^3}{n^4 G_x} + 1.7142 \frac{r^2}{n^4 G_x} - 7.9999 \frac{r^2}{n^3 G_x} + 12.3618 \frac{r}{n^4 G_x} + 0.5332 \frac{xr^2}{n^3 G_x} - 9.1428 \frac{rx}{n^2 G_x} - 9.1428 \frac{x^2}{n G_x} - 9.4285 \frac{x}{n^2 G_x} - 19.4285 \frac{r}{n^3 G_x} - \frac{24}{n^3 G_x} \right] + \frac{12x^2}{n^2} + \frac{36x}{n^3}.$

Now, we ready to state and prove Voronovskaja theorem

Theorem (2.4):[11] (Voronovskaja theorem)

Suppose that $f \in C_{\rho,q}(\mathbb{R}_0 \times \mathbb{R}_0)$, $\rho, q \in \mathbb{N}^0$, and suppose that

$\frac{\partial^2 f(x,y)}{\partial x^2}$ and $\frac{\partial^2 f(x,y)}{\partial y^2}$ exist at a point $(x, y) \in \mathbb{R}_0 \times \mathbb{R}_0$.

Then

$$\lim_{n \rightarrow \infty} n \left(L_{n,m}^{\sim}(f(t, u); x, y) - f(x, y) \right) = x \frac{\partial^2 f(x,y)}{\partial x^2} + y \frac{\partial^2 f(x,y)}{\partial y^2}.$$

Proof: By using Taylor's expansion for $f(t, u)$ about (x, y) , we have; [10]

$$f(t, u) = f(x, y) + f'_x(x, y)(t - x) + f'_y(x, y)(z - y) + \frac{1}{2}(f''_{xx}(x, y)(t - x)^2 + 2f''_{xy}(x, y)(u - y)^2) + \varphi(t, u; x, y)\sqrt{(t - x)^4 + (u - y)^4}$$

where $(t, u) \in (0, \infty) \times (0, \infty)$, and $\varphi(t, u) = \varphi(t, u; x, y)$ is a function belonging to $C_{\rho, q} \in [0, \infty) \times [0, \infty)$ and $\varphi(t, u) \rightarrow 0$ when $(t, u) \rightarrow (x, y)$ for $n, m \in \mathbb{N}$.

So, by depending on $L_{n,n}(f; x)$ is a sequence of linear positive operators, we have;

$$\begin{aligned} L_{n,n}(f(t, u); x, y) &= f(x, y) + \frac{\partial f(x, y)}{\partial x} L_n((t - x); x) \\ &+ \frac{\partial f(x, y)}{\partial y} L_n((u - y); y) \\ &+ \frac{1}{2} \left[\frac{\partial^2 f(x, y)}{\partial x^2} L_n((t - x)^2; x) + \frac{\partial^2 f(x, y)}{\partial x \partial y} L_n((t - x); x) L_n((u - y); y) + \frac{\partial^2 f(x, y)}{\partial y^2} L_n((u - y)^2; y) \right] + \\ &L_{n,n}(\varphi(t, u)\sqrt{(t - x)^4 + (u - y)^4}; x, y) \end{aligned}$$

Here, using lemma (2.3), we have:

$$\begin{aligned} \lim_{n \rightarrow \infty} n \left(L_{n,n}(f(t, u); x, y) - f(x, y) \right) &= \\ \frac{\partial f(x, y)}{\partial x} \frac{2r}{nG_x} d_{n,r}(x) + \frac{\partial f(x, y)}{\partial y} \frac{2r}{nG_y} d_{n,r}(y) + \\ \frac{1}{2} \left[\frac{\partial^2 f(x, y)}{\partial x^2} \left\{ r d_{n,r}(x) + \left(\frac{6nx+2r+6}{3n^2G_x} + \frac{4x}{nG_x} \right) \frac{2x}{n} \right\} + \frac{\partial^2 f(x, y)}{\partial x \partial y} \frac{2r}{nG_x} d_{n,r}(x) \frac{2r}{nG_y} d_{n,r}(y) + \frac{\partial^2 f(x, y)}{\partial y^2} \left\{ r d_{n,r}(y) + \left(\frac{6ny+2r+6}{3n^2G_y} + \frac{4y}{nG_y} \right) \frac{2y}{n} \right\} \right] + \\ n L_{n,n}(\varphi(t, u)\sqrt{(t - x)^4 + (u - y)^4}; x, y) \end{aligned}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} n \left(L_{n,n}(f(t, u); x, y) - f(x, y) \right) &= \\ x \frac{\partial^2 f(x, y)}{\partial x^2} + y \frac{\partial^2 f(x, y)}{\partial y^2} + \lim_{n \rightarrow \infty} n L_{n,n}(\varphi(t, u)\sqrt{(t - x)^4 + (u - y)^4}; x, y). \end{aligned}$$

By using the Cauchy - Schwartz inequality; we get:

$$\begin{aligned} & \left| L_{n,n}(\varphi(t, u)\sqrt{(t - x)^4 + (u - y)^4}; x, y) \right| \\ & \leq \left(L_{n,n}(\varphi^2(t, u); x, y) \right)^{1/2} \cdot \left(L_n((t - x)^4; x) + L_n((u - y)^4; y) \right)^{1/2} \\ & \leq \varphi^2 \left(L_n((t - x)^4; x) + L_n((u - y)^4; y) \right)^{1/2} \end{aligned}$$

Which tend to zero when n tend to ∞ , (by applying lemma {(2.3) (5)}).

Hence, $\lim_{n \rightarrow \infty} n L_{n,n}(\varphi(t, u)\sqrt{(t - x)^4 + (u - y)^4}; x, y) = 0$.

Therefore,

$$\lim_{n \rightarrow \infty} n \left(L_{n,n}(f(t, u); x, y) - f(x, y) \right) = x \frac{\partial^2 f(x, y)}{\partial x^2} + y \frac{\partial^2 f(x, y)}{\partial y^2}. \quad \blacksquare$$

References

- [1] Ali J. Mohammed and Haneen J. Sadiq, A sequence of linear and positive operators for The functions of growth 2^x , *Journal of Basrah Research ((Sciences))*, **40**, 87-93. (2014).
- [2] O. Agratini, On a sequence of linear and positive operators, *Facta Universitatis(Nis), Ser. Math. Inform.*, **14** (1999), 41-48.
- [3] Allan Pinkus: Weierstrass and Approximation Theory, *J.Approx. Theory* **107** (2000),1-66.

- [4] F. Dirik , Statistical convergence and rate of convergence of sequence of positive linear Operators, *Mathematical communications*, **12**(2007), 147-153.
- [5] A. Erencyn and Fatma Tasdelen, on a family of linear and positive operators in weighted spaces, *Jornal of inequalities in pure and applied mathematics*, **8** (2007), 1-11.
- [6] İbrahim Büyükyazıcı ^{a,*}, Ertan İbikli ^{b,*}: The approximation properties of generalized Bernstein polynomials of two variables. *Applied Mathematics and Computation* .**156** (2004), 367-380.
- [7] H.S. Kasana and P.N. Agrawal: On Sharp estimates and linear combinations of modified Bernstein polynomials, *Bull. Soc. Math. Belg.* **40**(1), Ser B(1988), 61-71.
- [8] Lucyna Rempulska and Mariola Skorupka : On modified Meyer- König and Zeller Operators of functions of two variables. *Archivum Mathematicum (BRNO). Tomus.* **42** (2006), 273-284.
- [9] A. Lupas, The approximation by some positive linear operators, In: *Proceeding of the International Dortmund Meeting on Approximation Theory. (M.W. Müller et al., eds.)*, Akademie Verlag, Berlin (1995), 201–209.
- [10] S. Tarabie, On some α - statistical approximation processes, *International journal of pure and applied mathematics*, **67**, (2012), 327-332.
- [11] E.Voronovskaja: Détermination de la forme asymptotique d'approximation des fonctions par les polynômes de S. N. Bernstein, *C. R. Acad. Sci. USSR*(1932),79-85.
- [12] Zbigniew Walczak: Approximation properties of certain linear positive operators in polynomial weighted spaces of functions of one and two variables. *Univ. Beograd. Publ. Elektrotehn. Fak.* **14**(2003), 51- 64.