

Reproducing Kernel Hilbert Space Method for Solving Nonlinear Integro-Differential Equations

Wafaa K. Al-Zuhairi ^{*} , Hameeda O. Al-Humedi 

Department of Mathematics, College of Education for Pure Sciences, University of Basrah, Basrah, Iraq.

ARTICLE INFO

Received 9 August 2024
Revised 20 September 2024
Accepted 25 September 2024
Published 31 December 2024

Keywords :

Reproducing Kernel Hilbert Space Method, Approximate Solution, Nonlinear Integro-Differential Equations, Hilbert Space, Function Space.

ABSTRACT

In this work, the reproducing kernel Hilbert space method (RKHSM) was used to find a numerical solution to nonlinear integro-differential equations (NIDEs) on the form of finite series. The results showed that the approximate solution are converges to the exact solution of the NIDEs, which confirms the effectiveness of RKHSM as a reliable and practical tool for solving this type of e. This method is of great importance in the field of numerical solutions of NIDEs, which has wide applications in many engineering and scientific fields

Citation: W. K. Al-Zuhairi, H. O. Al-Humedi, J. Basrah Res. (Sci.) 50(2), 182 (2024).
[DOI:https://doi.org/10.56714/bjrs.50.2.15](https://doi.org/10.56714/bjrs.50.2.15)

1. Introduction

Integro-differential equations (IDEs) are important in describing physical, natural, and biological phenomena, as well as in engineering and biological technologies, in recent years, they have received increasing attention in many different disciplines. Such as elasticity theory, biomechanics, electromagnetism, industry and scientific applications, and others [1, 2, 3].

RKHSM considered one of the effective methods to solve many papers, including differential and integral equations which have widely used in statistics and probability.

The following formula of NIDEs is given by [4]:

*Corresponding author email : pgs.Wafaa.kamel@uobasrah.edu.iq



$$\sum_{k=1}^q b_k(t) g^{(k)}(t) = w(t) + \sigma \int_{t_0}^{t_1 \text{ or } t} T(t, z) F(g(z)) dz, \quad (1)$$

the initial condition $g^{(k)}(t_0) = r_k \quad k = 1, \dots, q,$

where $b_k(t)$, $w(t)$ and $T(t, z)$ are known functions so that $t \in [t_0, t_1]$, $t_0, t_1, \sigma, r_k, q \in \mathbb{R}$, $F(g(t))$ is a nonlinear operator and $g(t)$ is an unknown function in $W_2^i[t_0, t_1]$, $i = 1, 2, \dots$.

IDEs have wide applications in various fields, during the last decades, many numerical methods have been developed to solve these equations, whether linear or nonlinear. These methods include Adomian decomposition method by [5] collection method [6, 7, 8, 9], Taylor polynomial [10], cubic B-spline least-square method combine with a quadratic weight function [11], cubic B-spline Galerkin method combine with a quadratic weight function [12], least-squares method [13, 14], homotopy perturbation method [15], the simplified reproducing kernel method and the homotopy perturbation method [16], Laplace discrete Adomian decomposition method [17], other numerical methods are used to solve IDEs, since these equations are difficult to solve using analytical methods.

The RKHSM has been used to solving many problems, such as singular IEs [18, 19, 20, 21], linear IEs [22, 23, 24], nonlinear IEs [25, 26, 27] and all equations were solved using the Gram-Schmidt algorithm. However, the researchers below worked on a new approach, where they solved a nonlinear IEs [28], solved a class of functional IEs [29] and solved third-order differential equations [30].

The main purpose of this paper is to apply RKHSM for solving NIDEs. The convergence and error analysis will be discussed.

It is organized into five parts, including the introduction. The second part gives the basic definitions of the reproducing kernel, inner product, norm, function space, and the form of the reproducing kernel. The third part explains how to apply the method to NIDEs, and also discusses the convergence theory. The fourth part gives the numerical results and explains the error magnitude, and the fifth part contains the conclusion.

2. Reproducing Kernel Hilbert Space

In this part, It has been defined the function space $W_2^i[t_0, t_1]$ where $i = 1, 2, \dots$ and define the kernel $G_s(t)$ in any space in the research that relies on $W_2^i[t_0, t_1]$, [31]:

Definition 2.1. Let it be S are abstract set so that $S \neq \phi$, a function $G: S \times S \rightarrow \mathbb{C}$, where $\mathbb{C}$ be the set of complex numbers G is a reproducing kernel of H if satisfied

- $\forall s \in S, G_s(t) \in H,$
 - $\forall s \in S, \forall g \in H, g(s) = \langle G_s(t), g(t) \rangle.$
- (2)

Definition 2.2. The definition of function space $W_2^i[t_0, t_1]$, $i = 1, 2, \dots$ is as follows:

$$W_2^i[t_0, t_1] = \{g(t): g: [t_0, t_1] \rightarrow \mathbb{R}, g^{(i-1)} \text{ continuous function}, g^{(i)} \in L^2[t_0, t_1]\},$$

The inner product and the norm in $W_2^i[t_0, t_1]$ are defined as follows.

$$\langle g, f \rangle_{W_2^i[t_0, t_1]} = \sum_{k=0}^{i-1} g^{(k)}(t_0) f^{(k)}(t_0) + \int_{t_0}^{t_1} g^{(i)}(t) f^{(i)}(t) dt, \quad (3)$$

$$\|g\|_{W_2^i[t_0, t_1]} = \sqrt{\langle g, g \rangle_{W_2^i[t_0, t_1]}} \quad (4)$$

Remark 2.1. [25] If $\forall t \in [t_0, t_1]$ and $\forall g(t) \in W_2^i[t_0, t_1] \exists G_s(t) \in W_2^i[t_0, t_1]$ such that $\langle G_s(t), g(s) \rangle_{W_2^i[t_0, t_1]} = g(t)$, where $s \in [t_0, t_1]$ then the Hilbert space $W_2^i[t_0, t_1]$ is a reproducing kernel space.

Define the reproducing kernel of $W_2^i[t_0, t_1]$ is

$$G_s(t) = \begin{cases} G_1(t) = \sum_{k=1}^{2i} a_k(s)t^{k-1}, & t \leq s \\ G_2(t) = \sum_{k=1}^{2i} v_k(s)t^{k-1}, & t > s \end{cases} \quad (5)$$

the coefficients $a_k(s)$, $v_k(s)$, $\{k = 1, 2, \dots, 2i\}$, can be obtained by.

$$\frac{\partial^k G_1(s)}{\partial t^k} = \frac{\partial^k G_2(s)}{\partial t^k}, \quad k = 0, 1, \dots, 2i - 2, \quad (6)$$

$$\text{and} \quad (-1)^i \left(\frac{\partial^{2i-1} G_1(s^+)}{\partial t^{2i-1}} - \frac{\partial^{2i-1} G_2(s^-)}{\partial t^{2i-1}} \right) = 1. \quad (7)$$

Remark 2.2. By using (6) and (7), we will calculate $G_s(t)$ in $W_2^i[0,1]$ when $i = 1, 2, 3, 4$
 $i = 4$,

$$G_s^4(t) = \begin{cases} \frac{-t^7}{5040} + \frac{(t^6s)}{720} - \frac{(t^5s^2)}{240} + \frac{(t^4s^3)}{144} + \frac{(t^3s^3)}{36} + \frac{(t^2s^2)}{4} + ts + 1, & t \leq s \\ \frac{-s^7}{5040} + \frac{(s^6t)}{720} - \frac{(s^5t^2)}{240} + \frac{(s^4t^3)}{144} + \frac{(s^3t^3)}{36} + \frac{(s^2t^2)}{4} + ts + 1, & t > s \end{cases}$$

$i = 3$,

$$G_s^3(t) = \begin{cases} \frac{t^5}{120} - \frac{(t^4s)}{24} + \frac{(t^3s^2)}{12} + \frac{(t^2s^2)}{4} + st + 1, & t \leq s \\ \frac{s^5}{120} - \frac{(s^4t)}{24} + \frac{(s^3t^2)}{12} + \frac{(s^2t^2)}{4} + st + 1, & t > s \end{cases}$$

$i = 2$

$$G_s^2(t) = \begin{cases} -\frac{t^3}{6} + \frac{(t^2s)}{2} + st + 1, & t \leq s \\ -\frac{s^3}{6} + \frac{(s^2t)}{2} + st + 1, & t > s \end{cases}$$

$i = 1$,

$$G_s^1(t) = \begin{cases} t + 1, & t \leq s \\ s + 1, & t > s \end{cases}$$

3. The Implementation of Method

In this part, it is mentioned how RKHSM is applied to solve NIDEs and related theories are mentioned. In many previous studies, researchers used the Gram-Schmidt algorithm to obtain an orthogonal system. However, there are some problems associated with using this algorithm, namely numerical instability due to the large number of computational steps required to implement it. Therefore, in this study, an alternative algorithm that is numerically stable and less computationally complicated will be used.

The equation (1) can be expressed in the following form:

$$Lg(t) = \sum_{k=1}^q b_k(t)g^{(k)}(t) - \sigma \int_{t_0}^{t_1 \text{ or } t} T(t, z)F(g(z))dz \quad (8)$$

$$Lg(t) = w(t),$$

where $L: W_2^i[t_0, t_1] \rightarrow W_2^1[t_0, t_1]$ is an invertible bounded linear operator.

Was chosen a countable set of points $\{t_k\}_{k=1}^\infty$ in the interval $[t_0, t_1]$ define

$$\delta_k(t) = G_{t_k}^1(t), \quad \mu_k(t) = L^* \delta_k(t),$$

where L^* is the adjoint operator of L .

Theorem 3.1. [31] Let $\{t_k\}_{k=1}^\infty$ is dense on $[t_0, t_1]$ then $\{\mu_k(t)\}_{k=1}^\infty$ is a complete function system of $W_2^i[t_0, t_1]$ and $\mu_k(t) = L_s G_{t_k}^i(t)$, where the operator L_s applies to function of s

Theorem 3.2. [28, 29, 30] Let $\{\mu_k(t)\}_{k=1}^{\infty}$ be a linear independent set in $W_2^i[t_0, t_1]$, and $\{\tilde{\mu}_k(t)\}_{k=1}^{\infty}$ be an orthogonal system in $W_2^i[t_0, t_1]$, so that $\tilde{\mu}_k(t) = \sum_{j=1}^k \beta_{kj} \mu_j(t)$, if

$$g(t) = \sum_{k=1}^{\infty} A_k \mu_k(t) \quad (9)$$

and the approximate solution

$$g_M(t) = \sum_{k=1}^M A_k \mu_k(t) = \sum_{k=1}^M \tilde{A}_k \tilde{\mu}_k(t) \quad (10)$$

then $\Gamma A = \bar{w}$,

where,

$$\Gamma = \begin{pmatrix} \langle L\mu_1, \mu_1 \rangle_{W_2^i[t_0, t_1]} & \cdots & \langle L\mu_1, \mu_M \rangle_{W_2^i[t_0, t_1]} \\ \vdots & \ddots & \vdots \\ \langle L\mu_M, \mu_1 \rangle_{W_2^i[t_0, t_1]} & \cdots & \langle L\mu_M, \mu_M \rangle_{W_2^i[t_0, t_1]} \end{pmatrix}, \quad A = \begin{pmatrix} A_1 \\ \vdots \\ A_M \end{pmatrix}, \quad \tilde{A} = \begin{pmatrix} \tilde{A}_1 \\ \vdots \\ \tilde{A}_M \end{pmatrix},$$

$$\bar{w} = \begin{pmatrix} \langle w, \mu_1 \rangle_{W_2^i[t_0, t_1]} \\ \vdots \\ \langle w, \mu_M \rangle_{W_2^i[t_0, t_1]} \end{pmatrix}, \quad \beta = \begin{pmatrix} \beta_{11} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ \beta_{1M} & \cdots & \beta_{MM} \end{pmatrix}, \text{ and } \beta_{jh} \text{ are orthogonal coefficients, } (\beta_{kk} >$$

$0, k, h = 1, 2, \dots, M$).

Proof. Let $g(t) \in W_2^i[t_0, t_1]$ then $g(t) = \sum_{k=1}^{\infty} A_k \mu_k(t) = \sum_{k=1}^{\infty} \tilde{A}_k \tilde{\mu}_k(t)$

To get an approximate solution, was cut the series at M .

$$\begin{aligned} g_M(t) &= \sum_{k=1}^M A_k \mu_k(t) = \sum_{k=1}^M \tilde{A}_k \tilde{\mu}_k(t) \\ &= \sum_{k=1}^M \tilde{A}_k \sum_{j=1}^k \beta_{kj} \mu_j(t) = \sum_{j=1}^M (\sum_{k=j}^M \tilde{A}_k \beta_{kj}) \mu_j(t) \end{aligned}$$

$\{\mu_k(t)\}_{k=1}^{\infty}$ be a linear independence, the $A_j = \sum_{k=j}^M \tilde{A}_k \beta_{kj}, k = 1, \dots, M$ then

$$A = \beta^T \tilde{A}, \quad (11)$$

using (8) leads to $Lg_M(t) = w(t)$, for $k = 1, \dots, M$,

$$\begin{aligned} &\rightarrow \sum_{h=1}^M \tilde{A}_h \langle L\tilde{\mu}_h, \tilde{\mu}_k \rangle_{W_2^i[t_0, t_1]} = \langle w, \tilde{\mu}_k \rangle_{W_2^i[t_0, t_1]} \\ &\rightarrow \sum_{h=1}^M \tilde{A}_h \sum_{j=1}^k \beta_{kj} \sum_{l=1}^h \beta_{hl} \langle L\mu_l, \mu_j \rangle_{W_2^i[t_0, t_1]} = \sum_{j=1}^k \beta_{kj} \langle w, \mu_j \rangle_{W_2^i[t_0, t_1]} \\ &\rightarrow \sum_{h=1}^M \tilde{A}_h \sum_{j=1}^j \sum_{l=1}^h \beta_{kj} \langle L\mu_l, \mu_j \rangle_{W_2^i[t_0, t_1]} \beta_{lh}^T = \sum_{j=1}^k \beta_{kj} \langle w, \mu_j \rangle_{W_2^i[t_0, t_1]} \\ &\rightarrow \sum_{h=1}^M \tilde{A}_h (\beta \Gamma \beta^T)_{kh} = \sum_{j=1}^j \beta_{kj} \langle w, \mu_j \rangle_{W_2^i[t_0, t_1]} \rightarrow (\beta \Gamma \beta^T) \tilde{A} = \beta \bar{w}, \end{aligned}$$

from (11) then $\Gamma A = \bar{w}$. ■

Theorem 3.3. [25, 28] If $g(t) \in W_2^i[t_0, t_1]$, then $g(t)$ is bounded.

Theorem 3.4. [25, 26, 28, 31] Let $\|g_M(t)\|_{W_2^i[t_0, t_1]}$ is bounded, and Eq. (1) has a unique solution. If the dense sequence $\{t_k\}_{k=1}^{\infty}$ on $[t_0, t_1]$, then M -term approximate solution $g_M(t)$ converges to $g(t)$ of (1) and the exact solution is expressed $g(t) = \sum_{k=1}^{\infty} A_k \mu_k(t)$, where $A_k = \sum_{j=1}^k \Gamma_{kj}^{-1} \bar{w}_j$.

Proof.

i. To prove that $g_M(t)$ convergent to $g(t)$, we conclude that:

$g_{M+1}(t) = g_M(t) + A_{M+1}\mu_{M+1}(t) = g_M(t) + \tilde{A}_{M+1}\tilde{\mu}_{M+1}(t)$,
 $\{\tilde{\mu}_k(t)\}_{k=1}^\infty$ is orthogonal, and leads to $\|g_{M+1}\|_{W_2^i[t_0, t_1]} = \|g_M\|_{W_2^i[t_0, t_1]} + \tilde{A}_k^2 = \sum_{k=1}^M \tilde{A}_k^2$,
 from the boundedness of $\|g_M\|_{W_2^i[t_0, t_1]}, \sum_{k=1}^M \tilde{A}_k^2 < \infty$, i.e. $\{\tilde{A}_k^2\}_{k=1}^\infty \in J^2$, where
 $J^2 = \{\tilde{A}_k; \sum_{k=1}^M \tilde{A}_k^2 < \infty\}, (k = 1, 2, \dots)$ if $N > M$ then

$$\begin{aligned} & \|g_N - g_M\|_{W_2^i[t_0, t_1]}^2 \\ &= \|g_N - g_{N-1} + g_{N-1} - \dots + g_{M+1} - g_M\|_{W_2^i[t_0, t_1]}^2 \\ &\leq \|g_N - g_{N-1}\|_{W_2^i[t_0, t_1]}^2 + \dots + \|g_{M+1} - g_M\|_{W_2^i[t_0, t_1]}^2 \\ &= \sum_{k=N}^{M+1} \|g_k - g_{k-1}\|_{W_2^i[t_0, t_1]}^2 \end{aligned}$$

So $\|g_k - g_{k-1}\|_{W_2^i[t_0, t_1]}^2 = \tilde{A}_k^2$, consequently $\|g_N - g_M\|_{W_2^i[t_0, t_1]}^2 = \sum_{k=1}^M \tilde{A}_k^2 \rightarrow 0$ as
 $M \rightarrow \infty$. Considering the completeness of $W_2^i[t_0, t_1]$,

$$g_M(t) \rightarrow g(t), \text{ as } M \rightarrow \infty$$

where $g(t) \in W_2^i[t_0, t_1]$.

ii. Can prove that $g(t)$ is a solution of (8). By (9)

$$g(t) = \sum_{k=1}^\infty A_k \mu_k(t) = \sum_{k=1}^\infty \tilde{A}_k \tilde{\mu}_k(t),$$

$$\text{Since } Lg(t_j) = \sum_{k=1}^\infty \tilde{A}_k < L\tilde{\mu}_k, \delta_j >_{W_2^i[t_0, t_1]}$$

$$= \sum_{k=1}^\infty \tilde{A}_k < \tilde{\mu}_k, L^* \delta_j >_{W_2^i[t_0, t_1]} = \sum_{k=1}^\infty \tilde{A}_k < \tilde{\mu}_k, \mu_j >_{W_2^i[t_0, t_1]},$$

$$\rightarrow \sum_{j=1}^M \beta_{Mj} * (Lg(t_j) = \sum_{k=1}^\infty \tilde{A}_k < \tilde{\mu}_k, \mu_j >_{W_2^i[t_0, t_1]})$$

$$\rightarrow \sum_{j=1}^M \beta_{Mj} Lg(t_j) = \sum_{k=1}^\infty \tilde{A}_k < \tilde{\mu}_k, \sum_{j=1}^M \beta_{Mj} \tilde{\mu}_j >_{W_2^i[t_0, t_1]}$$

$$= \sum_{k=1}^\infty \tilde{A}_k < \tilde{\mu}_k, \tilde{\mu}_M >_{W_2^i[t_0, t_1]} = \tilde{A}_M = \sum_{j=1}^M \beta_{Mj} w(t_j).$$

$$\text{if } M = 1 \text{ then } \beta_{11} Lg(t_1) = \beta_{11} w(t_0) \rightarrow Lg(t_1) = w(t_1),$$

$$\text{if } M = 2 \text{ then } \beta_{21} Lg(t_1) + \beta_{22} Lg(t_2) = \beta_{21} w(t_0) + \beta_{22} w(t_2) \rightarrow Lg(t_2) = w(t_2),$$

$$\text{moreover, it is clear } Lg(t_j) = w(t_j),$$

Since $\{t_k\}_{k=1}^\infty$ is dense on $[t_0, t_1]$, $\forall t \in [t_0, t_1], \exists \{t_{nk}\}_{k=1}^\infty$ such that $t_{nk} \rightarrow t$, as $k \rightarrow \infty$.
 Hence, when $k \rightarrow \infty$, by $Lg(t) = w(t_k)$, Obtained $g(t)$ is the solution of (8). ■

Theorem 3.5. [25, 26] If $g(t) \in W_2^i[t_0, t_1]$ then $e_M^2 = \|g_M(t) - g(t)\|_{W_2^i[t_0, t_1]}^2$ convergent to 0, and the sequence $\{e_M\}$ is monotonically decreasing.

Proof. From $g_M(t), g(t)$ in (9), (10) then

$$\begin{aligned} e_M^2 &= \|g_M - g\|_{W_2^i[a, b]}^2 = \|\sum_{k=M+1}^\infty A_k \mu_k(t)\|_{W_2^i[t_0, t_1]}^2 \\ &= \|\sum_{k=M+1}^\infty \tilde{A}_k \tilde{\mu}_k(t)\|_{W_2^i[t_0, t_1]}^2 = \sum_{k=M+1}^\infty \tilde{A}_k^2, \end{aligned}$$

and $e_{M-1}^2 = \sum_{k=M}^\infty \tilde{A}_k^2$, clearly, $e_M^2 \leq e_{M-1}^2$, and consequently $\{e_M\}$, is monotonically decreasing. ■

Algorithm

1. $t \in [t_0, t_1]$.
2. $t_k = \frac{(k-1)}{(M-1)}, k = 1, 2, \dots, M$.
3. $\mu_k(t) = L_s G_{t_k}^i(t)$.
4. $\Gamma_{kj} = \langle L\mu_j(t), \mu_k(t) \rangle_{W_2^i[t_0, t_1]}$.
5. $N=1$.
6. $g_0(t_1) = g(t_1)$.
7. $A_N = \sum_{j=1}^N \Gamma_{Nj}^{-1} \bar{w}_j$.
8. $g_N(t) = \sum_{k=1}^N A_k \mu_k(t)$.
9. If $N = M$ stop.
10. Else $N = N + 1$ go to (7) .

Numerical experiment

In this study, three examples of NIDEs have been solved to demonstrate the effectiveness of the RKHSM and its ability to converge toward the exact solution after a few number of iterations. Our error rate was also compared with other methods using absolute error as a measure.

The absolute error:

$$e_k = |g(t_k) - g_M(t_k)|, \quad k = 1, 2, \dots, M.$$

Example 1: Consider the Volterra NIDE [32]:

$$g'(t) + g(t) = w(t) + 2 \int_0^t \sin(t) g^2(z) dz, \quad t \in [0, 1] \quad (12)$$

where $w(t) = \cos(t)(1 + \sin^2(t)) + (1 - t)\sin(t)$, subject to initial condition $g(0) = 0$ and the exact solution is $g(t) = \sin(t)$.

Table 1. Comparing the numerical results of our method with [32]

t	$g(t)$	$g_M(t)$ in W_2^4	e_k	e_k in [32]
$2.0000E - 01$	$1.9866E - 01$	$1.9866E - 01$	0	$6.6518E - 06$
$4.0000E - 01$	$3.8941E - 01$	$0.3894E - 01$	$5.5622E - 14$	$2.2069E - 05$
$6.0000E - 01$	$5.6464E - 01$	$5.6464E - 01$	0	$4.0620E - 05$
$8.0000E - 01$	$7.1735E - 01$	$7.1735E - 01$	0	$2.5674E - 05$
$1.0000E - 00$	$8.4147E - 01$	$8.4147E - 01$	0	$9.6104E - 05$




Fig. 1 The comparisons between approximate and exact solutions for Example 1. at M=6.

Example 2: Consider the Fredholm NIDE [9]:

$$g'(t) + g(t) = w(t) - \int_0^1 g^2(z)dz , \quad t \in [0,1] \quad (13)$$

where $w(t) = \frac{1}{2}(e^{-2} - 1)$, subject to initial condition $g(0) = 1$, and the exact solution is $g(t) = e^{-t}$.

Table 2. Comparing the numerical results of our method with [9]

t	$g(t)$	$g_M(t)$ in W_2^5	e_k	e_k in [9]
$1.2500E - 01$	$8.8249E - 01$	$8.8249E - 01$	0	$6.2983E - 13$
$2.5000E - 01$	$7.7880E - 01$	$7.7880E - 01$	$1.1657E - 14$	$6.4948E - 13$
$3.7500E - 01$	$6.8728E - 01$	$6.8728E - 01$	0	$6.7090E - 13$
$5.0000E - 01$	$6.0653E - 01$	$6.0653E - 01$	0	$6.9033E - 13$
$6.2500E - 01$	$5.3526E - 01$	$5.3526E - 01$	0	$7.0754E - 13$
$7.5000E - 01$	$4.7236E - 01$	$4.7236E - 01$	$5.8287E - 15$	$7.2214E - 13$
$8.7500E - 01$	$4.1686E - 01$	$4.1686E - 01$	0	$7.3024E - 13$




Fig. 2 The comparisons between approximate and exact solution at $M=10$.

Example 3: Consider the Volterra NIDE [33]:

$$g'(t) = w(t) - \int_0^t (g^2(z) - 2)dz \quad t \in [0,1] \quad (14)$$

where $w(t) = \frac{1}{5}t^5$, the initial condition $g(0) = 0$, the exact solution is $g(t) = t^2$

Table 3. Comparing the numerical results of our method with [21]

t	$g(t)$	$g_M(t)$ in W_2^6	e_k	e_k in [21]
0	0	0	0	0
$2.0000E-01$	$0.4000E-01$	$0.4000E-01$	0	$4.6837E-17$
$4.0000E-01$	$1.6000E-01$	$1.6000E-01$	0	$2.7755E-17$
$6.0000E-01$	$3.6000E-01$	$3.6000E-01$	$1.1657e-14$	$1.11022E-16$
$8.0000E-01$	$6.4000E-01$	$6.4000E-01$	0	$2.2204E-16$
$1.0000E-00$	$1.000E-00$	$1.000E-00$	0	$1.1102E-15$




Fig. 3 The comparisons between approximate and exact solutions at $M=4$.

5. Conclusion

In this study, the reproducing kernel Hilbert space method was applied on three examples of nonlinear integro- differential equations that have analytical solutions for the purpose of knowing the accuracy and effectiveness of proposed method, using MATLAB R2023a to obtain a numerical result. A new algorithm was used instead of the Gram-Schmidt algorithm, which was It is numerically unstable and contains many complicated mathematical operations. A comparison was also made between the resulting approximate solutions and the approximate solutions of other methods. The results showed that the absolute error in the solutions is equal to zero or converges to zero, which means that it is better than the results of [32], [9] and [33] at an equal number of iterations. The algorithm is simpler, more efficient and contains fewer mathematical steps, leading to more accurate approximate solutions compared to other methods.

References

- [1] F. Bloom, "Asymptotic bounds for solutions to a system of damped integro-differential equations of electromagnetic theory," *Journal of Mathematical Analysis and Applications*, vol. 73, pp. 524–542, 1980. Doi: [https://doi.org/10.1016/0022-247X\(80\)90297-8](https://doi.org/10.1016/0022-247X(80)90297-8).
- [2] K. Holm aker, "Global asymptotic stability for a stationary solution of a system of integro-differential equations describing the formation of liver zones," *SIAM Journal on Mathematical Analysis*, vol. 24, pp. 116–128, 1993. Doi: <https://doi.org/10.1137/S0036139995283110>.
- [3] L. K. Forbes, S. Crozier and D. M. Doddrell, "Calculating current densities and fields produced by shielded magnetic resonance imaging probes," *SIAM Journal on Applied Mathematics*, vol. 57, pp. 401–425, 1997. Doi: <https://doi.org/10.1137/0524008>.
- [4] X. Lv and S. Shi, "The combined RKM and ADM for solving nonlinear weakly singular Volterra integro-differential equations," *Abstract and Applied Analysis*, vol. 2012, pp.1-10, 2012. Doi: <https://doi.org/10.1155/2012/258067>.

- [5] R. H. Khan and H. O. Bakodah, "Adomian decomposition method and its modification for nonlinear Abel's integral equations," *Computers and Mathematics with Applications*, vol. 7, pp. 2349 – 2358, 2013. Doi: <http://dx.doi.org/10.12988/ijma.2013.37179>.
- [6] G. Ajileye and S. A. Amoo, "Numerical solution to Volterra integro-differential equations using collocation approximation," *Mathematics and Computational Sciences*, vol. 4, pp. 1–8, 2023. Doi: <http://doi.org/10.30511/mcs.2023.1978083.1099>.
- [7] O. A. Agbolade and T. A. Anake, "Solutions of first-order Volterra type linear integro-differential equations by collocation method," *Journal of Applied Mathematics*, vol. 2017, pp. 1-5, 2017. Doi: <https://doi.org/10.1155/2017/1510267>.
- [8] S. Nemati, P. Lima and Y. Ordokhani, "Numerical method for the mixed Volterra-Fredholm integral equations using hybrid Legendre functions," *Application of Mathematics* 2015, pp. 184–193, 2015. Available: <http://dml.cz/dmlcz/702975>.
- [9] B. Durak, A. Sezin, H.Ö. ÖZER, L. E. Sakman and Ş. Kapkin, "Collocation method applied to numerical solution of integro-differential equations," *Kahramanmaraş Sütçü İmam Üniversitesi Mühendislik Bilimleri Dergisi*, vol. 26, pp. 1010–1020, 2022.
- [10] K. Maleknejad and Y. Mahmoudi, "Taylor polynomial solution of high-order nonlinear Volterra–Fredholm integro-differential equations," *Applied Mathematics and Computation*, vol. 145, p. 641–653, 2003. Doi: [https://doi.org/10.1016/S0096-3003\(03\)00152-8](https://doi.org/10.1016/S0096-3003(03)00152-8).
- [11] H. O. Al-Humedi and Z. A. Jameel, "Cubic B-spline least-square method combine with a quadratic weight function for solving integro-differential equations," *Earthline Journal of Mathematical Sciences*, pp. 99–113, 2020. Doi: <https://doi.org/10.34198/ejms.4120.99113>.
- [12] H. O. Al-Humedi and Z. A. Jameel, "Combining Cubic B-spline Galerkin method with quadratic weight function for solving partial integro-differential equations," *Journal of Al-Qadisiyah for Computer Science and Mathematics*, vol. 12, pp. 9-20, 2020. Doi: <https://doi.org/10.29304/jqcm.2020.12.1.660>.
- [13] H. O. Al-Humedi and A. F. Shoushan, "A combination of the orthogonal polynomials with least-squares method for solving high-orders Fredholm-Volterra integro-differential equations," *Al-Qadisiyah Journal of pure science*, vol. 26, pp. 20–38, 2021. Doi: <https://doi.org/10.29350/jops.2021.26.1.1207>.
- [14] A. F. Shoushan and H. O. Al-Humedi, "The numerical solutions of integro-differential equations by Euler polynomials with least-squares method," *PalArch's Journal of Archaeology of Egypt/Egyptology*, vol. 18, pp. 1740–1753, 2021.
- [15] J. Biazar, H. Ghazvini and M. Eslami, "He's homotopy perturbation method for systems of integro-differential equations," *Chaos, Solitons & Fractals*, vol. 39, pp. 1253–1258, 2009. Doi: <https://doi.org/10.1016/j.chaos.2007.06.001>.
- [16] J. Hou, J. Niu and W. Ngolo, "A new numerical method to solve nonlinear Volterra-Fredholm Integro-differential equation," *Mathematical Modelling and Analysis*, vol. 26, pp. 469–478, 2021. Doi: <https://doi.org/10.3846/mma.2021.12923>.
- [17] H. O. Bakodah, M. Al-Mazmumy, S. O. Almuhalbedi and L. Abdullah, "Laplace discrete Adomian decomposition method for solving nonlinear integro differential equations," *Journal of Applied Mathematics and Physics*, vol. 7, pp. 1388–1407, 2019. Doi: <https://doi.org/10.4236/jamp.2019.76093>.
- [18] H. Du and J. Shen, "Reproducing kernel method of solving singular integral equation with cosecant kernel," *Journal of Mathematical Analysis and Applications*, vol. 348, pp. 308–314, 2008. Doi: <http://dx.doi.org/10.1016/j.jmaa.2008.07.037>.

- [19] Z. Chen and W. Jiang, "The exact solution of a class of Volterra integral equation with weakly singular kernel," *Applied Mathematics and Computation*, vol. 217, pp. 7515–7519, 2011. Doi: <http://dx.doi.org/10.1016/j.amc.2011.02.059>.
- [20] H. Beyrami, T. Lotfi and K. Mahdiani, "Stability and error analysis of the reproducing kernel Hilbert space method for the solution of weakly singular Volterra integral equation on graded mesh," *Applied Numerical Mathematics*, vol. 120, pp. 197–214, 2017. Doi: <https://doi.org/10.1016/j.apnum.2017.05.010>.
- [21] H. O. Al-Humedi and A. S. Abdul-hasan, "The reproducing kernel Hilbert space method for solving system of linear weakly singular Volterra integral equations," *J. Adv. Math.*, vol. 15, pp. 8070–8080, 2018.
- [22] L.-H. Yang, H.-Y. Li and J.-R. Wang, "Solving a system of linear Volterra integral equations using the modified reproducing kernel method," in *Abstract and Applied Analysis* voll. 2013, pp.1-5, 2013. Doi: <http://dx.doi.org/10.1155/2013/196308>.
- [23] X. Lv and Y. Gao, "Reproducing kernel space method for the solution of linear Fredholm integro-differential equations and analysis of stability," *Abstract and Applied Analysis*, vol. 2012, pp.1-10, 2012. Doi: <http://dx.doi.org/10.1155/2012/971593>.
- [24] A. Alvandi and M. Paripour, "Reproducing kernel method with Taylor expansion for linear Volterra integro-differential equations," *Communications in Numerical Analysis*, vol. 1, pp. 1–10, 2017. Doi: <http://dx.doi.org/10.5899/2017/cna-00264>.
- [25] O. Abu Arqub, M. Al-Smadi and S. Momani, "Application of reproducing kernel method for solving nonlinear Fredholm-Volterra integro-differential equations," *Abstract and Applied Analysis*, vol. 2012, pp.1-16, 2012. Doi: <http://dx.doi.org/doi:10.1155/2012/839836>.
- [26] O. A. Arqub, M. Al-Smadi and N. Shawagfeh, "Solving Fredholm integro-differential equations using reproducing kernel Hilbert space method," *Applied Mathematics and Computation*, vol. 219, pp. 8938–8948, 2013. Doi: <http://dx.doi.org/10.1016/j.amc.2013.03.006>.
- [27] A. Alvandi and M. Paripour, "The combined reproducing kernel method and Taylor series for solving nonlinear Volterra-Fredholm integro-differential equations," *International Journal of Mathematical Modelling & Computations*, vol. 6, pp. 301–312, 2016. Doi: <https://doi.org/10.1155/2017/7498136>.
- [28] S. Farzaneh Javan, S. Abbasbandy and M. A. Fariborzi Araghi, "Application of Reproducing Kernel Hilbert Space Method for Solving a Class of Nonlinear Integral Equations," *Mathematical Problems in Engineering*, vol. 2017, pp. 1–10, 2017. Doi: <https://doi.org/10.1155/2017/7498136>.
- [29] S. Javadi, E. Babolian and E. Moradi, "New implementation of reproducing kernel Hilbert space method for solving a class of functional integral equations," *Communications in Numerical Analysis*, vol. 2014, pp. 1–7, 2014. Doi: <https://doi.org/10.5899/2014/cna-00205>.
- [30] E. Moradi, A. Yusefi, A. Abdollahzadeh and E. Tila, "New implementation of reproducing kernel Hilbert space method for solving A class of third-order differential equations," *Journal of Mathematics and Computer Science*, vol. 12, pp. 253–262, October 2014.
- [31] M. Cui, *Nonlinear Numerical Analysis in Reproducing Kernel Space.*, Y. Lin, Ed., New York: Nova Science Publishers, Incorporated, 2009.
- [32] G. Ajileye, O. O. Aduroja, I. G. Amakoromo and S. A. Amoo, "A numerical approach to the solution of nonlinear Volterra-Fredholm integro-differential

equations," Songklanakarin Journal of Science & Technology, vol. 46, pp.8-15, 2024.

- [33] J. Hou, J. Niu and W. Ngolo, "A new numerical method to solve nonlinear Volterra-Fredholm integro-differential equations," Mathematical Modelling and Analysis, vol. 26, pp. 469–478, 2021. Doi: <https://doi.org/10.3846/mma.2021.12923>.

طريقة استنساخ نواة فضاء هيلبرت لحل المعادلات التفاضلية التكاملية غير الخطية

وفاء كامل الزهيري* ، حميدة عودة الحميدي

قسم الرياضيات، كلية التربية للعلوم الصرفة، جامعة البصرة، بصرة، العراق.

معلومات البحث	الملخص
الاستلام 9 آب 2024 المراجعة 20 ايلول 2024 القبول 25 ايلول 2024 النشر 31 كانون الأول 2024	في هذا العمل تم استخدام طريقة استنساخ نواة فضاء هيلبرت لايجاد حلول التقريبية للمعادلات التفاضلية التكاملية غير الخطية على شكل متسلسلة منتهية، وقد أظهرت النتائج أن الحلول التقريبية يقترَب بشكل وثيق من الحلول الدقيق للمعادلات، مما يؤكد فعالية الطريقة كأداة موثوقة وعملية لحل هذا النوع من المعادلات. هذه الطريقة ذات أهمية كبيرة في مجال ايجاد الحلول العددية للمعادلات التفاضلية التكاملية غير الخطية، والتي لها تطبيقات واسعة في العديد من المجالات الهندسية والعلمية.
الكلمات المفتاحية	طريقة استنساخ نواة الفضاء هيلبرت ، المعادلات التفاضلية التكاملية غير الخطية ، الحل التقريبي ، فضاء هيلبرت ، فضاء الدوال.

Citation: W. K. Al-Zuhairi, H. O. Al-Humedi, J. Basrah Res. (Sci.) 50(2), 182 (2024).
[DOI:https://doi.org/10.56714/bjrs.50.2.15](https://doi.org/10.56714/bjrs.50.2.15)

*Corresponding author email : pgs.Wafaa.kamel@uobasrah.edu.iq

