# **ARTICLE / INVESTIGACIÓN**

# Synthesis, characterization, biological studies and DFT study of Schiff Bases and their complexes derived from aromatic diamine compounds with cobalt (II)

Abduljleel Mohammed Abduljleel, Jassim Mohammed Saleh Alshawi, kawkab Ali Hussein\*, Sadiq M-H. Ismael

DOI. 10.21931/RB/2023.08.01.61 Department of Chemistry, College of Education for Pure Sciences, University of Basrah, Basrah, 61004 Iraq. Corresponding author: kawkab.ali@uobasrah.edu.ig.

**Abstract:** Based on vanillin and diamine compounds (ortho phenylene diamine, meta phenylene diamine, 3,4- diamine toluene), derivation of two new Schiff base ligands (L<sup>1</sup> and L<sup>2</sup>) was done, after which synthesis and treatment with Co (II) chloride was performed at a metal-to-ligand ratio of 1:1 to get two new complexes, i.e.  $[CoL^3(H_2O)_2]Cl_2$  and  $[CoL^1(H_2O)_2]Cl_2$ . These complexes and ligands were characterized by employing NMR, IR, atomic absorption, UV visible absorption, molecular weight determination, molar conductance, and magnetic measurement techniques. As per the data, the ligands were found to be bidentate ligands that were linked to two azomethine nitrogen sites. It was suggested that these complexes were paramagnetic electrolyte compounds that possessed coordination number four. Screening of the ligands and metal complexes was done to assess their antimicrobial activities against gram-negative and gram-positive bacteria, which was found to show biological activity. Calculations using quantum chemistry were done to examine the molecule geometry. The investigation includes several quantum chemical characteristics derived from frontier molecular orbitals.

Key words: Schiff bases, transition metal complexes, vanillin, diamine aromatic compounds, antibacterial activity, DFT study.

#### Introduction

The general structure of Schiff bases includes R-CH=N-Ar, wherein Ar and R represent the aromatic and aliphatic groups due to the condensation of primary amines with ketones or aldehydes<sup>1</sup>. These can be characterized based on the (-N=CH-) groups, which include biological characteristics like antimicrobial<sup>2</sup>, antifungal<sup>3</sup> and anticancer activities<sup>4</sup>. Schiff bases derived based on diamines and aldehydes have one of the most critical synthetic ligand systems for asymmetric catalysis. They are considered significant for a wide variety of progress metal catalysis responses, including epoxidation of ole-fins<sup>5</sup>, lactide polymerisation<sup>6</sup>, and the opening of the asymmetric rings of epoxides7. The chelating Schiff base ligands derived based on different carbonyl compounds, and diamines include a highly notable class of compounds with numerous applications in synthetic<sup>8</sup>, catalytic clinical<sup>9</sup>, analytical<sup>10</sup>, and biochemical<sup>11</sup> processes. Previous studies have shown that enhancement of the coordinating potential of diamine compounds like meta phenylene diamine happens when they are condensed with different carbonyl compounds. However, the literature review did not show any studies or research work conducted regarding the transition metal complexes of Schiff bases that have been derived from vanillin and certain diamines. The current research focuses on preparing Schiff bases based on the condensation of vanillin along with certain aromatic diamines and synthesizing their complexes with cobalt (II) ions. These Schiff bases and complexes have been identified based on 1HNMR, 13CNMR, UV/visible and IR, and atomic absorption techniques. This focus also examined these complexes' conductivity, antibacterial activities, and magnetic characteristics. DFT calculations upheld the research findings. The ground state characteristics of the ligand and its complexes were studied by DFT at the B3LYP/6-31+G (d,p) and B3LYP/LANL2DZ, respectively.

### **Materials and methods**

Merck provided all the employed reagents, which have been used without purification. The melting point was determined based on a Thermo Fisher apparatus. An FTIR – 8400 S – Shimadzu spectrometer was employed to record the IR spectra in the 400–4000 cm<sup>-1</sup> via KBr pellets. An NMR spectrophotometer (Brucker – 500 MHZ) was used to record the 1HNMR spectra of CDCl<sub>3</sub>, while a Jenway Pcmb conductivity meter was used to measure the conductance in DMF at room temperature. An SPV-725 (Germany) spectrophotometer was used to record the UV visible spectra in THF. An atomic absorption apparatus (Analyst 200 Atomic Absorption Spectrometer) determined the metal-to-ligand molar ratio. The magnetic properties were measured at room temperature by employing Hg [Co (NCS)4] as the calibrant based on the Gouy method.

## **Results**

#### Synthesis of Schiff Base Ligands add L<sup>1</sup> and L<sup>2</sup>

Vanillin (3.041 gm and 20 mmol) was mixed with diamine (10 mmol) in ethanol (50 mL), and a few drops of glacial acetic acid were added. Then reflux mixture for 14 - 20 hrs. According to the TLC examination, the starting materials

**Citation:** Abduljleel M A, Jassim M S A, kawkab A H and Sadiq M-H I, Synthesis, Characterization and Biological Studies of Schiff Bases and Their Complexes Derived from Aromatic Diamine Compounds with Cobalt (II). Revis Bionatura 2023;8 (1) 61. http://dx.doi. org/10.21931/RB/2023.08.01.61

Received: 15 January 2023 / Accepted: 25 February 2023 / Published: 15 March 2023

Publisher's Note: Bionatura stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1