

<https://doi.org/10.34883/PI.2025.15.1.046>



Abbas Abed Khazaal<sup>1</sup> ✉, Miami Kadhim Yousif<sup>2</sup>, Hasnah Abdulzahra Mousa<sup>3</sup>

<sup>1</sup> Basrah Medical College, University of Basrah, Basrah, Iraq

<sup>2</sup> Al-Zahraa Medical College, University of Basrah, Basrah, Iraq

<sup>3</sup> Al-Mawani Teaching Hospital, Basra Health Directorate, Basrah, Iraq

## Mothers' Knowledge and Practice on Optimal Complementary Feeding

**Conflict of interest:** nothing to declare.

**Authors' contribution:** Abbas Abed Khazaal – conceptualization, data curation, investigation, methodology, project administration, validation, visualization, writing – original draft and writing – review & editing; Miami Kadhim Yousif – conceptualization, data curation, investigation, methodology, resources, software, supervision, validation, visualization, and writing – review & editing; Hasnah Abdulzahra Mousa – conceptualization, data curation, investigation, methodology, software, validation, visualization, writing – original draft and writing – review & editing.

The article is published in author's edition.

Submitted: 20.01.2025

Accepted: 14.02.2025

Contacts: [abbas.hatab@uobasrah.edu.iq](mailto:abbas.hatab@uobasrah.edu.iq)

### Abstract

**Introduction.** Optimal complementary feeding during first two year of life is crucial for optimal development and growth of infant and young children during first two years of life.

**Purpose.** The goal of this study is to investigate mothers' knowledge and practices about the complementary feeding among infants and young children age from four months to two years of life.

**Materials and methods.** A cross-sectional study was designed to interview 210 mothers of infant and young children, aged from 4 month to two year who attended outpatient pediatrics clinic in Al-Mawani Teaching Hospital in Basrah, the relation between mothers' age and educational level, with their knowledge and practices about complementary feeding was evaluated.

**Results.** The results revealed only 28.57% of the participated mothers know the proper time of starting complementary food, and about 52% of the mothers start complementary food before six months of age. We didn't found statistical significant relation between mothers' age, mothers' education level and mothers' knowledge and practices, apart from a significant relation between level of mothers' education level and correct frequency of introduction new food per week (p-value 0.001).

**Conclusion.** There is a need to promote optimal feeding practices among mothers to insure optimal complementary feeding.

**Keywords:** complementary feeding, feeding practices, passive immunity, nutritional requirements, infants

## ■ INTRODUCTION

The process of giving of appropriate food at the proper time in addition to mothers' breast milk to face increased needing nutrients for the growing child is called complementary feeding. The exclusive breast feeding of the babies during the first six

months of life endows the baby a passive immunity that is essential to protect them from various bacterial and viral infections with the lack of a well-developed immune system at this critical period of time. In addition to that, the infants and young children who are provided with complementary food after the first half year will meet the rising nutritional requirements necessary for development and growth. This process called weaning, and mothers who do weaning should be knowledgeable about the timing and types of complementary feeding [1, 2].

In addition to that first two years of life considers as a critical period of time for development and growth of children. Most cases of malnutrition occur during this time, thus appropriate nourishment is critical. Malnutrition in the first two years of life can cause permanent impairment to physical and intellectual development. As a result, improving food habits during this time is critical to improve children's health and nutrition levels [3, 4].

Suboptimal complementary feeding practices are associated with a large burden of malnutrition and mortality in children [5].

Mother's information and work on with respect to reciprocal food play a significant part on their taking care of practices. The capacity of the mothers to do the suggested reciprocal taking care of practices is related with their insight and demeanor on ideal integral taking care of. Numerous variables could influence mothers' information and disposition towards complementary feeding [6–8].

The goal of our study is to determine the knowledge and practices of the mothers about complementary feeding and to identify the factors like mothers' education and mothers' age which could be affecting their knowledge and practice of complementary food.

## ■ MATERIALS AND METHODS

### **Study design**

A cross-sectional descriptive study was carried out among mothers who visited the outpatient department of Al-Mawani Teaching Hospital in Basrah, over the period of four months, from January 2023 to April 2023.

### **Sample size**

210 mothers of up to two years children were included in this study aged (17–44 years) who visited the pediatrics outpatient clinic in Al-Mawani Teaching Hospital in Basrah during the study period.

### **Exclusion criteria**

Mothers of infants aged less 4 months or more than 24 months were excluded.

### **Inclusion criteria**

Mothers of infants and young children between 4–24 months of age visiting the outpatient clinic of pediatrics were involved in this study.

### **Data collection**

The study is based on interviews using questionnaire, data were included information about age of starting complementary food, type of food start with, causes of delay

complementary feeding, correct frequency of introducing new foods per week, and the way of feeding whether by the mother or baby-led which means babies are allowed to feed themselves.

Other variables addressed were the age of the mother, educational level and occupation of mother, family size, child age and sex and feeding pattern of child whether breast, bottle or mixed feeding.

### **Ethical approval**

The study protocol received approval from the College Council and Ethical Committee at Basrah College of Medicine, University of Basrah, with reference number 030411-036-2022 on 3/4/2022.

### **Data analysis**

The data collected was analysed using SPSS version 23. Chi-square test was used and a  $P < 0.05$  is considered significant.

## **■ RESULTS**

210 mothers were involved in this study, majority of the mothers were belong to age group 20–40 years (88.57%) and majority of the mothers were housewives (92.85%), and about 74% of the mothers had primary and secondary school education. 15.7% of the mothers were illiterate and only 10% have college and above education. About 57% of children included in this study were male, and about 43% of them were female. Regarding the age of children 39.05% of them belong to age group 6–12 month, and 7.62% aged less than 6 months and 25.71% belonged to age group 12–18 and 27.62 belonged to age group 18–24 month. About tow third of children (66.67%) are bottle fed, were the remaining children were breast and mixed feeding with 23.81% and 9.52% respectively (Table 1).

Age of starting of complementary feeding was less than 6 months in 52.38% of children, and at age of 6 months of age in 28.57% of children and older than 6 months in about 19.05% of studied children. Way of giving complementary feeding was by mother in 91.43% of children and by baby-led in only 8.57% of studied children. Correct frequency of introduction new food per week was found in only 29.5% of children while about 70.5% of children had incorrect frequency of introduction new food per week. Frequency of complementary feeding was twice a day in about 41.9% of children and thrice a day in 58.1 of children. There was no significant correlation between mothers' age and age of initiation of complementary feeding, way of giving complementary feeding, and correct frequency of introduction new food per week,  $p$ -value is 0.635, 0.969, 0.789 respectively. There was a significant correlation between mothers' education level and correct frequency of introduction new food per week ( $p=0.001$ ). There was no significant correlation between mothers' education level and the way of giving complementary feeding, age of initiation of complementary feeding  $p$ -value was 0.09 and 0.747 respectively (Table 2).

**Table 1**  
**General and sociodemographic characters of the mothers and their children**

Character	No.	%
Mothers' Age (year)		
<20	14	6.67
20–40	186	88.57
>40	10	4.76
Mothers' Education Status		
Illiterate	33	15.71
Primary	84	40.00
Secondary	72	34.29
College and above	21	10
Mothers' Occupation		
Housewife	195	92.85
Government Employee	14	6.67
Private Employee	1	0.48
Student	0	0
Sex of the child		
Male	120	57.14
Female	90	42.86
Childs' Age(month)		
<6 m	16	7.62
6–12	82	39.05
12–18	54	25.71
18–24	58	27.62
Childs' feeding pattern		
Breast	50	23.81
Bottle	140	66.67
Mixed	20	9.52

**Table 2**  
**Mothers' Knowledge and their practice about complementary feeding**

Variable	No.	%
Age of initiation of complementary feeding		
<6 m	110	52.38
At 6 m	60	28.57
>6 m	40	19.05
Way of giving complementary feeding		
By Mother	192	91.43
Baby – Led	18	8.57
Correct Frequency of introduction new food per week		
Yes	62	29.52
No	148	70.48
Frequency of complementary feeding		
Twice a day	88	41.90
Thrice a day	122	58.10

## ■ DISCUSSION

Complementary feeding plays a crucial role for healthy growth and development of infants and young children during the first two years of life [9].

According to World Health Organization guidelines, complementary feeding should begin at six months of age, and breastfeeding should continue until two years of age [10, 11]. Delayed beginning of complementary foods may enhance the risk of

malnutrition, immune diseases, and development of diabetes mellitus in later life in high-risk population [12].

In this study we investigated mothers' knowledge and practices toward optimal complementary feeding of children from the age 4–24 months in Basrah.

The study shows that the most of participated mothers were young housewives and had low education level, these results are comparable with other studies was done in Baghdad [13], Erbil [14], and Al-Najaf [15].

In this study 28.57% of mothers know the correct age of introduction of complementary feeding, which is lower compering with other regions universally [16].

In this study (52.38%) of the participated mothers start complementary feeding before age of 6 months. Many mothers start complementary feeding before the recommended age of 6 months [17] these no add benefits from this practice [18] this may also increase risk of infants' alimentary infections and may increase the incidence of childhood obesity in the future [19].

The majority of the involving mothers (70.48%) didn't had an idea about the correct frequency of introduction new food per week, this may be due to low educational level of the participated mothers.

Regarding the way of giving complementary feeding, we found that the majority of the babies (91.42%) was fed by the mother and only (8.58%) by baby-led method.

In our study there was no significant relation between mothers' age and age of initiation of complementary feeding, way of giving complementary feeding, and correct frequency of introduction new food per week. Also, there was no significant relation between mothers' education level and the age of initiation of complementary feeding, way of giving complementary feeding, apart from a significant relation between level of mothers' education and correct frequency of introduction new food per week, similar results were obtained in study done in Kosova [20].

There were some limitations of our study is that cross-sectional design and small-sized sample which collected from single site of Basrah, in spite of that this study can be used as a foundation for future researches.

## ■ CONCLUSION

Optimal complementary feeding during first two year of play a crucial role in healthy development and growth of infant and young children. Many factors influence mothers' understanding and practices on complementary feeding, including their age, educational level, and family socioeconomic status. There is a need to promote optimal feeding knowledge and practices among mothers to insure ideal complementary feeding.

## ■ REFERENCES

1. Nuzzi G, Gerini C, Comberiati P, Peroni DG. The weaning practices: a new challenge for pediatricians? *Pediatr Allergy Immunol*. 2022;33:44–46.
2. Verga MC, Scotese I, Bergamini M, et al. Timing of complementary feeding, growth, and risk of non-communicable diseases: systematic review and meta-analysis. *Nutrients*. 2022;14:702.
3. Indicators for assessing infant and young child feeding practices. Conclusions of a Consensus Meeting Held 6–8 November 2007 in Washington DC. 2007:1–19.
4. World Health Organization. Indicators for Assessing Infant and Young Child Feeding Practices. World Health Organization; 2010. Available from: <https://apps.who.int/iris/bitstream/10665/42590/1/9241562218.pdf>. Accessed November 20, 2019.
5. WHO Guideline. Improving childhood development-summary. 2019. Available from: [www.who.int/maternal\\_child\\_adolescent/child/summary\\_guideline\\_improving\\_early\\_childhood\\_development.pdf](http://www.who.int/maternal_child_adolescent/child/summary_guideline_improving_early_childhood_development.pdf). Accessed January 02, 2020.

6. Egata G, Berhane Y, Worku A. Predictors of non-exclusive breastfeeding at 6 months among rural mothers in east Ethiopia: a community-based analytical cross-sectional study. *Int Breastfeed J*. 2013;8:8.
7. Gyampoh S, Otoo GE, Aryeetey RN. Child feeding knowledge and practices among women participating in growth monitoring and promotion in Accra, Ghana. *BMC Pregnancy Childbirth*. 2014;14(1):180.
8. Fadare O, Id MA, Mavrotas G, et al. Mother's nutrition-related knowledge and child nutrition outcomes: empirical evidence from Nigeria. *PLoS One*. 2019;14:e0212775.
9. Dewey KG. The challenge of meeting nutrient needs of infants and young children during the period of complementary feeding: An evolutionary perspective. *J Nutr*. 2013;143(12):2050–4.
10. World Health Organization. Global strategy for infant and young child feeding [Internet]. Geneva: World Health Organization. 2003 [cited January 23, 2019]. Available from: <https://www.who.int/nutrition/publications/infantfeeding/9241562218/en/>
11. World Health Organization. Complementary feeding of young children in developing countries: a review of current scientific knowledge. Geneva: World Health Organization. 1998 Available from: [https://www.who.int/nutrition/publications/infantfeeding/WHO\\_NUT\\_98.1/en/](https://www.who.int/nutrition/publications/infantfeeding/WHO_NUT_98.1/en/)
12. Przyrembel H. Timing of introduction of complementary food: short- and long-term health consequences. *Ann Nutr Metab*. 2012;60 Suppl 2:8–20.
13. Abdul Ameer AJ, Al-Hadi AH, Abdulla MM. Knowledge, attitudes and practices of Iraqi mothers and family child-caring women regarding breastfeeding. *East Mediterr Health J*. 2008;14(5):1003–14.
14. AL-Azzawi SI, Hussein KA, Shaker NZ. Knowledge, Attitude and Practices (KAP) of Mothers toward Infant and Young Child Feeding in Primary Health Care (PHC) Centers, Erbil City. *Kufa Journal for Nursing sciences*. 2012;2(2):118–126.
15. AL-Abedi NFH, Al-Asadi KMN. Assessment of mother's knowledge toward breastfeeding at AL-Najaf City. *International Journal of Scientific and Research Publications*. 2016;6(12):31–38.
16. UNICEF's Report. From the First Hour of Life, Making the case for improved infant and young child feeding everywhere, 2016. Available from: <https://data.unicef.org/resources/firsthour-life-new-report-breastfeeding-practices/>.
17. Alder EM, et al. What influences the timing of the introduction of solid food to infants? *Br J Nutr*. 2004;92(3):527–31.
18. Dewey KG. Nutrition, growth, and complementary feeding of the breastfed infant. *Pediatr Clin North Am*. 2001;48(1):87–104.
19. Pearce J, Taylor MA, Langley-Evans SC. Timing of the introduction of complementary feeding and risk of childhood obesity: a systematic review. *Int J Obes (Lond)*. 2013;37(10):1295–306.
20. Merita Berisha Naser Ramadani, Rina Hoxha1, Sanije Gashi, et al. Knowledge, Attitudes and Practices of Mothers in Kosova About Complementary Feeding for Infant and Children 6–24 Months. *Med Arch*. 2017;71(1):37–41.