



Analysis and evaluation of toxic heavy metals in sediments of seabed cores using inductively coupled plasma optical emission spectroscopy

Muhand K. AL-Tememi ^a, Ahmed Y. Hammood^a, Zuhair A. Abdulnabi^a, and Ahmed Abdalhadi Majed ^{b,c,*}

^a Marine Science Center, University of Basrah, zip code is 61004, Basrah, Iraq

^b College of Education for Pure Science, University of Basrah, Iraq

^c Scientific Research Center, Al-Ayen University, Thi-Qar, Iraq

ARTICLE INFO:

Received 3 May 2025

Revised form 23 Jul 2025

Accepted 19 Aug 2025

Available online 30 Sep 2025

Keywords:

Heavy metals,
Analysis of sediments
Inductively coupled plasma optical
emission spectroscopy,
Seabed,
Geoaccumulation Index

ABSTRACT

Soil sediments were studied in three areas in the northern Gulf within Iraqi territorial waters, where sediments were collected to cover all depths from 0 to 40 meters below the sea floor. The concentrations of the elements lead (Pb), cadmium (Cd), nickel (Ni), chromium (Cr), and vanadium (V) were measured using the inductively coupled plasma optical emission spectroscopy (ICP-OES), and arsenic (As) was determined by the graphite furnace atomic absorption spectrometry (GF-AAS) (%RSD < 5). The detection limits of Pb, As, Cd, Ni, Cr, and V were obtained at 0.511, 0.554, 0.525, 0.528, 0.501, and 1.221 $\mu\text{g L}^{-1}$, respectively. The Geoaccumulation Index (I-geo) compares the concentration of heavy metals in six classes in a sediment sample to its background value. The results were compared using I-geo calculations, which indicated high levels of contamination for arsenic and cadmium, with I-geo values ranging from 2.5 to 6.7 for arsenic, with moderately to significantly contaminated (Class 3 and 6), and from 2.4 to 3.9 for cadmium, with moderately to heavily contaminated (Class 3 and 4). There were lower contamination rates for nickel, as the I-geo values ranged from 0.23 to 2.09, while the concentrations of lead, chromium, and vanadium remained within safe levels. So, no dangerous concentrations for Pb, Ni, Cr, and V were found in any areas or depths from 0 to 40 meters below the seafloor.

1. Introduction

Heavy metals are among the key components of the Earth's crust. The level of heavy metals in soil varies widely depending on geological formation, depth, and location [1]. These elements can have a significant impact on the environment, economy, and human health when present in soil. Therefore, it is essential to understand the levels and concentrations of these minerals in sediment,

especially in areas where human activities, such as commercial, industrial, agricultural, and marine fishing, are closely linked to the sediment [2]. The Earth's crust contains about 15 $\mu\text{g g}^{-1}$ of lead. Lead levels in global soils vary from 2 to 200 $\mu\text{g g}^{-1}$ on average. Lead is also present in many rocks, including argillaceous and magmatic types, where it can be found in significant amounts (10–40 $\mu\text{g g}^{-1}$), as well as in ultramafic rocks (0.1–10 $\mu\text{g g}^{-1}$) [3]. Cadmium is regarded as a rare element. Although its approximate abundance in the earth's crust is 0.1 $\mu\text{g g}^{-1}$, it is not found in nature as a

*Corresponding Author: [Ahmed Abdalhadi Majed](mailto:Ahmed.Abdalhadi.Majed@uobasrah.edu.iq)

Email: eduppg.ahmed.majed@uobasrah.edu.iq

<https://doi.org/10.24200/amecj.v8.i03.1047>