

PAPER • OPEN ACCESS

Curdlan Gum, Properties, Benefits and Applications

To cite this article: Y. S. S. Al-Rmedh *et al* 2023 *IOP Conf. Ser.: Earth Environ. Sci.* **1158** 112011

View the [article online](#) for updates and enhancements.

You may also like

- [Investigating SUMER Coronal Hole Observations: A Robust Method of Raster Reduction](#)
Alisdair R. Davey, Scott W. McIntosh and Donald M. Hassler
- [Biological Activity of Quinolinium Salts](#)
B M Gutsulyak
- [Preface](#)

Curdlan Gum, Properties, Benefits and Applications

Y. S. S. Al-Rmedh¹, H. I. Ali² and S. T. G. Al-Sahlany³

¹⁻³Department of Food Science, College of Agriculture, University of Basrah, Basrah, Iraq.

²E-mail: Haider.ali@uobasrah.edu.iq

Abstract. Curdlan gum is a new polymer produced by the microbial fermentation of sugar raw materials. It has the unique property of forming a gel under heating conditions and is also called a thermal gel. In 1996, the FDA approved the application of curdlan gum as a food raw material in the food industry. Curdlan gum does not dissolve in water, but it will swell and disperse easily in cold water. It is heated to form a colloid, according to its properties can be divided into low colloid and high colloid. It is produced by heating to 55–65°C and cooling to about 40°C, forming low-grade colloid with thermal reflection. When the water dispersion of curdlan gum is heated to more than 80°C, a higher colloid is formed. The Curdlan gel can reduce the water loss rate of chilled meat products and extend their shelf life.

Keywords. Curdlan gum, Production source, Applications, Composition, Properties.

1. Agrobacterium

The *Agrobacterium* was discovered in 1853, and it is a group of bacteria that live naturally in the soil and was also found related to plants as well as mentioned by [1]. That in the year 1853 the first description was made of neoplastic diseases that affect different types of plants and the strains of some species of this bacteria infect the roots and stems of a large variety of dicotyledonous plants and some gymnosperms through wounds.

[2] The genus *Agrobacterium* is split into different species based on their pathogenicity, with *A. Radiobacter* (non-pathogenic), *Agrobacterium tumefaciens* (the cause of coronary gallbladder tumors), and *Agrobacterium rhizogenes* being the most well-known (the causative agent of the disease). *A. vitis* (the causative agent of coronary infection of grape roots).

The strains that cause coronary pathogens are classified as *A. tumefaciens*. They cause disease in dicotyledons. The disease causes the formation of tumor-like swellings and can generally be found on the crown of plants directly above the soil or in the roots. *Agrobacterium* causes coronary disease by transferring part From their DNA to plants [3].

2. Microbial Extracellular Polysaccharides

Polysaccharides are produced by plants, algae, fungi or bacteria, and they are bio-polymers of high molecular weight produced by microorganisms during the processes of demolition and building inside the cell. They are used industrially as thickeners, stabilizers and gelling agents in foodstuffs. It has also entered into the manufacture of biomedical materials that help heal wounds and adhesives used in surgical operations, antioxidant activity and biological antecedents are just two examples of how they might be useful in the body [4].



[5] stated that microbial polysaccharides are multiple substances produced outside the cell that have an important role in the metabolic processes of the microorganism. Polysaccharides can resist harsh conditions such as salinity and drought, and they are used in various industrial and agricultural applications, because they have unique properties such as vital activity, its ability to crystallize, as well as the presence of very large molecular weights that may reach several million.

Polysaccharides were produced from different sources such as plants, algae and fungi, but the increase in their production costs and their impact on seasonal climatic conditions and the increased demand for them due to their importance and applications in the fields of biomedicine, pharmaceuticals, nutrients and cosmetics, which made researchers in recent times to study, characterize and produce many microbial polysaccharides and study their properties. It is cheaper than other natural polysaccharides such as Arabic gum and Carrageenans produced by plants and seaweed, as well as the speed of their production compared to those produced from plants [4].

The Polysaccharides used commercially in the food industry are xanthan gum [6], gellan gum, carrageenan, alginate, pullulan, manan, galactomannan, hyaluronic acid and uronic acid [7].

Microbial polysaccharides are widely used in various fields to improve food quality and have an important role in health through their antimicrobial, antioxidant and cholesterol-lowering effect, as well as their antiviral activity. And some of them have electrical charges that help to stabilize the cells in the suspension evenly, so the cells remain diffuse without collecting or precipitating [8].




Figure 1. Homologous polysaccharides that can be produced by microorganisms Stewart-Tull, D E S (1980).

3. Curdlan Gum

Curdlan is a resinous substance resulting from various properties of microorganisms and the reason for its unique properties for its tolerance of high temperatures is called thermal gel. The basic building block of curdlan is glucose and the units are bound mediated by (1-3) β -glucose which is a linear, molecular homopolymer of $5.3 \times 10^4 - 2.0 \times 10^6$ Daltons. [9].

Many previous studies have shown that Curdlan is synthesized in the medium by the enzyme Uridine-diphosphate-glucose phosphorylase to form Uridine-diphosphate-glucose, as it is a pioneer to the biosynthesis of Curdlan. The presence of the carbon source in the growth medium is a factor that stimulates bacteria to secrete special enzymes in production, so the enzymes decompose sugars and convert them to curdlan [10].

3.1. A brief History of the Production of the Curdlan

Started the idea of producing Curdlan, which is one of the polysaccharides, after increasing interest in its use as a food additive, and it has several functional properties. Therefore, it is used as a stabilizer, supporter, and thickener. The researcher Harada was the first to discover it in 1966 and it was called Curdlan. Its capacity to "coagulate" in a water bath is where the term comes from [11].

In 1996, Curdlan (then known as Pure Glucan) received FDA approval for commercial usage. Curdlan's physical and chemical features set it apart from other polysaccharides like starch, garnering significant interest from the food and non-food industries. Cordlan is odorless, tasteless, and insoluble in water and alcohol but soluble in alkaline solutions, giving it excellent stability in industrial sterilization, frying, and freezing processes [12].

In 1996, Curdlan was granted registration with the FDA in the United States. It is approved for use in the European Union as a stabilizer under the symbol E424 and its formula $(C_6H_{10}O_5)_n$ [13].

3.2. Curdlan Production Source

The genus *Agrobacterium* is the most productive of linear curdlan, mainly (3-1)-Glucan, which may have a few chains, producing the branched sugar Glucan (1→3,1→6) [14]. It was discovered for the first time in *Alcaligenes faecalis* and was produced with other extracellular exogenous sugars [15]. The table below shows the most important microorganisms producing Curdlan sugar.

Table 1. The most important microorganisms producing curdlan gum.

Type of glucan	Bacterial Source	Sources
(1→3)-β-D-glucan,linear	<i>Agrobacterium</i> sp. 10C3 and derivatives	[16]
	<i>Agrobacterium</i> sp. ATCC 31749 and derivatives	[17]
	FO12607,12665,13127,13256 <i>A.radiobacter</i>	[16]
	<i>A. rhizogenes</i> IFO13259	[16]
	<i>Rhizobium trifolii</i> J60	[18]
	<i>Rhizobium</i> sp. TISTR 64B	[19]
	<i>Cellulomonas</i> spp.	[20]
(1→3,1→6)-β-Dglucan,cyclic	<i>C. flavigena</i> KU	[20]
	<i>Bradyrhizobium japonicum</i> USDA 110	[21]
	<i>R. loti</i> NZP 2309	[22]
	<i>Azorhizobium caulinodans</i> HAMBI 216	[23]
	<i>A. brasilense</i> ATCC29710	[24]
(1→3)-β-D-glucan,cyclic	<i>Bradyrhizobium japonicum</i> ndvC mutants	[25]
	<i>Sinorhizobium meliloti</i> ndvB mutant with the <i>B. japonicum</i> ndv locus	[25]
(1→3,1→2)-β-Dglucan, side-chainbranched	<i>Streptococcus pneumoniae</i> type 37	[26]

3.3. Curdlan Gum Composition

Curdlan is a homopolymer. The basic building unit is glucose. The units are related by a β-1,3-glucan glycosidic bond as shown in Figure 1. Some of its types contain β-(1→3, 1→2) glucose subunits. Their molecular weight is up to 5.3×10^4 - 2.0×10^6 dalton and is approved for use in the European Union as a stabilizer under the symbol E424 and formula $(C_6H_{10}O_5)_n$ [27].




Figure 2. Curdlan Structure Produced by Bacteria. [28]

Curdlan consists of a carbohydrate concentration of at least 90% and a maximum water content of 10%. Curdlan lacks acidic components despite the abundance of glucose subunits linked by beta bonds between the ring's first and third carbon atoms [28]. The D-glucose that curdlan is made of is similar to cellulose, however the bonds are different. Curdlan is insoluble in water although it can be dissolved in aqueous sodium hydroxide due to the ionization of hydrogen bonds. When hydrated NaOH is introduced, the bonds break due to swelling and the granular structure of Curdlan is lost [29].

3.4. Curdlan Gum Properties

Curdlan is characterized by many important properties. It is a resinous substance characterized by its tolerance to high temperatures and the behavior of crystallization at high temperature. It is colorless, tasteless, and odorless. It is insoluble in water, alcohol, or acidic solutions, but it is soluble in alkaline solutions (pH12), as sodium hydroxide and trisodium phosphate. Although it is insoluble in water, it is able to form gels after heating, and it is a biodegradable, highly polar, non-toxic polymer. It can be easily decomposed into carbon dioxide and water. It can be used as an edible film which has good elongation when broken and the ability to form heat-resistant gel at the same time. It is considered a dietary fiber and characterized by its ability to form hard and flexible gels that cannot be thermally reversed when heated in aqueous suspensions at temperatures of 80 ° C or higher. 130 ° C and is not affected by freezing and thawing processes [11,30].

It has the ability to absorb water, bind it and prevent separation, so it is used in many food industries as a support material, such as the cream and water ice cream industry, the meat industry of all kinds, and the pasta industry. Curdlan was used as a catalyst for prebiotic bacteria in therapeutic food and beverages, as it showed great stability during industrial processes such as frying and freezing and solubility [31].

[32] Describe how curdlan can soak up water at a rate greater than 100 times its own weight. At 60 degrees Celsius, a low-density, thermally reversible gel forms, while at 80 degrees Celsius, a flexible, high-density gel is produced.

3.5. Curdlan Gum Synthesis

By sequencing the genome of ATCC 31749, we were able to determine regulatory mechanisms for curdlan biosynthesis and localize the genes involved in this process to two distinct genomic locations. In one location, you'll find the *crdR* gene, which encodes a positive transcriptional regulatory protein, and two other genes, *crdS* (which is homologous to glycosyltransferases like cellulose synthase), *crdA* (whose function is unclear), and *crdC* (whose function is also unclear) [33,34].

Curdlan gum synthesis requires the presence of the enzyme Hexokinase to form glucose-6-phosphate via glucose phosphorylation. The process takes place inside the cell after glucose enters the cytoplasm by active transport, UDP-glucose phosphorylase synthesizes UDP glucose from uridine triphosphate (UTP) as the initiator of Curdlan biosynthesis. A polymerization reaction catalyzed by a polymerase enzyme leads to the formation of Glucose bonds and the release of the polymer into the extracellular

environment after chain elongation, after which the transferase binds glucose to a fat molecule, isoprenoid fat phosphate, releasing UDP. Thereafter, UTP is synthesized from UDP by UDP kinase, and the curdian production cycle is resumed [35].




Figure 3. The metabolic pathway for the production of Curdian from bacteria [36]

4. Optimal Conditions for the Production of Curdian Gum

The study development conditions of the microorganisms producing Curdian were studied in order to find the best conditions that lead to an increase in the quantities of Curdian produced and thus reduce production costs. It was found that the processes of producing curdian gum from microorganisms are affected by many factors such as carbon source, temperature, ventilation, pH, type of fermentation, and the size of the bacterial inoculum [37,38].

The best conditions for the production of curdian gum are from *Agrobacterium sp.* ATCC was using different sources of carbon and nitrogen when using sucrose as a carbon source by 140 g/L, was obtained 32 g/L curdian, and when using a nitrogen source, Using urea (28.16 and 9.58 g/L), ATCC 31749 was obtained, while using NH_4Cl resulted in (15.17 and 6.25 g/L) and incubation time of (95 hours) [39]. [40], found that the best optimum conditions for the production of curdian gum from *Agrobacterium sp.* are the use of sucrose 140 g / l, NH_4Cl 2.4 g / l, KH_2PO_4 1.0 g / l and $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.5 g / l at a temperature of is 30 °C, the optimal pH is 7, the incubation period is 96 hours, and by using the vibrating incubator at a speed of 150 rpm, the highest production rate of curdian gum reached 5.2 g / liter.

Researchers [41] investigated the most effective carbon sources for curdian formation by *Agrobacterium sp.*, which include sugars like glucose, sucrose, and maltose. A specific group of the three sources were examined from 5 to 25% to obtain maximum yield The results showed 38.1 g/L at 15% sucrose, 27.3 g/L at 20% glucose and 37.4 g/L at 20% maltose, and the maximum yield of Curdian was obtained using sucrose under limited nitrogen concentration. [11], indicated that Curdian can be produced from bacteria using soluble condensed corn syrup as a source of carbon and nitrogen. As the soluble condensed corn juice material concentration rose from 50 g/L to 400 g/L, the Curdian concentration rose as well, though not in direct proportion. After 120 hours, the strain produced the maximum concentration of Curdian when fed 400 g/L soluble condensed corn juice. Its concentration exceeds that seen in curdian formation that relies on glucose. Similarly, ATCC 31749's biomass production was greater after 120 hours of culture on 400 g/L soluble condensed corn juice than it was after growing on glucose.

[42] studied the effect of nitrogen sources on curd production, potassium nitrate, sodium nitrate, ammonium nitrate, ammonium acetate, ammonium citrate, ammonium chloride, ammonium sulfate, and ammonium acetate. Based on the observations, a maximum achieved was yield of 4.8 g/L Curd. They [43] indicated that the optimum acidity function for the production of Curd is 5.5 and to obtain an increase in production, it is the optimum nutrition for microorganisms to obtain 60 g/L Curd. [33], was able to obtain a maximum Curd yield of 47.97 ± 0.57 g/L from *Agrobacterium sp. ATCC 31749* using optimized medium containing 60 g/L sucrose, 6 g/L yeast, 2 g/L KH_2PO_4 , 0.4 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/L CaCO_3 , 0.1 g/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.04 g/L MnSO_4 , and 0.02 g/L ZnCl_2 at 30 °C and 280 rpm after 96 hours of fermentation.

5. The Use of Alternatives in the Production of Curd Gum

Curd gum and other biopolymers have gained a lot of attention in recent years due to their useful physical and chemical properties and wide range of applications. The chemical, culinary, petroleum, medical, and biotech industries are just a few of the many that can benefit from microbial sugars. Many researchers have resorted to using agricultural waste as an alternative carbon source for sucrose, thus reducing production costs. Sugars constitute only a small part of the current polymer market due to their high cost production and recovery. For this reason a lot of effort has been put into developing cost-effective production processes using cheaper fermentation substrates [44].

Found [41] that the effect of different carbon sources on biomass production and Curd was distinguished by using sucrose among other sugar sources (glucose, fructose, maltose, xylose, mannose) at the pH level 7 after 72 hours of a period of time. Fermentation at 30 °C on a rocking incubator at a shaking rate of 100 rpm with 50 mL of modified fermentation medium. Explain [45], the use of sugar beet molasse in the production of curd gum from *Agrobacterium sp.* It gave good yield when incubating for 120 hours using the incubator at a speed of 150 rpm at a temperature of 30°C and the pH function was 7, and the yield was 42 g/liter of molasses as a carbon source instead of sucrose.

6. Curd Gum Purification

Biotechnology is divided, depending on the nature of the microorganism, into two main groups, the first of which is easy to separate, which is the biological products outside the cell, compared with the second group whose products are inside the cell and requires a number of steps to obtain them in pure form [46]. [47], the possibility of purifying the Curd produced from *Agrobacterium* by adding one ml of the bacterial medium with two ml of sodium hydroxide (1N), the mixture was kept for one hour at room temperature, then the mixture was expelled at 4000 rpm to remove the bacterial cells, then mixed. The filtrate was mixed with hydrogen chloride (3N) until the pH reached 5-7 and the mixture was kept at 2-8 m overnight. To obtain the curd, a centrifuge was carried out at 4000 rpm. In order to get rid of the salts, the residues were washed with distilled water several times and then washed using acetone. To provide a dry product and store at room temperature until use.

[48] showed that the *agrobacterium*-derived curd was purified by the following steps: dissolving the curd in 5 ml of fermentation medium containing 100 ml of sodium hydroxide 0.5 M at 60 °C for one hour; centrifuging the mixture at 10,000 g for 10 minutes; adding 100 mL of hydrogen chloride 1 M to the supernatant; centrifuging the suspension for 30 minutes; washing three times with distilled water.

7. Description and Diagnosis of Curd Gum

The product obtained during fermentation processes must know its physical and chemical composition in order to characterize the compound, so some researchers used modern characterization techniques that determine the physical and chemical properties, molecular mass, molecular structure, shape, thermal and mechanical properties and some other chemical properties, some researchers do not consider characterization and diagnosis results. Multiple chemical assays and biomolecules are clear [49].

8. Curdlan Gum Applications

There is currently a growing interest in microbial sugars as an important source of new biopolymers for industrial use. One of the industrially important microbial sugars is curdlan, which has many uses due to its physical and chemical properties that make it important in food, industrial and pharmaceutical applications. It is safe and non-toxic and is used industrially as thickeners, stabilizers and agents. Crystallization in foodstuffs is also used for its biological functions, such as its antioxidant and prebiotic activity. [50].

Dissect this [40] As -glucan is consumed as an edible fiber, curdlan has been utilized as a stabilizer in jelly foods, a bio-thickener for pasta, and in other applications. Its unusual qualities [tasteless, odorless, and colorless] and its capacity to contain water make it useful as a textile material in the meat, dairy, and baking sectors.

Explain [51] the possibility of using curdlan gum in foods as a thickener, stabilizer, flavor enhancer, and in other therapeutic materials. On the structure of milk, which encouraged a decrease in the synergy of milk compared with milk without curdlan, the curdlan which also contributed to a decrease in the growth of lactic acid bacteria during the storage period, which and helps the gradual conversion of lactose to lactic acid by the initiator.

As shown in [52], curdlan sulfate has potential as both a standalone immunotherapeutic agent and as an adjuvant to an antiviral vaccination for the treatment of hepatitis infection. Curdlan, when combined with activated carbon, can be used to purge heavy metals from herbs used in the creation of oriental medicine, making it a valuable industrial use.

[33] studied that curdlan was used to determine the quality of pasta and texture frequently, as a strong gluten network tended to enhance the firmness and malleability of the pasta, and that the preparation of pasta with curdlan improved the eating quality of both raw and cooked pasta, which showed a gelatinous strength. High resulting in excellent edible quality noodles. Also, the tensile properties of the noodles changed, the rupture strength and distance, and the percentage of breaking of curdlan-containing noodles was much higher than that of the no-curdlan pasta which led to a significant improvement in the eating quality of raw and cooked pasta.

Conclusions

Curdlan has been studied for over fifty years due to its distinct gel formation, poor water solubility, mild processing conditions, indigestibility, and process ability. Researchers have only recently begun to investigate its applications in the food, health, and biomedical sectors. It has a nutritional benefit because it has several functional properties, so it is used as a stabilizer, fortifier, and thickener, and it has health benefits from its prebiotic use. It can be introduced into many food industries. Agricultural waste was used as a carbon source, such as date juice, molasses, grape juice, and whey protein, to reduce the economic cost of curdlan production. In addition to the disposal of agricultural waste that causes environmental damage, The yield of curdlan varies according to the type of strain and the culture media used.

References

- [1] Pacurar, D.I.; Thordal-Christensen, H.; Pacurar, M.L.; Pamfil, D.; Botez, C.; Bellini, C.(2011). *Agrobacterium tumefaciens: From Crown Gall Tumors to Genetic Transformation*. *Physiol. Mol. Plant Pathol.* 76, 76–81. <https://doi.org/10.1016/j.pmpp.2011.06.004>.
- [2] Flores-Félix, José David; Menéndez, Esther; Peix, Alvaro; García-Fraile, Paula; Velázquez, Encarna (2019). History and current taxonomic status of genus *Agrobacterium*. *Systematic and Applied Microbiology*, , 126046–. S0723- (19)30341-8. <https://doi.org/10.1016/j.syapm.2019.126046>.
- [3] Portier, P. ; Saux, M. F. ; Mougél, C. ; Lerondelle, C. ; Chapulliot, D. ; Thioulouse, J. ; and Nesme, X. (2006). Identification of genomic species in *Agrobacterium* biovar 1 by AFLP genomic markers. *Appl. Environ Microbiol.* 72(11):7123–7131. <https://doi.org/10.1128//AEM.00018-06>.
- [4] Ahmad Qamar, Sarmad; Asgher, Muhammad; Bilal, Muhammad; Iqbal, Hafiz M.N. (2020). Bio-based active food packaging materials: Sustainable alternative to conventional petrochemical-based packaging materials. *Food Research International*, (1370, 109625. <https://doi.org/10.16/j.foodres.2020.109625>.
- [5] Muhammad Asgher;Aneesa Rani;Nimrah Khalid;Sarmad Ahmad Qamar;Muhammad Bilal; (2021). Bioconversion of sugarcane molasses waste to high-value exopolysaccharides by engineered

- Bacillus licheniformis*. Case Studies in Chemical and Environmental Engineering, Vol.3. 100084 <https://doi.org/10.16/j.cscee.2021.100084>.
- [6] Al-Roomi, F. W., & Al-Sahlany, S. T. G. (2022). Identification and Characterization of Xanthan Gum Produced from Date Juice by a Local Isolate of Bacteria *Xanthomonas campestris*. *Basrah Journal of Agricultural Sciences*, 35[1], 35–49. <https://doi.org/10.37077/25200860.2022.35.1.03>.
 - [7] Karataş, Meltem Altin; Gökkaya, Hasan (2018). A review on machinability of carbon fiber reinforced polymer [CFRP] and glass fiber reinforced polymer (GFRP) composite materials. *Defence Technology*, 14 (4), 318–326. <https://doi.org/10.1016/j.dt.2018.02.001>.
 - [8] Nakanishi I, Kimura K, Suzuki T, Ishikawa M, Banno I, Sakane T, Harada T (1976) Demonstration of curdlan-type polysaccharide and some other β -1,3-glucan in microorganisms with aniline blue. *J. Gen Appl. Microbiol.* 22:1–11. <https://doi.org/10.2323/jgam.22.1>.
 - [9] Ortiz Martinez C, Pereira Ruiz S, Carvalho Fenelon V, Rodrigues de Moraes G, Luciano Baesso M, Matioli G (2016) Characterization of curdlan produced by *Agrobacterium* sp. IFO 13140 cells immobilized in a loofa sponge matrix, and application of this biopolymer in the development of functional yogurt. *J Sci Food Agric* 96(7):2410–2417.
 - [10] Yuan, M., Fu, G., Sun, Y., & Zhang, D. [2021]. Biosynthesis and applications of curdlan. *Carbohydrate Polymers*, 273, 118597. <https://doi.org/10.1016/j.carbpol.2021.118597>.
 - [11] West TP (2020) Production of the polysaccharide curdlan by *Agrobacterium* species on processing coproducts and plant lignocellulosic hydrolysates. *Fermentation* 6:16. <https://doi.org/10.3390/fermentation6010016>.
 - [12] Kumar Verma, Deepak; Kareem Niamah, Alaa; Patel, Ami R.; Thakur, Mamta; Singh Sandhu, Kawaljit; ChÃ¡vez-GonzÃ¡lez, MÃ³nica L.; Shah, Nihir; Noe Aguilar, Cristobal (2020). Chemistry and Microbial Sources of Curdlan with Potential Application and Safety Regulations as Prebiotic in Food and Health. *Food Research International*, Vol. (133), 109136.
 - [13] Suflet DM, Popescu I, Prisacaru AI, Pelin IM (2020) Synthesis and characterization of curdlan—phosphorylated curdlan based hydrogels for drug release. *Int. J. Polym. Mater* 70(12):870–879. <https://doi.org/10.1080/00914037.2020.1765360>.
 - [14] Venkatachalam, Geetha; Arumugam, Senthilkumar; Doble, Mukesh (2020). Industrial production and applications of linear and branched glucans. *Indian Chemical Engineer*, 1–15. <https://doi.org/10.80/00194506.2020.1798820>.
 - [15] Prakash, S.; Rajeswari, K.; Divya, P.; Ferlin, M.; Rajeshwari, C. T.; Vanavil, B. (2018). Optimization and production of curdlan gum using *Bacillus cereus* pr3 isolated from rhizosphere of leguminous plant. <https://doi.org/10.1080/10826068.2018.1451886>.
 - [16] Nakanishi I, Kimura K, Suzuki T, Ishikawa M, Banno I, Sakane T, Harada T (1976) Demonstration of curdlan-type polysaccharide and some other β -1,3-glucan in microorganisms with aniline blue. *J Gen Appl Microbiol* 22(1):1–11.
 - [17] Kim, M.K.; K.E. Ryn, W.A. Choi, Y.H. Rhee, I.Y. Lee, (2003). Enhanced production of (1→3)- β -d-glucan by mutant strain of *Agrobacterium* species, *Biochem. Eng. J.* 16(2), 163–168. [https://doi.org/10.16/S1369-703X\(03\)00032-9](https://doi.org/10.16/S1369-703X(03)00032-9).
 - [18] Ghai SK, Hisamatsu A, Amemura A, Harada T (1981) Production and chemical composition of extracellular polysaccharides of *Rhizobium*. *J Gen Microbiol* 122:33–40. <https://doi.org/10.1099/00221287-122-1-33>.
 - [19] Footrakul K, Suyanandana P, Amemura A, Harada T (1981) Extracellular polysaccharides of *Rhizobium* from Bangkok MIRCEN collection. *J. Ferment Technol* 59:9–14. <https://doi.org/10.1128/jb.158.3.1168-1171.1984>
 - [20] Kenyon WJ, and Buller C.S. (2002) Structural analysis of the curdlan-like exopolysaccharide produced by *Cellulomonas flavigena* KU. *J. Ind. Microbiol. Biotechnol.* 29(4):200–203. <https://doi.org/10.1038/sj.jim.7000277>.
 - [21] Miller KJ, Gore RS, Johnson R, Benesi AJ, Reinhold VN (1990) Cell-associated oligosaccharides of *Bradyrhizobium* spp. *J Bacteriol* 172(1):136–142. <https://doi.org/10.1128/jb.172.1.136142.1990>.
 - [22] Estrella MJ, Pfeffer PE, Brouillette JN, Ugalde RA, Iannino NI de (2000) Biosynthesis and structure of cell associated glucans in slow growing *Rhizobium loti* strain NVP2309. *Symbiosis* 29:173–19.
 - [23] Komaniecka I, and Choma A (2003) Isolation and characterisation of periplasmic cyclic β -glucans of *Azorhizobium caulinodans*. *FEMS Microbiol. Lett* Vol.227(2):263–269. [https://doi.org/10.16/S0378-1097\(03\)00690-6](https://doi.org/10.16/S0378-1097(03)00690-6).
 - [24] Altabe SG, Talaga P, Wieruszkeski J-M, Lippens G, Ugalde R, Bohin J-P (1998) Periplasmic glucans of *Azospirillum brasilense*. In Elmerich C, Kondorosi A, Newton WE (eds) *Biological nitrogen fixation for the 21st century*. Kluwer, Dordrecht, p. 390. <https://doi.org/10.5483/BMBRep.2009.42.12.769>.

- [25] Joanne M. Holden; Alison L. Eldridge; Gary R. Beecher; I. Marilyn Buzzard; Seema Bhagwat; Carol S. Davis; Larry W. Douglass; Susan Gebhardt; David Haytowitz; Sally Schakel (1999). Carotenoid Content of U.S. Foods: An Update of the Database. *Journal of Food Composition and Analysis* , 12(3), 169–196. <https://doi.org/10.1006/jfca.1999.0827>.
- [26] Llull D, García E, and López R (2001) Tts, a processive β -glucosyltransferase of *Streptococcus pneumoniae*, directs the synthesis of branched Type 37 capsular polysaccharide in pneumococcus and other Gram-positive species. *J. Biol. Chem.* 276:21053–21061. <https://doi.org/10.1074/jbc.M010287200>.
- [27] Nakashima, Y., K. Fukasawa, and H. Samejima. (2018). Estimating animal density without individual recognition using information derivable exclusively from camera traps. *Journal of Applied Ecology* 55: 735–744. <https://doi.org/10.1111/1365-2664.13059>.
- [28] Ying Liang; Zhuoting Qu; Mei Liu; Mengfei Zhu; Xia Zhang; Le Wang; Feng Jia; Xiaobei Zhan; Jinshui Wang; (2021). Further interpretation of the strengthening effect of curdlan on frozen cooked noodles quality during frozen storage: Studies on water state and properties . *Food Chemistry*. Vol. 346(1). 128908. <https://doi.org/10.1016/j.foodchem.2020.128908>.
- [29] Jia, Xiaoyu; Wang, Cheng; Du, Xueqing; Peng, Hui; Liu, Lin; Xiao, Yazhong; He, Chao (2021). Specific hydrolysis of curdlan with a novel glycoside hydrolase family 128 β -1,3-endoglucanase containing a carbohydrate-binding module. *Carbohydrate Polymers*, 253(1), 117276. <https://doi.org/10.1016/j.carbpol.2020.117276>.
- [30] Dominguez-Martinez, B. M., Martínez-Flores, H. E., Berrios, J. D. J., Otoni, C. G., Wood, D. F., & Velazquez, G. (2016). Physical Characterization of Biodegradable Films Based on Chitosan, Polyvinyl Alcohol and Opuntia Mucilage. *Journal of Polymers and the Environment*, 25(3), 683–691. <https://doi.org/10.1007/s10924-016-0851-y>.
- [31] Zhai, W., & Iwata, T. (2019). Synthesis and properties of curdlan branched and linear mixed ester derivatives. *Polymer Degradation and Stability*, 161, 56–50. <https://doi.org/10.1016/j.polymdegradstab.2019.01.008>.
- [32] Zhu, Jingyi; Wang, Qian; Han, Lu; Zhang, Chong; Wang, Yuanyuan; Tu, Kang; Peng, Jing; Wang, Jiahong; Pan, Lei (2021). Effects of caprolactam content on curdlan-based food packaging film and detection by infrared spectroscopy. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 245(5), 118942. <https://doi.org/10.1016/j.saa.2020.118942>.
- [33] Gao, H., Zhang, W., Zhang, J., Huang, Y., Zhang, J., Tian, J., ... Jiang, D. (2021). Methionine biosynthesis pathway genes affect curdlan biosynthesis of *Agrobacterium* sp. CGMCC 11546 via energy regeneration. *International Journal of Biological Macromolecules*, 185, 821–831. <https://doi.org/10.1016/j.ijbiomac.2021.06.189>.
- [34] Yu, X.; Zhang, C.; Yang, L.; Zhao, L.; Lin, C.; Liu, Z.; Mao, Z. (2015). CdR function in a curdlan-producing *Agrobacterium* sp. ATCC31749 strain. *BMC Microbiol*, 15, 25. <https://doi.org/10.1186/s12866-015-0356-1>.
- [35] Jianrong Wu; Ruoyu Yang; Minjie Gao; Hongtao Zhang; Xiaobei Zhan; (2021). Synthesis of functional oligosaccharides and their derivatives through cocultivation and cellular NTP regeneration . *Advances in Applied Microbiology*, (115), <https://doi.org/10.1016/bs.aambs.2021.02.002>.
- [36] Lee, J. H., & Park, Y. H. (2001). Optimal production of curdlan by *Agrobacterium* sp. with feedback inferential control of optimal pH profile. *Biotechnology Letters*, 23(7), 525–530. <https://doi.org/10.1023/A:1010374519891>.
- [37] Jin, L.-H.; Um, H.-J.; Yin, C.-J.; Kim, Y.-H.; Lee, J.-H. (2008) Proteomic analysis of curdlan-producing *Agrobacterium* sp. in response to pH downshift. *J. Biotechnol.*, 138, 80–87. <https://doi.org/10.1016/j.jbiotec.2008.08.010>.
- [38] Minjie Gao; Guoshuai Yang; Feifei Li; Zichao Wang; Xiuyu Hu; Yun Jiang; Jiajun Yan; Zhitao Li; Xiaobei Zhan; (2021). Efficient endo- β -1,3-glucanase expression in *Pichia pastoris* for co-culture with *Agrobacterium* sp. for direct curdlan oligosaccharide production . *International Journal of Biological Macromolecules*, Vol. 182(1), 1611–1617 . <https://doi.org/10.1016/j.ijbiomac.2021.05.142>.
- [39] Jiang, L. (2013). Effect of nitrogen source on curdlan production by *Alcaligenes faecalis* ATCC 31749. *International Journal of Biological Macromolecules*, 52, 218–220. <https://doi.org/10.1016/j.ijbiomac.2012.10.010>.
- [40] Shih IngLung ; Yu JinYee ; Hsieh Chienyan ; Wu JaneYii. (2009) Production and characterization of curdlan by *Agrobacterium* sp. *Biochemical Engineering Journal*. 43. 33–40. <https://doi.org/10.1016/j.bej.2008.08.006>.

- [41] Wan, Jie; Wang, Yifeng; Jiang, Deming; Gao, Hongliang; Yang, Guang; Yang, Xuexia (2020). Effects of carbon sources on production and properties of curdlan using *Agrobacterium* sp. DH-2. *Preparative Biochemistry & Biotechnology*, 1–8. <https://doi.org/10.1080/10826068.2020.1777423>.
- [42] Xia, Z. (2013). Effect of Tween 80 on the production of curdlan by *Alcaligenes faecalis* ATCC 31749. *Carbohydr. Polym.* 98, 178–180. <https://doi.org/10.1016/j.carbpol.2013.05.073>.
- [43] Shobana S. Ponnusami V and Sugumaran K. R. (2015) Documentation on current developments in production and applications of a β -(1-3)-D glucan (Curdlan). *J. Chem. Pharm. Res.* 7(2):807-810. www.jocpr.com.
- [44] Srikanth, R.; Reddy, C.H.; Siddartha, G.; Ramaiah, M.J. & Uppuluri, K.B. (2015). Review on production, characterization and applications of microbial levan. *Carbohydr. Polym.*, 120: 102-114. <https://doi.org/10.1016/j.carbpol.2014.12.003>.
- [45] Lee, I-Y; W T Seo; G J Kim; M K Kim; C S Park; Y H Park (1997). Production of curdlan using sucrose or sugar cane molasses by two-step fed-batch cultivation of *Agrobacterium* species. *Journal of Industrial Microbiology and Biotechnology* , 18(4), 255–259. <https://doi.org/10.1038/sj.jim.2900378>.
- [46] Al-Khafaji, Z.M. (2008) Probiotics (For Life). Published and Printed by House and Documentation Baghdad. Library of Control No. 84, 3-11. <https://doi.org/10.4236/health.2013.512A007>.
- [47] El-Sehemy I.A., Haroun B.M., El-Diwanly A.I., El-Shahed K.Y., Awad H.M., and Tolba I.H.(2012). isolation and identification of native Egyptian curdlan production *agrobacterium* isolate. *Az. J. Pharm Sci.* Vol. 45, March, 2012. DOI: 10.21608/ajps.2012.7250
- [48] Liu, J., Tang, J., Li, X., Yan, Q., Ma, J., & Jiang, Z. (2019). Curdlan (*Alcaligenes faecalis*) (1 $\rightarrow$ 3)- β -d-glucan oligosaccharides Drive M1 phenotype polarization in murine bone marrowderived macrophages via activation of MAPKs and NF- κ B pathways. *Molecules*, 24 (23), 4251. <https://doi.org/10.3390/molecules24234251>.
- [49] Khassaf, W. H. ., Niamah, A. K. ., & Al-Manhel, A. J. A. . (2019). Study of the Optimal Conditions of Levan Production from a Local Isolate of *Bacillus subtilis* subsp. *subtilis* w36. *Basrah Journal of Agricultural Sciences*, 32(2), <https://doi.org/10.37077/25200860.2019.211>.
- [50] Mangolim, C. S., da Silva, T. T., Fenelon, V. C., Koga, L. N., de Souza Ferreira, S. B., Bruschi, M. L., & Matioli, G. (2017). Description of recovery method used for curdlan produced by *Agrobacterium* sp. IFO 13140 and its relation to the morphology and physicochemical and technological properties of the polysaccharide. *PloS One*, 12 (2), e0171469. <https://doi.org/10.1371/journal.pone.0171469>.
- [51] Wei, Yuanyuan; Guo, Yalong; Li, Ruiqi; Ma, Aiqin; Zhang, Hongbin (2020). Rheological characterization of polysaccharide thickeners oriented for dysphagia management: Carboxymethylated curdlan, konjac glucomannan and their mixtures compared to xanthan gum. *Food Hydrocolloids*, (110), 106198. <https://doi.org/10.1016/j.foodhyd.2020.106198>.
- [52] Shobana S. Ponnusami V and Sugumaran K. R. (2015) Documentation on current developments in production and applications of a β -(1-3)-D glucan (Curdlan). *J. Chem. Pharm. Res.* 7(2):807-810. www.jocpr.com