


Chapter 6

Carboxylic Acids

1. Structure

Carboxylic acids are organic compounds containing the carboxyl group (-COOH), where in the hydroxyl group (-OH) is directly attached to the carbonyl (C=O) group.



Nomenclature of Carboxylic Acids

The common names of some basic carboxylic acids are derived from Latin names that indicate the first original natural source of the carboxylic acid.

Structure of Acid	Natural Source	Common Name
$\text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$	Ants (Formica)	Formic acid
$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$	Vinegar (Acetum)	Acetic acid
$\text{CH}_3\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$	Basic Fat (Propio)	Propionic acid
$\text{CH}_3\text{CH}_2\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$	Rancid butter (Butyrum)	Butyric acid
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$	Present in a Valerian herb	Valeric acid
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$	Goat (Caper)	Caproic acid

Common Names of Carboxylic Acids

The common name of a carboxylic acid (R-COOH) is derived by adding the suffix **-ic acid** to a prefix representing the chain length of the carboxylic acid.

# of Carbons	Prefix	Common Name of Acid
1	Form-	Formic acid
2	Acet-	Acetic acid
3	Propion-	Propionic acid
4	Butyr-	Butyric acid
5	Valer-	Valeric acid
6	Capro-	Caproic acid
Aromatic acid	Benzo-	Benzoic acid

IUPAC Nomenclature of Aliphatic Carboxylic Acids

IUPAC names of straight chain aliphatic carboxylic acids are derived by adding the suffix **-oic acid** to the systematic name of the parent hydrocarbon. They are named as alkanonic acids.

# of Carbons	Structure & IUPAC Name of Alkane	Structure & IUPAC Name of Acid
1	$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array}$ Methane	$\begin{array}{c} \text{O} \\ \\ \text{H}-\text{C}-\text{OH} \end{array}$ Methanoic acid
2	CH_3-CH_3 Ethane	$\begin{array}{c} \text{O} \\ \\ \text{CH}_3-\text{C}-\text{OH} \end{array}$ Ethanoic acid
3	$\text{CH}_3\text{CH}_2-\text{CH}_3$ Propane	$\begin{array}{c} \text{O} \\ \\ \text{CH}_3\text{CH}_2-\text{C}-\text{OH} \end{array}$ Propanoic acid



Systematic Nomenclature of Substituted Carboxylic Acids

The systematic names of substituted aliphatic carboxylic acids are derived by:

- (i) First identifying the parent chain that contains most, if not all, the carboxyl groups.
- (ii) Number the parent chain from the carbon of the carboxyl group i.e the carboxyl carbon is C-1.
- (iii) Identify the substituents and assign each substituent a locator/address number (2,3,4...etc.) consistent with the numbering in the parent chain.
- (iv) Arrange the names of the substituents in alphabetical order in the systematic name of the poly-substituted carboxylic acid.

Physical properties of Carboxylic Acids

The physical properties of carboxylic acids can be explained from the perspective of the bond polarization in the carboxyl group and its capacity to engage in hydrogen-bonding. Carboxylic acids boil at considerably higher temperatures than alcohols, ketones, or aldehydes of similar molecular weight.




Strong intermolecular attractive forces hold the acid molecules together

The high boiling point of carboxylic acids is attributed to their capacity to readily form stable, hydrogen-bonded dimers.


Preparation of Carboxylic Acids

1. Oxidation of Primary Alcohols


The synthesis of carboxylic acids requires the generation or incorporation of the carboxyl group in a substrate.




The best conditions for the oxidation of primary alcohols to carboxylic acids is under the basic conditions employing potassium permanganate.



Example



2. Oxidation of Aldehydes



Aldehydes can be oxidized to carboxylic acids by a variety of oxidizing agents. Both strong and mild oxidizing agents may be employed successfully.



Strong Oxidizing Agents

- (a) CrO_3
Oxidations in water or dilute mineral acid
- (b) $\text{Na}_2\text{Cr}_2\text{O}_7$ or $\text{K}_2\text{Cr}_2\text{O}_7$
Oxidation in dilute mineral acid
- (c) KMnO_4
Oxidations in basic media in the presence of KOH


Mild Oxidizing Agents

- (a) Ag_2O

Note that Ag_2O does not oxidize alcohols





3. Carboxylation of Grignard Reagents




4. Hydrolysis of Nitriles

Acid-catalyzed hydrolysis of nitriles provides carboxylic acids.






Example



Reactions of Carboxylic Acids

The reactions of carboxylic acids can be directed to various sites on the carboxyl group.




Reactions of carboxylic acids can be placed into four categories:


- (1) Reactions at the acidic hydrogen on the carboxyl group.
- (2) Reactions at the carbonyl group
- (3) Reactions at the carboxylate oxygen
- (4) Reactions that lead to loss of the carboxyl group as CO_2

a) Reaction of Carboxylic Acids with Sodium Bicarbonate


The most reliable test for carboxylic acids employs NaHCO_3 leading to evolution of CO_2 . This is commonly called the bicarbonate test for carboxylic acids.



Example




b) Reactions of Carboxylic Acids with Strong Bases




Bases such as metal hydroxides (NaOH and KOH) and amines abstract the acidic proton on carboxylic acids to form carboxylate salts.

Mechanism




Example




c) Acid-Catalyzed Esterification of Carboxylic Acids

The traditional method for converting carboxylic acids to esters is through an acid-catalyzed esterification in the presence of an alcohol: Commonly referred to as the Fischer esterification.




The acid-catalyst can be provided by strong mineral acids such as H_2SO_4 , HCl and H_3PO_4 or organic acids such as benzenesulphonic acid or *p*-toluenesulphonic acid.

Example




d) Reaction of Carboxylic Acids with Amines


Amines, being organic bases, react with carboxylic acids to form ammonium salts.



Mechanism




Example




e) Reduction of Carboxylic Acids to Primary Alcohols

Carboxylic acids are reduced to primary alcohols when treated with a strong reducing agent such as LiAlH_4 .



Example



Medium strength reducing agents such as sodium borohydride (NaBH_4) that reduce aldehydes and ketones are not sufficiently strong to reduce carboxylic acids.

